

 <p><b>U.S. ENVIRONMENTAL PROTECTION AGENCY</b> Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460</p>	<p><b>EPA Registration Number:</b> 42750-208</p>	<p><b>Date of Issuance:</b> DEC 22 2009</p>
	<p><b>NOTICE OF PESTICIDE:</b> <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)</p>	
	<p><b>Term of Issuance:</b> Conditional</p>	
<p><b>Name of Pesticide Product:</b> Metalaxyl 28.35% FL ST</p>		
<p><b>Name and Address of Registrant (include ZIP Code):</b>  Albaugh Inc 1525 NE 36th ST Ankeny, Iowa 50021</p>		
<p><small>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</small></p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3 (c) (7) (A) provided that you:</p> <ol style="list-style-type: none"> <li>1. Submit/cite all data required for the registration/ reregistration of your product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.</li> <li>2. Make the following label changes:</li> </ol> <p>(a) <b>Page 1</b> – Change the EPA Registration Number to 42750-208</p>		
<p><b>Signature of Approving Official:</b>  <i>Mary L. Waller</i> Mary L. Waller, Product Manager 21 Fungicide Branch Registration Division (7505P)</p>		<p><b>Date:</b>  12/22/2009</p>

**(b) Page 1– Company Address:**

Under “Albaugh, Inc.” add the line: “1525 NE 36th ST”

**(c) Page 2– Personal Protective Equipment –PPE:**

Change the third bullet under “Applicators and other handlers” to read:

“Chemical-resistant gloves (such as nitrile ≥ 14 mils in thickness, neoprene ≥ 14 mils in thickness, Viton ≥ 14 mils in thickness and/or barrier laminate).”

**(d) Page 2– User Safety Recommendations :**

In the first sentence of the third bullet add a period and a space after “. . . this product”

**(e) Page 3– Directions for Use - Storage and Disposal – Container Disposal :**

Delete the text “(non-refillable > 5 gallons): Triple rinse as follows . . . after the flow begins to drip.”

Put a box around the entire “Storage and Disposal” section and move it so that it immediately follows the Agricultural Use Requirements” box.

**(f) Page 4– use with other registered fungicides and insecticides:**

In the sentence “Metalaxyl 28.25% FL ST may be applied as a water-based slurry . . . commercial seed treatment equipment.” remove the quotation mark at the beginning of the word fungicides.

Immediately following that sentence add the sentences: “Follow more restrictive labeling requirements of tank mix products. Do not tank mix with any product that contains a prohibition on tank mixing.”

**(g) Page 4– Section beginning “Note: Seed treated commercially . . .” :**

Immediately following the end of the first sentence, “Seed treated commercially . . . before being sold.” start a new paragraph and add the heading “Commercially treated seed must be labeled:”

Place the entire text of this paragraph, which begins “This seed is treated with . . . and ends “. . . from treated foodstuffs.” in quotation marks.

**(h) Page 6– Application Instructions Table – Leafy Vegetables:**

Change the sentence: “Containers or bags containing METALAXYL 28.35% FL ST treated spinach seed must be labeled with this restriction.” to “Containers or bags containing METALAXYL 28.35% FL ST treated spinach seed must be labeled with the restriction: “Do not grow spinach from treated seed in greenhouses.” ”

**(i) Page 8– Seed Treatment – Export Use:**

Seed treated in accordance with the export use application instructions must be labeled as follows: “For export only – NOT for domestic sale or use.”

3. Provide the following data as conditions of registration:
  - a. OPPTS number 830.6317 - storage stability - due April 1, 2011.
  - b. OPPTS number 830.6320 - corrosion characteristics - due April 1, 2011.
4. Submit one (1) copy of the revised final printed label before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions, contact Lisa Jones of my team at (703) 308-9424 or at [jones.lisa@epa.gov](mailto:jones.lisa@epa.gov).

Sincerely,



Mary L. Waller  
Product Manager (21)  
Fungicide Branch  
Registration Division (7505P)

Enclosure: Stamped copy of label  
Product Chemistry review dated 14 October 2009  
Acute Toxicity review dated 6 November 2009

4  
12

METALAXYL 28.35% FL ST  
Seed Treatment Fungicide

A SEED TREATMENT CHEMICAL FOR CONTROL OF SEED ROT AND DAMPING-OFF DISEASES OF CERTAIN CROPS

ACTIVE INGREDIENTS:

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl)alanine methyl ester . . . . .	28.35%
OTHER INGREDIENTS: . . . . .	71.65%
TOTAL: . . . . .	100.00%

Contains 2.65 lbs. of metalaxyl per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF SWALLOWED:	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
IF IN EYES:	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
In case of an emergency call CHEMTREC toll free at 1-800-424-9300. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	

EPA Reg. No. 42750-appl.

EPA Est. No. xxxxxx-xx-xxx

NET CONTENT: \_\_\_\_\_

Manufactured For:  
ALBAUGH, INC.  
ANKENY, IA 50021

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

**DEC 22 2009**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.**

**42750-208**

5/12

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### WARNING AVISO

Causes skin irritation. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with eyes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants,
- Socks and chemical-resistant footwear and
- Chemical-resistant gloves (such as nitrile, neoprene, and/or barrier laminate).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

#### ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate:

Groundwater Advisory: This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where solids are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not contaminate bodies of water when disposing of planting equipment wash water.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

6  
12

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in a cool dry location. Do not store in direct hot sunlight. Do not store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

### PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER DISPOSAL:

Non-refillable containers (1, or 2.5 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

(non-refillable <5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(non-refillable ≥5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: if the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls,
- socks and chemical resistant footwear,
- chemical resistant gloves, such as barrier laminate or nitrile rubber or neoprene rubber or Viton
- protective eyewear.

In California - Only for use in commercial seed treating establishments. Not allowed for "on-farm" use.

NOTE: Seed treated commercially or for sale must be dyed an unnatural color using an EPA approved dye before being sold. This seed is treated with METALAXYL 28.35% FL ST, a metalaxyl product. Do not allow children, pets or livestock to have access to treated seeds. Wear long pants, long sleeved shirt and protective gloves when handling treated seed. Treated seeds exposed on soil surface may e hazardous to wildlife. Cover or collect treated seed spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not contaminate bodies of water when disposing of planting equipment wash water. Dispose of seed packaging or containers in accordance with local requirements. Do not use for food feed, oil production or any other purpose except planting. Store away from treated foodstuffs.

METALAXYL 28.35% FL ST is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium anfrPhytophthora spp.* For control of other soilborne diseases, combination of Gustafson Captan and VITA VAX® registered formulations ate compatible.

Do not use with other seed treatment products unless previous experience assures compatibility.

METALAXYL 28.35% FL ST may be applied as a water-based slurry with other registered seed treatment insecticides and "fungicides through standard slurry or mist-type commercial seed treatment equipment.

NOTE: METALAXYL 28.35% FL ST is a systemic fungicide having a specific mode of action. METALAXYL 28.35% FL ST could be subject to envelopment of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Albaugh cannot assume liability for crop damage resulting from resistant strains of fungi. Consult your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible METALAXYL 28.35% FL ST resistant strains of fungi which may occur.

FAILURE to FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

CROP	APPLICATION INSTRUCTIONS
Canola* Mustard Seed* Rapeseed*	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.25 - 0.5 fl. oz. per 100 lbs of seed.  *Not registered for use in California.
Cotton	For <i>Pythium</i> seed rot and damping off: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed.
Soybeans	For <i>Pythium</i> damping off and early season <i>Phytophthora</i> control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed.
Peanuts*	For <i>Pythium</i> seed rot and damping off: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed.  * Not registered for use in California.

CROP	APPLICATION INSTRUCTIONS
Sorghum	<p>For control of <i>Pythium</i> damping-off apply 0.375 - 0.75 fl.oz. of METALAXYL 28.35% FL ST per 100 lbs. of seed applied as a seed treatment.</p> <p>For control of systemic downy mildew (races susceptible to chemical control), apply 0.75 - 1.5 fl. oz. METALAXYL 28.35% FL ST per 100 lbs. of seed on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew in heavily infected areas with recent severe history, apply 1.5 - 3.0 fl. oz. of METALAXYL 28.35% FL ST per 100 lbs. of seed as a seed treatment.</p> <p>Only high quality sorghum seed (90% germination and above) should be treated with METALAXYL 28.35% FL ST.</p>
Sunflowers	<p>For control of systemic downy mildew: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 1.5 - 3.0 fl. oz. per 100 lbs. of seed.</p>
Forage Legumes	<p>Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage).</p> <p>For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed.</p>
Grain Crops	<p>Wheat, barley, rye, oats, millet, milo, corn (field corn, sweet corn and popcorn), buckwheat and triticale.</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p> <p>For systemic downy mildew control of sweet corn: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.</p>
Rice*	<p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed. If using ALLEGIENCE-FL treated rice do not use Ridomil for control of <i>Pythium</i> in water-seeded rice.</p> <p>* Not registered for use in California.</p>
Seed and Pod Vegetables	<p>Black-eyed peas, chickpeas (Garbanzos), cowpeas, dill, edible soybeans, field beans, field peas, garden peas, green beans, kidney beans, lima beans, lupines, navy beans, okra, peas, pole beans, snap beans, string beans, wax beans, and lentils.</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p> <p>For control of <i>Pythium</i> damping-off and systemic downy mildew of peas: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.</p>
Turf Grasses	<p>For <i>Pythium</i> damping-off control on golf courses, home lawns reclamation and erosion control projects: Apply METALAXYL 28.35% FL ST as a seed treated at the rate of 1.5 fl. oz. per 100 lbs of seed.</p>



CROP	APPLICATION INSTRUCTIONS
<b>Forage Grasses</b>	<p>Grasses grown for hay, grazing or silage, corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay, grazing or silage.</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p>
<b>Brassica (Cole) Leafy Vegetables</b>	<p>Broccoli (including Chinese and raab), Brussels sprouts, cabbage (including Chinese bok choy, Chinese napa and mustard), cauliflower, cavalo broccoli, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p> <p>Not registered for this use in California.</p>
<b>Cucurbit Vegetables</b>	<p>Chayote (fruit), Chinese waxgourd, citron melon, cucumber, gherkin, edible gourd (includes hyotan, cucuza, Chinese okra, and hechima), Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), muskmelon (includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), pumpkin, summer squash (includes crookneck squash, scallop squash, straightneck squach, vegetable marrow, zucchini), winter squash (includes butternut squash, calabaza, hubbard squash), Cucumis mixta, Cucumis pepo (includes acorn squash, spaghetti squash), watermelon (includes hybrids and/or varieties of Citrullus lanatus).</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.</p> <p>Not registered for this use in California.</p>
<b>Leafy Vegetables</b>	<p>Amaranth (leafy, Chinese spinach, tampala), cardoon, celery (including Chinese), celtuce, chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden and upland), dandelion, dock, endive, fennel (finocchio), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio, rhubarb, spinach* (including New Zealand and vine), and Swiss chard.</p> <p>* Not registered for use in California.</p> <p>NOTE: Do not apply METALAXYL 28.35% FL ST to spinach which is destined to be grown in the greenhouse. Containers or bags containing METALAXYL 28.35% FL ST treated spinach seed must be labeled with this restriction.</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.</p>
<b>Fruiting Vegetables</b>	<p>Eggplants, ground cherry, pepino, pepper (including bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), tomatillo, and tomato</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 fl. oz. per 100 lbs of seed.</p>

CROP	APPLICATION INSTRUCTIONS
Onions	Onion (dry bulb and green)  For Pythium damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.
Root and Tuber Vegetables (except potatoes)	Arracacha, arrowroot, artichoke (Chinese and Jerusalem), beet garden and sugar), burdock (edible), canna (edible), carrot, cassava (bitter and sweet), celery root, chayote, chervil, chicory, chufa, dasheen, ginger, ginseng, horseradish, leren, parsley (turnip-rooted), parsnip, radish (includes oriental dalkon), rutabaga, salsify (includes black and Spanish), skirret, sweet potato, tanier, turmeric, turnip, yam bean (jicama, manico pea), yam  For Pythium damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 fl. oz./100 lbs. of seed.  NOTE: Do not apply METALAXYL 28.35% FL ST to potatoes.  Not registered for this use in California.

11/12

SEED TREATMENT – EXPORT USE

For treatment of the following seed for export only to countries requiring treatment with metalaxyl before importation.

Directions for use at reduced rates in combination with other fungicides METALAXYL 28.35% FL ST is a systemic fungicide seed dressing which may be applied in combination with other EPA-registered fungicides to aid in control of Pythium. METALAXYL 28.35% FL ST may be applied as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment.

CROP	APPLICATION INSTRUCTIONS
Brassica (Cole) Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 - 1.50 fl. oz. per 100 lbs. of seed.
Canola	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.50 - 1.70 fl. oz. per 100 lbs. of seed. (32 – 100 ml per 100 kg of seed)
Corn	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.  For systemic downy mildew control: Apply METALAXYL 28.35% FL ST as a seed treatment at 3.0 fl. oz. per 100 lbs. of seed.
Cucurbit Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.
Fruiting Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75-1.50 fl. oz. per 100 lbs. of seed.
Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.
Onions	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.
Millet	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.  For systemic downy mildew control: Apply METALAXYL 28.35% FL ST as a seed treatment at 2.5 fl. oz. per 100 lbs. of seed.
Peas	For <i>Pythium</i> damping-off and early season Phytophthora control: Apply METALAXYL 28.35% FL ST as a seed treatment at 3.5 fl. oz. per 100 lbs. of seed.
Root and Tuber Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.  Do not apply METALAXYL 28.35% FL ST to potatoes.
Sorghum	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.  For systemic downy mildew control: Apply METALAXYL 28.35% FL ST as a seed treatment at 5.25 fl. oz. per 100 lbs. of seed.
Sunflowers	For systemic downy mildew control: Apply METALAXYL 28.35% FL ST as a seed treatment at 9.0 fl. oz. per 100 lbs. of seed.

CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Albaugh, no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Albaugh warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. TO THE EXTENT PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, ALBAUGH SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, ALBAUGH NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. To the extent permitted by applicable law, Buyer assumes the responsibility to handle, use, and store this product in accordance with safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Albaugh. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.