

42750-208

07/29/2010

1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

JUL 29 2010

Mr. Morris Gaskins
Product Registration
Albaugh, Inc.
P.O. Box 2127
Valdosta, GA 31604-2127

SUBJECT: Application for Pesticide Notification (PRN 98-10)
Request Alternate Brand Names "TEBUSTAR 250 ST"
EPA Reg. No. 42750-130 and
"METALAXYL 265 ST"
EPA Reg. No. 42750-208
Application Dated June 18, 2010

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 06/18/10 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-5335 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Paul J. Mastradone, Ph.D., Acting
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
 Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 42750-208	2. EPA Product Manager M. Waller	3. Proposed Classification <input type="checkbox"/> None <input checked="" type="checkbox"/> Restricted
4. Company/Product (Name) Metalaxyl 28.35% FL ST	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) Albaugh, Inc. P.O. Box 2127 Valdosta, GA 31604 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

Amendment - Explain below.
 Resubmission in response to Agency letter dated _____
 Notification - Explain below.
 Final printed labels in response to Agency letter dated _____
 "Me Too" Application.
 Other - Explain below.

NOTIFICATION

JUL 29 2010

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of alternate brand name from "Metalaxyl 265 ST".

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no further changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under section 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted			
If "Yes" Unit Packaging wgt.		No. per container	
If "Yes" Package wgt		No. per container	

3. Location of Net Contents Information
 Label Container

4. Size(s) Retail Container
1 gal, 2.5 gal

5. Location of Label Directions
 Attached to container

6. Manner in Which Label is Affixed to Product
 Lithograph Paper glued Stenciled Other _____

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Morris Gaskins	Title Registrations Manager	Telephone No. (Include Area Code) 229-244-3286
------------------------	--------------------------------	---

Certification
 I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature

3. Title
Registrations Manager

4. Typed Name
Morris Gaskins

5. Date
June 18, 2010

6. Date Application Received
(Stamped)

CORPORATE OFFICE

525 NE 36th Street
Ankeny, IA 50021
515.964.9444 - Office
800.247.8013 - Toll Free
515.964.7813 - Facsimile

ALBAUGH, INC.

Valdosta Office
P.O. Box 2127
304 Janet Street, Suite H
Valdosta, GA 31604
229.244.3288 - Office
229.244.5841 - Facsimile

3/13

Fed-X

June 18, 2010

Document Processing Desk (NOTIFY)
Ms. Mary Waller (PM 21)
Registration Division
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard (South Bldg)
2777 South Crystal Dr.
Arlington, VA 22202

RE: Tebuconazole 28% FL ST
EPA Reg. No. 42750-130

Metalaxyl 28.35% FL ST
EPA Reg. No. 42750-208

Dear Ms. Waller,

The enclosed submission for the above referenced registrations are label notifications under PR Notice 98-10 for the following alternate brand names:


TEBUSTAR 250 ST

METALAXYL 265 ST

Three copies of each label are enclosed.

Please call if you have any questions.

Regards,



Morris Gaskins
Registrations Manager
Albaugh, Inc.
229-244-3288



PREMIER SUPPLIER OF OFF-PATENT CROP PROTECTION PRODUCTS

www.albaughinc.com

4/12

METALAXYL 265 ST
Seed Treatment Fungicide

A SEED TREATMENT CHEMICAL FOR CONTROL OF SEED ROT AND DAMPING-OFF DISEASES OF CERTAIN CROPS

ACTIVE INGREDIENTS:

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl)alanine methyl ester 28.35%

OTHER INGREDIENTS: 71.65%

TOTAL: 100.00%

Contains 2.65 lbs. of metalaxyl per gallon.

NOTIFICATION

JUL 29 2010

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
In case of an emergency call CHEMTREC toll free at 1-800-424-9300. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	

EPA Reg. No. 42750-208
AD122209

EPA Est. No. xxxxxx-xx-xxx

NET CONTENT: _____

Manufactured For:
ALBAUGH, INC.
1525 NE 36th ST
ANKENY, IA 50021

5/12

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING AVISO

Causes skin irritation. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with eyes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants,
- Socks and chemical-resistant footwear and
- Chemical-resistant gloves (such as nitrile \geq 14 mils in thickness, neoprene \geq 14 mils in thickness, Viton \geq 14 mils in thickness and/or barrier laminate).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where solids are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not contaminate bodies of water when disposing of planting equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your

State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: if the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls,
- socks and chemical resistant footwear,
- chemical resistant gloves, such as barrier laminate or nitrile rubber or neoprene rubber or Viton
- protective eyewear.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool dry location. Do not store in direct hot sunlight. Do not store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Non-refillable containers (1, or 2.5 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.
(non-refillable <5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

In California - Only for use in commercial seed treating establishments. Not allowed for "on-farm" use.

NOTE: Seed treated commercially or for sale must be dyed an unnatural color using an EPA approved dye before being sold.

Commercially treated seed must be labeled:

"This seed is treated with METALAXYL 265 ST, a metalaxyl product. Do not allow children, pets or

livestock to have access to treated seeds. Wear long pants, long sleeved shirt and protective gloves when handling treated seed. Treated seeds exposed on soil surface may e hazardous to wildlife. Cover or collect treated seed spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not contaminate bodies of water when disposing of planting equipment wash water. Dispose of seed packaging or containers in accordance with local requirements. Do not use for food feed, oil production or any other purpose except planting. Store away from treated foodstuffs."

METALAXYL 265 ST is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium anfrPhytophthora spp.* For control of other soilborne diseases, combination of Gustafson Captan and VITA VAX® registered formulations ate compatible.

Do not use with other seed treatment products unless previous experience assures compatibility.

METALAXYL 265 ST may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment. Follow more restrictive labeling requirements of tank mix products. Do not tank mix with any product that contains a prohibition on tank mixing.

NOTE: METALAXYL 265 ST is a systemic fungicide having a specific mode of action. METALAXYL 265 ST could be subject to envelopment of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Albaugh cannot assume liability for crop damage resulting from resistant strains of fungi. Consult your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible METALAXYL 265 ST resistant strains of fungi which may occur.

FAILURE to FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

CROP	APPLICATION INSTRUCTIONS
Canola* Mustard Seed* Rapeseed*	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.25 - 0.5 fl. oz. per 100 lbs of seed. *Not registered for use in California.
Cotton	For <i>Pythium</i> seed rot and damping off: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed.
Soybeans	For <i>Pythium</i> damping off and early season <i>Phytophthora</i> control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed.
Peanuts*	For <i>Pythium</i> seed rot and damping off: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed. * Not registered for use in California.

8/12

CROP	APPLICATION INSTRUCTIONS
Sorghum	<p>For control of <i>Pythium</i> damping-off apply 0.375 - 0.75 fl.oz. of METALAXYL 265 ST per 100 lbs. of seed applied as a seed treatment.</p> <p>For control of systemic downy mildew (races susceptible to chemical control), apply 0.75 - 1.5 fl. oz. METALAXYL 265 ST per 100 lbs. of seed on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew in heavily infected areas with recent severe history, apply 1.5 - 3.0 fl. oz. of METALAXYL 265 ST per 100 lbs. of seed as a seed treatment.</p> <p>Only high quality sorghum seed (90% germination and above) should be treated with METALAXYL 265 ST.</p>
Sunflowers	<p>For control of systemic downy mildew: Apply METALAXYL 265 ST as a seed treatment at the rate of 1.5 - 3.0 fl. oz. per 100 lbs. of seed.</p>
Forage Legumes	<p>Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage).</p> <p>For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed.</p>
Grain Crops	<p>Wheat, barley, rye, oats, millet, milo, corn (field corn, sweet corn and popcorn), buckwheat and triticale.</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p> <p>For systemic downy mildew control of sweet corn: Apply METALAXYL 265 ST as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.</p>
Rice*	<p>For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed. If using ALLEGIENCE-FL treated rice do not use Ridomil for control of <i>Pythium</i> in water-seeded rice.</p> <p>* Not registered for use in California.</p>
Seed and Pod Vegetables	<p>Black-eyed peas, chickpeas (Garbanzos), cowpeas, dill, edible soybeans, field beans, field peas, garden peas, green beans, kidney beans, lima beans, lupines, navy beans, okra, peas, pole beans, snap beans, string beans, wax beans, and lentils.</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p> <p>For control of <i>Pythium</i> damping-off and systemic downy mildew of peas: Apply METALAXYL 265 ST as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.</p>
Turf Grasses	<p>For <i>Pythium</i> damping-off control on golf courses, home lawns reclamation and erosion control projects: Apply METALAXYL 265 ST as a seed treated at the rate of 1.5 fl. oz. per 100 lbs of seed.</p>

CROP	APPLICATION INSTRUCTIONS
Forage Grasses	<p>Grasses grown for hay, grazing or silage, corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay, grazing or silage.</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p>
Brassica (Cole) Leafy Vegetables	<p>Broccoli (including Chinese and raab), Brussels sprouts, cabbage (including Chinese bok choy, Chinese napa and mustard), cauliflower, cavalo broccoli, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p> <p>Not registered for this use in California.</p>
Cucurbit Vegetables	<p>Chayote (fruit), Chinese waxgourd, citron melon, cucumber, gherkin, edible gourd (includes hyotan, cucuza, Chinese okra, and hechima), Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), muskmelon (includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), pumpkin, summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), winter squash (includes butternut squash, calabaza, hubbard squash), Cucumis mixta, Cucumis pepo (includes acorn squash, spaghetti squash), watermelon (includes hybrids and/or varieties of Citrullus lanatus).</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.</p> <p>Not registered for this use in California.</p>
Leafy Vegetables	<p>Amaranth (leafy, Chinese spinach, tampala), cardoon, celery (including Chinese), celtuce, chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden and upland), dandelion, dock, endive, fennel (finocchio), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio, rhubarb, spinach* (including New Zealand and vine), and Swiss chard.</p> <p>* Not registered for use in California.</p> <p>NOTE: Do not apply METALAXYL 265 ST to spinach which is destined to be grown in the greenhouse. Containers or bags containing METALAXYL 265 ST treated spinach seed must be labeled with the restriction: "Do not grow spinach from treated seed in greenhouses."</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.</p>
Fruiting Vegetables	<p>Eggplants, ground cherry, pepino, pepper (including bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), tomatillo, and tomato</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 fl. oz. per 100 lbs of seed.</p>

10/12

CROP	APPLICATION INSTRUCTIONS
Onions	Onion (dry bulb and green) For Pythium damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.
Root and Tuber Vegetables (except potatoes)	Arracacha, arrowroot, artichoke (Chinese and Jerusalem), beet garden and sugar), burdock (edible), canna (edible), carrot, cassava (bitter and sweet), celery root, chayote, chervil, chicory, chufa, dasheen, ginger, ginseng, horseradish, leren, parsley (turnip-rooted), parsnip, radish (includes oriental dalkon), rutabaga, salsify (includes black and Spanish), skirret, sweet potato, tanier, turmeric, turnip, yam bean (jicama, manico pea), yam For Pythium damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 fl. oz./100 lbs. of seed. NOTE: Do not apply METALAXYL 265 ST to potatoes. Not registered for this use in California.

11/12

SEED TREATMENT – EXPORT USE

For treatment of the following seed for export only to countries requiring treatment with metalaxyl before importation.

Seed treated in accordance with the export use application instructions must be labeled as follows: "For export only – NOT for domestic sale or use."

Directions for use at reduced rates in combination with other fungicides METALAXYL 265 ST is a systemic fungicide seed dressing which may be applied in combination with other EPA-registered fungicides to aid in control of Pythium. METALAXYL 265 ST may be applied as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment.

CROP	APPLICATION INSTRUCTIONS
Brassica (Cole) Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.75 - 1.50 fl. oz. per 100 lbs. of seed.
Canola	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at the rate of 0.50 - 1.70 fl. oz. per 100 lbs. of seed. (32 – 100 ml per 100 kg of seed)
Corn	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply METALAXYL 265 ST as a seed treatment at 3.0 fl. oz. per 100 lbs. of seed.
Cucurbit Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.
Fruiting Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75- 1.50 fl. oz. per 100 lbs. of seed.
Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.
Onions	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.
Millet	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply METALAXYL 265 ST as a seed treatment at 2.5 fl. oz. per 100 lbs. of seed.
Peas	For <i>Pythium</i> damping-off and early season Phytophthora control: Apply METALAXYL 265 ST as a seed treatment at 3.5 fl. oz. per 100 lbs. of seed.
Root and Tuber Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed. Do not apply METALAXYL 265 ST to potatoes.
Sorghum	For <i>Pythium</i> damping-off control: Apply METALAXYL 265 ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed. For systemic downy mildew control: Apply METALAXYL 265 ST as a seed treatment at 5.25 fl. oz. per 100 lbs. of seed.
Sunflowers	For systemic downy mildew control: Apply METALAXYL 265 ST as a seed treatment at 9.0 fl. oz. per 100 lbs. of seed.

12/12

CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Albaugh, no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Albaugh warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. TO THE EXTENT PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, ALBAUGH SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, ALBAUGH NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. To the extent permitted by applicable law, Buyer assumes the responsibility to handle, use, and store this product in accordance with safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Albaugh. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.