

42750-208

5/15/2014

jack
1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Mr. Morris Gaskins
Albaugh Inc.
P.O. Box 2127
Valdosta, GA 31604

MAY 15 2014

Subject: Metalaxyl 28.35% FL ST
EPA Reg. No. 42750-208
EPA Decision Number: 490708
Your label submitted on 4/18/14 and resubmitted on 5/14/14 to add instructions for corn, cotton and soybean to include rates per seed count

Dear Mr. Gaskins:

The label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Heather Garvie by phone at: 703-308-0034 or via email at: garvie.heather@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Hope A. Johnson".

Hope A. Johnson
Product Manager 21
Fungicide Branch
Registration Division

2/11

METALAXYL 28.35% FL ST
Seed Treatment Fungicide

A SEED TREATMENT CHEMICAL FOR CONTROL OF SEED ROT AND
DAMPING-OFF DISEASES OF CERTAIN CROPS

ACTIVE INGREDIENTS:

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl)alanine methyl ester 28.35%

OTHER INGREDIENTS: 71.65%

TOTAL: 100.00%

Contains 2.65 lbs. of metalaxyl per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID


IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
In case of an emergency call CHEMTREC toll free at 1-800-424-9300. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	

EPA Reg. No. 42750-208

EPA Est. No. xxxxxx-xx-xxx

NET CONTENT: _____

Manufactured For:
ALBAUGH, INC.
1525 NE 36th ST
ANKENY, IA 50021



3/11

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING AVISO

Causes skin irritation. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with eyes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants,
- Socks and chemical-resistant footwear and
- Chemical-resistant gloves (such as nitrile \geq 14 mils in thickness, neoprene \geq 14 mils in thickness, Viton \geq 14 mils in thickness and/or barrier laminate).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: This chemical as properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not contaminate bodies of water when disposing of planting equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: if the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls,
- socks and chemical resistant footwear,
- chemical resistant gloves, such as barrier laminate or nitrile rubber or neoprene rubber or Viton
- protective eyewear.

In California - Only for use in commercial seed treating establishments. Not allowed for "on-farm" use.

NOTE: Seed treated commercially or for sale must be dyed an unnatural color using an EPA approved dye before being sold.

Commercially treated seed must be labeled:

"This seed is treated with METALAXYL 28.35% FL ST, a fungicide containing metalaxyl. Do not allow children, pets or livestock to have access to treated seeds. Wear long pants, long sleeved shirt and protective gloves when handling treated seed. Treated seeds exposed on soil surface maybe hazardous to wildlife. Cover or collect treated seed spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not contaminate bodies of water when disposing of planting equipment wash water. Dispose of seed packaging or containers in accordance with local requirements. Do not use for food, feed, oil production or any other purpose except planting. Store away from treated foodstuffs."

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.

METALAXYL 28.35% FL ST is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium* and *Phytophthora spp.* For control of other soilborne diseases, combination of Gustafson Captan and VITA VAX® registered formulations are compatible.

Do not use with other seed treatment products unless previous experience assures compatibility.

METALAXYL 28.35% FL ST may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment. Follow more restrictive labeling requirements of tank mix products. Do not tank mix with any product that contains a prohibition on tank mixing.

NOTE: METALAXYL 28.35% FL ST is a systemic fungicide having a specific mode of action.

5/11

METALAXYL 28.35% FL ST could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Consult your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible METALAXYL 28.35% FL ST resistant strains of fungi which may occur.

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

CROP	APPLICATION INSTRUCTIONS					
Canola* Mustard Seed* Rapeseed*	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.25 - 0.5 fl. oz. per 100 lbs of seed. *Not registered for use in California.					
Cotton	For <i>Pythium</i> seed rot and damping-off. Apply METALAXYL 28.35% FL ST as a seed treatment:					
	METALAXYL 28.35% FL ST Application rate			Maximum Planting rate - seeds/A*		
	seeds/lb	fl oz/CWT	mg Al/seed	@ 0.75 fl oz/CWT	@1.5 fl oz/CWT	
	3600	0.75 - 1.5	0.019 - 0.038	96,320	48,160	
	4400	0.75 - 1.5	0.015 - 0.031	117,700	58,860	
	5200	0.75 - 1.5	0.013 - 0.026	139,100	69,560	
	6000	0.75 - 1.5	0.011 - 0.023	160,500	80,270	
* Do not apply by seed treatment more than 0.004 pounds of Metalaxyl (0.2 fl. oz. METALAXYL 28.35% FL ST) per acre per year.						
Soybeans	For <i>Pythium</i> damping off and early season <i>Phytophthora</i> control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed.					
	METALAXYL 28.35% FL ST Application rate			Maximum Planting rate - seeds/A*		
	seeds/lb	fl oz/CWT	Fl. Oz./140,000 seed unit	mg Al/seed	@ 0.75 fl oz/CWT	@1.5 fl oz/CWT
	2000	0.75 - 1.5	0.53 - 1.05	0.034 - 0.068	222,100	111,000
	2400	0.75 - 1.5	0.44 - 0.88	0.028 - 0.057	266,500	133,200
	2800	0.75 - 1.5	0.38 - 0.75	0.024 - 0.048	310,900	155,400
	3200	0.75 - 1.5	0.33 - 0.66	0.021 - 0.042	355,300	177,700
3600	0.75 - 1.5	0.29 - 0.58	0.019 - 0.038	399,700	199,900	
* Do not apply by seed treatment more than 0.0166 pounds of Metalaxyl (0.83 fl. oz. METALAXYL 28.35% FL ST) per acre per year.						
Peanuts*	For <i>Pythium</i> seed rot and damping off: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed. * Not registered for use in California.					
Sorghum	For control of <i>Pythium</i> damping-off apply 0.375 - 0.75 fl.oz. of METALAXYL 28.35% FL ST per 100 lbs. of seed applied as a seed treatment. For control of systemic downy mildew (races susceptible to chemical control), apply 0.75 - 1.5 fl. oz. METALAXYL 28.35% FL ST per 100 lbs. of seed on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew in heavily infected areas with recent severe history, apply 1.5 - 3.0 fl. oz. of METALAXYL 28.35% FL ST per 100 lbs. of seed as a seed treatment. Only high quality sorghum seed (90% germination and above) should be treated with METALAXYL 28.35% FL ST.					

6/11

CROP	APPLICATION INSTRUCTIONS
Sunflowers	For control of systemic downy mildew: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 1.5 - 3.0 fl. oz. per 100 lbs. of seed.
Forage Legumes	<p>Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage).</p> <p>For <i>Pythium</i> damping-off and early season <i>Phytophthora</i> control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed.</p>
Grain Crops	<p>Wheat, barley, rye, oats, millet, milo, corn (field corn, sweet corn and popcorn), buckwheat and triticale.</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p> <p>For systemic downy mildew control of sweet corn: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.</p>
Rice*	<p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 - 1.5 fl. oz. per 100 lbs. of seed. If using metalaxyl treated rice do not use Ridomil for control of <i>Pythium</i> in water-seeded rice.</p> <p>* Not registered for use in California.</p>
Seed and Pod Vegetables	<p>Black-eyed peas, chickpeas (Garbanzos), cowpeas, dill, edible soybeans, field beans, field peas, garden peas, green beans, kidney beans, lima beans, lupines, navy beans, okra, peas, pole beans, snap beans, string beans, wax beans, and lentils.</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p> <p>For control of <i>Pythium</i> damping-off and systemic downy mildew of peas: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 1.5 fl. oz. per 100 lbs. of seed.</p>
Turf Grasses	For <i>Pythium</i> damping-off control on golf courses, home lawns reclamation and erosion control projects: Apply METALAXYL 28.35% FL ST as a seed treated at the rate of 1.5 fl. oz. per 100 lbs of seed.
Forage Grasses	<p>Grasses grown for hay, grazing or silage, corn (fodder or silage), sorghum (hay or silage), and small grain grown for hay, grazing or silage.</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p>
Brassica (Cole) Leafy Vegetables*	<p>Broccoli (including Chinese and raab), Brussels sprouts, cabbage (including Chinese bok choy, Chinese napa and mustard), cauliflower, cavalo broccoli, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 fl. oz. per 100 lbs. of seed.</p> <p>*Not registered for this use in California.</p>
Cucurbit Vegetables*	Chayote (fruit), Chinese waxgourd, citron melon, cucumber, gherkin, edible gourd (includes hyotan, cucuza, Chinese okra, and hechima), Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), muskmelon (includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), pumpkin, summer

7/11

CROP	APPLICATION INSTRUCTIONS
	<p>squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), winter squash (includes butternut squash, calabaza, hubbard squash), Cucumis mixta, Cucumis pepo (includes acorn squash, spaghetti squash), watermelon (includes hybrids and/or varieties of Citrullus lanatus).</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.</p> <p>*Not registered for this use in California.</p>
<p>Leafy Vegetables*</p>	<p>Amaranth (leafy, Chinese spinach, tampala), cardoon, celery (including Chinese), celtuce, chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden and upland), dandelion, dock, endive, fennel (finocchio), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio, rhubarb, spinach* (including New Zealand and vine), and Swiss chard.</p> <p>* Not registered for use in California.</p> <p>NOTE: Do not apply METALAXYL 28.35% FL ST to spinach which is destined to be grown in the greenhouse. Containers or bags containing METALAXYL 28.35% FL ST treated spinach seed must be labeled with the restriction: "Do not grow spinach from treated seed in greenhouses."</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.</p>
<p>Fruiting Vegetables</p>	<p>Eggplants, ground cherry, pepino, pepper (including bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), tomatillo, and tomato</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 fl. oz. per 100 lbs of seed.</p>
<p>Onions</p>	<p>Onion (dry bulb and green)</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.</p>
<p>Root and Tuber Vegetables (except potatoes)*</p>	<p>Arracacha, arrowroot, artichoke (Chinese and Jerusalem), beet (garden and sugar), burdock (edible), canna (edible), carrot, cassava (bitter and sweet), celery root, chayote, chervil, chicory, chufa, dasheen, ginger, ginseng, horseradish, leren, parsley (turnip-rooted), parsnip, radish (includes oriental dalkon), rutabaga, salsify (includes black and Spanish), skirret, sweet potato, tanier, turmeric, turnip, yam bean (jicama, manico pea), yam</p> <p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 fl. oz./100 lbs. of seed.</p> <p>NOTE: Do not apply METALAXYL 28.35% FL ST to potatoes.</p> <p>*Not registered for this use in California.</p>

Directions for Use at Reduced Rates in Combination With Other Fungicides:

This product is a systemic fungicide seed dressing which may be applied in combinations with other EPA registered fungicides to aid in control of *Pythium*. This product may be applied as a water-based slurry through standard slurry or mist type commercial seed treatment equipment.

SEED	INSTRUCTIONS FOR USE
Corn (field), Wheat, Oats, Barley, Rye	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.10 - 0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of seed treatment fungicides.
Soybeans, Field beans, Kidney beans, Navy beans, Pinto beans, Lentils, Peas	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.10 - 0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides.
Sunflowers	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.10 - 0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides.
Peanuts*	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.10 - 0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides. *Not registered for use in California.
Rice*	To aid in the control of seed decay and damping-off caused by <i>Pythium</i> : Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.10 - 0.375 fl. oz. per 100 lbs. of seed. Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides. *Not registered for use in California.

SEED TREATMENT – EXPORT USE

For treatment of the following seed for export only to countries requiring treatment with metalaxyl before importation.

Seed treated in accordance with the export use application instructions must be labeled as follows: "For export only – NOT for domestic sale or use."

Directions for use at reduced rates in combination with other fungicides METALAXYL 28.35% FL ST is a systemic fungicide seed dressing which may be applied in combination with other EPA-registered fungicides to aid in control of *Pythium*. METALAXYL 28.35% FL ST may be applied as a water-based slurry through standard slurry or mist-type commercial seed treatment equipment.

CROP	APPLICATION INSTRUCTIONS
Brassica (Cole) Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.75 - 1.50 fl. oz. per 100 lbs. of seed.
Canola	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at the rate of 0.50 - 1.70 fl. oz. per 100 lbs. of seed. (32 – 100 ml per 100 kg of seed)

CROP	APPLICATION INSTRUCTIONS																																				
Corn	<p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.</p> <p>For systemic downy mildew control: Apply METALAXYL 28.35% FL ST as a seed treatment at 3.0 fl. oz. per 100 lbs. of seed.</p> <table border="1" data-bbox="391 380 1377 611"> <thead> <tr> <th colspan="4" data-bbox="391 380 1062 436">METALAXYL 28.35% FL ST Application rate</th> <th colspan="2" data-bbox="1062 380 1377 436">Maximum Planting rate - seeds/A*</th> </tr> <tr> <th data-bbox="391 436 532 491">seeds/lb</th> <th data-bbox="532 436 695 491">fl oz/CWT</th> <th data-bbox="695 436 867 491">Fl. oz./80,000 seed unit</th> <th data-bbox="867 436 1062 491">mg AI/seed</th> <th data-bbox="1062 436 1159 491">@ 0.1 fl oz/CWT</th> <th data-bbox="1159 436 1377 491">@ 0.375 fl oz/CWT</th> </tr> </thead> <tbody> <tr> <td data-bbox="391 491 532 525">1200</td> <td data-bbox="532 491 695 525">0.1 - 0.375</td> <td data-bbox="695 491 867 525">0.07 - 0.25</td> <td data-bbox="867 491 1062 525">0.008 - 0.028</td> <td data-bbox="1062 491 1159 525">457,500</td> <td data-bbox="1159 491 1377 525">122,000</td> </tr> <tr> <td data-bbox="391 525 532 558">1600</td> <td data-bbox="532 525 695 558">0.1 - 0.375</td> <td data-bbox="695 525 867 558">0.05 - 0.19</td> <td data-bbox="867 525 1062 558">0.006 - 0.021</td> <td data-bbox="1062 525 1159 558">610,000</td> <td data-bbox="1159 525 1377 558">162,700</td> </tr> <tr> <td data-bbox="391 558 532 592">2000</td> <td data-bbox="532 558 695 592">0.1 - 0.375</td> <td data-bbox="695 558 867 592">0.04 - 0.15</td> <td data-bbox="867 558 1062 592">0.005 - 0.017</td> <td data-bbox="1062 558 1159 592">762,500</td> <td data-bbox="1159 558 1377 592">203,300</td> </tr> <tr> <td data-bbox="391 592 532 625">2400</td> <td data-bbox="532 592 695 625">0.1 - 0.375</td> <td data-bbox="695 592 867 625">0.03 - 0.13</td> <td data-bbox="867 592 1062 625">0.004 - 0.014</td> <td data-bbox="1062 592 1159 625">915,000</td> <td data-bbox="1159 592 1377 625">244,000</td> </tr> </tbody> </table> <p data-bbox="391 611 1377 674">* Do not apply by seed treatment more than 0.0076 pounds of Metalaxyl (0.46 fl. oz. METALAXYL 28.35% FL ST) per acre per year.</p>	METALAXYL 28.35% FL ST Application rate				Maximum Planting rate - seeds/A*		seeds/lb	fl oz/CWT	Fl. oz./80,000 seed unit	mg AI/seed	@ 0.1 fl oz/CWT	@ 0.375 fl oz/CWT	1200	0.1 - 0.375	0.07 - 0.25	0.008 - 0.028	457,500	122,000	1600	0.1 - 0.375	0.05 - 0.19	0.006 - 0.021	610,000	162,700	2000	0.1 - 0.375	0.04 - 0.15	0.005 - 0.017	762,500	203,300	2400	0.1 - 0.375	0.03 - 0.13	0.004 - 0.014	915,000	244,000
METALAXYL 28.35% FL ST Application rate				Maximum Planting rate - seeds/A*																																	
seeds/lb	fl oz/CWT	Fl. oz./80,000 seed unit	mg AI/seed	@ 0.1 fl oz/CWT	@ 0.375 fl oz/CWT																																
1200	0.1 - 0.375	0.07 - 0.25	0.008 - 0.028	457,500	122,000																																
1600	0.1 - 0.375	0.05 - 0.19	0.006 - 0.021	610,000	162,700																																
2000	0.1 - 0.375	0.04 - 0.15	0.005 - 0.017	762,500	203,300																																
2400	0.1 - 0.375	0.03 - 0.13	0.004 - 0.014	915,000	244,000																																
Cucurbit Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 fl. oz. per 100 lbs. of seed.																																				
Fruiting Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75-1.50 fl. oz. per 100 lbs. of seed.																																				
Leafy Vegetables	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.																																				
Onions	For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.																																				
Millet	<p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.</p> <p>For systemic downy mildew control: Apply METALAXYL 28.35% FL ST as a seed treatment at 2.5 fl. oz. per 100 lbs. of seed.</p>																																				
Peas	For <i>Pythium</i> damping-off and early season Phytophthora control: Apply METALAXYL 28.35% FL ST as a seed treatment at 3.5 fl. oz. per 100 lbs. of seed.																																				
Root and Tuber Vegetables	<p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.</p> <p>Do not apply METALAXYL 28.35% FL ST to potatoes.</p>																																				
Sorghum	<p>For <i>Pythium</i> damping-off control: Apply METALAXYL 28.35% FL ST as a seed treatment at 0.75 - 1.50 fl. oz. per 100 lbs. of seed.</p> <p>For systemic downy mildew control: Apply METALAXYL 28.35% FL ST as a seed treatment at 5.25 fl. oz. per 100 lbs. of seed.</p>																																				
Sunflowers	For systemic downy mildew control: Apply METALAXYL 28.35% FL ST as a seed treatment at 9.0 fl. oz. per 100 lbs. of seed.																																				

10/11

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool dry location. Do not store in direct hot sunlight. Do not store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

(non-refillable ≤ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(non-refillable >5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container:

Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinseate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Albaugh, no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Albaugh warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. TO THE EXTENT PERMITTED BY APPLICABLE

11/11

LAW, ALBAUGH SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, ALBAUGH NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. To the extent permitted by applicable law, Buyer assumes the responsibility to handle, use, and store this product in accordance with safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Albaugh. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.