

42750-229

8/15/2012

1/32



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Mr Morris Gaskins
Registrations Manager
Albaugh Inc
P O Box 2127
Valdosta GA 31604

AUG 15 2012

Subject Label Amendment – Adding Use on Potatoes Incorporating Cotton Dry Bean and Snap Bean
Uses from Supplemental Labels and Making Numerous Other Edits and Revisions
Product Name Fomesafen 22 1% SC
EPA Registration Number 42750 229
Submission Date May 10 2012
Decision Number 465115

Dear Mr Gaskins

The label amendment referred to above which also included submitted in connection with registration under the Federal Insecticide Fungicide and Rodenticide Act (FIFRA) as amended is acceptable provide the following item(s) are addressed

- 1) Correct the typographical error on page 25 under the GLYPHOSATE TOLERANT SOYBEAN TANK MIXES header in the third to last item changing ti to to for the text in the parentheses to read (refer to the Regional Use Map)
- 2) NOTE Marketing claims made on the pesticide label must be substantiated by data maintained in your files If data supporting the marketing claims made on the product label is not available then those claims must be removed
- 3) NOTE Should you wish to add/retain a reference to the company s website on your label then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency If the website is false or misleading the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E) 40 CFR 156 10(a)(5) list examples of statements EPA may consider false or misleading In addition regardless of whether a website is referenced on your product s label claims made on the website may not substantially differ from those claims approved through the registration process Therefore should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration the website will be referred to the EPA s Office of Enforcement and Compliance

SEE NEXT PAGE

Page 2 of 2
Label Amendment
Product Name Fomesafen 22 1% SC
EPA Registration Number 42750 229
Submission Date May 10 2012
Decision Number 465115

Products shipped after 18 months from the date on this notice or the next printing of the master label whichever occurs first, must bear the new label language If these conditions are not complied with the registration will be subject to cancellation in accordance with FIFRA Your release for shipment of this product constitutes acceptance of these conditions You must submit one copy of the final printed label before the product is released for shipment If you have any questions please contact Michael Walsh by phone at (703) 308 2972 or via email at walsh michael@epa gov

Sincerely



Kathryn V Montague
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

Enclosure

EDITOR S NOTE 5/10/12 marked draft label amendment to add potato use and roll up previously approved bulletins for cotton and beans

FOMESAFEN 22 1% SC

For Control of Weeds in Cotton, Dry Beans, Potatoes, Snap Beans and Soybeans

ACTIVE INGREDIENT	Sodium salt of fomesafen	
	5 [2 chloro 4 (trifluoromethyl)phenoxy] N (methylsulfonyl) 2 nitrobenzarnide	22 1%
OTHER INGREDIENTS		<u>77 9%</u>
TOTAL		100 0%

Equivalent to 21 0% fomesafen or 1 88 lbs fomesafen active ingredient per gal

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you do not understand the label find someone to explain it to you in detail)

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing Call a Poison Control Center or doctor for treatment advice
IF SWALLOWED	<ul style="list-style-type: none"> Call a Poison Control Center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to by a Poison Control Center or doctor Do not give anything by mouth to an unconscious person
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a Poison Control Center or doctor for treatment advice
IF INHALED	<ul style="list-style-type: none"> Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth to mouth if possible Call a Poison Control Center or doctor for further treatment advice
NOTE TO PHYSICIAN Probable mucosal damage may contraindicate the use of gastric lavage	
Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment	
HOT LINE NUMBER For 24 Hour Emergency Assistance call CHEMTREC at 1 800 424 9300	

EPA Reg No 42750 229

EPA Est No 42750 MO 001

NET CONTENTS _____ Gals

MANUFACTURED BY
Albaugh Inc
Ankeny IA 50021

ACCEPTED
with COMMENTS
In EPA Letter Dated
AUG 15 2012

Under the Federal Fungicide and Rodenticide Act as amended for the pesticide registered under EPA Reg No

42750-229

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER/PELIGRO

This product contains fomesafen which has been determined to cause tumors in laboratory animals (mice) Risks can be reduced by closely following use directions and precautions and by wearing the protective clothing specified elsewhere on this label

Corrosive Causes irreversible eye damage Causes skin irritation Harmful if absorbed through skin Harmful if swallowed Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals Do not get on skin Do not get in eyes or on clothing

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below If you want more options follow the instructions for category E on an EPA chemical resistance category selection chart

Applicators and other handlers must wear

Coveralls over short sleeved shirt and short pants

Chemical resistant gloves such as barrier laminate nitrile rubber neoprene rubber or Viton®

Chemical resistant footwear plus socks

Protective eyewear

Chemical resistant apron when cleaning equipment mixing or loading

In addition for aerial applications mixers and loaders handling more than 150 gallons of FOMESAFEN 22 1% SC in any single workday must wear

Dust/mist filtering NIOSH approved respirator with any N R P or HE filter

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product s concentrate Do not reuse them Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables exist, use detergent and hot water Keep and wash PPE separately from other laundry

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170 240 (d) (4 6)], the handler PPE requirements may be reduced or modified as specified in the WPS

USER SAFETY RECOMMENDATIONS

Users should

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Do not contaminate water when disposing of equipment wash water or rinsate Do not apply when weather conditions favor drift from target area

5/32

Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to spray drift and run off of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months after application. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of fomesafen from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. See the manual for Conservation Buffers to Reduce Pesticide Losses at the following internet address: <http://www.wsi.nrcs.usda.gov/products/W2Q/pest/core4.html>

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate, nitrile rubber, neoprene rubber, or Viton
- Chemical-resistant footwear plus socks

Protective eyewear

PRODUCT INFORMATION

Read all label directions before using.

FOMESAFEN 22.1% SC is a selective herbicide which may be applied preplant, preemergence, or postemergence for control or suppression of broadleaf weeds, grasses, and sedges in **cotton, dry beans, potatoes, snap beans** and soybeans.

Preplant Surface and Preemergence Applications

Certain germinating broadleaf weeds grasses and sedges may be controlled or suppressed by soil residual activity from either preplant preemergent or postemergent applications if rainfall occurs shortly after application The extent and consistency of soil activity is dependent upon soil characteristics ground cover amount of rainfall following application and the rate of FOMESAFEN 22 1% SC used

Postemergence Applications

FOMESAFEN 22 1% SC is generally most effective and consistent when used postemergence working through contact action Therefore emerged weeds must have thorough spray coverage for effective control

Some bronzing crinkling or spotting of soybean leaves may occur following a postemergent application but soybeans soon outgrow these effects and develop normally

Optimum weed control is achieved by postemergent applications of FOMESAFEN 22 1% SC to young actively growing broadleaf weeds that are not under stress from moisture temperature low soil fertility mechanical or chemical injury

Soil Characteristics

Application of FOMESAFEN 22 1% SC to soils with high organic matter and/or high clay content may require higher rates than soils with low organic matter and/or low clay content Refer to the Regional Use Map, weed control tables, and specific crop use sections for recommendations on use rates based on soil texture

Environmental and Agronomic Conditions

Always apply FOMESAFEN 22 1% SC under favorable environmental conditions that promote active weed growth Avoid applying FOMESAFEN 22 1% SC to weeds or labeled crops which are under stress from drought, extreme temperatures, excessive water, low humidity, low soil fertility, mechanical or chemical injury as reduced weed control and/or increased crop injury may result

Rainfastness

FOMESAFEN 22 1% SC requires a 1 hour rain free period for best results when applied postemergence

Cultivation

Cultivation prior to postemergence application is not recommended Cultivation may put weeds under stress, reducing weed control Timely cultivation 1 3 weeks after applying FOMESAFEN 22 1% SC may assist weed control

Information on Weed Resistance

Naturally occurring biotypes of certain broadleaf species with resistance to this herbicide and related products (same mode of action) are known to exist Selection of resistant biotypes through repeated use of these herbicides may result in control failures

If poor performance cannot be attributed to adverse weather conditions or improper application methods a resistant biotype may be present In such a case additional treatments with this herbicide or similar mode of action products are not recommended Consult your local company representative or agricultural advisor for assistance

APPLICATION DIRECTIONS

Application Timing

Best broad spectrum postemergence control of susceptible broadleaf weeds is obtained when FOMESAFEN 22 1% SC is applied early to actively growing weeds. This usually occurs 14 to 28 days after planting. Refer to the weed control tables for specific recommendations on weed growth stages and rates.

Spray Additives

Only spray additives cleared for use on growing crops under 40 CFR 180.1001 may be used in the spray mixture.

For best broad spectrum postemergence control of susceptible broadleaf weeds in Regions 2, 3, 4 and 5 (see Regional Use Maps), FOMESAFEN 22 1% SC should be used with 1.0-2.5% v/v liquid nitrogen (28% or similar) or a minimum of 8.5 lbs ammonium sulfate per 100 gals of spray volume.

For Postemergence Applications Always Add One of the Following (except in tank mix with products prohibiting spray additives) (See Tank Mix Directions for Use)

Nonionic Surfactant (NIS)

Use NIS containing at least 80% active ingredient at 0.25 to 0.5% v/v (2.4 qts /100 gals) of finished spray volume (Region 1 and East of Interstates 79 and 77 for Regions 2 and 3).

Crop Oil Concentrate (COC) or Methylated Seed Oil (MSO)

Use a nonphytotoxic COC or MSO containing 15-20% approved emulsifier at 0.5-1% v/v (2.4 qts /100 gals) of finished spray volume. COC or MSO can improve weed control but may slightly reduce crop tolerance.

Other Adjuvants

Adjuvants other than COC or NIS may be used providing the product meets the following criteria:

1. Contains only EPA exempt ingredients
2. Is nonphytotoxic to the target crop
3. Is compatible in mixture (May be established through a jar test)
4. Is supported locally for use with FOMESAFEN 22 1% SC on the target crop through proven field trials and through university and extension recommendations

Note: no adjuvants are needed for preplant or preemergence applications unless FOMESAFEN 22 1% SC is being used in a burndown.

Recommended Mixing Order

1. Fill spray tank with half the required amount of water and begin agitation
2. Add fertilizer (UAN, AMS)
3. Add dry pesticide formulations
4. Add FOMESAFEN 22 1% SC
5. Add liquid pesticide formulation
6. Add adjuvant (MSO, COC or NIS)
7. Add remainder of water and then maintain constant agitation

Compatibility agent: 1 gal /500 gals of water or 0.2% v/v may be added as needed.

Tank Mix Compatibility Test

A jar test is recommended prior to tank mixing to ensure compatibility of FOMESAFEN 22 1% SC with mixture partners Add proportion amounts of tank mixture components in a clear quart jar one at a time in the recommended mixing order Gently shake or invert capped jar and let stand for 15 30 minutes If the mixture clumps, forms flakes, oily films or layers or other precipitates, it is not compatible and the tank mixture should not be used

GROUND APPLICATION

Preplant Surface and Preemergence Application Use a minimum of 10 gallons per acre Nozzle selection should meet manufacturer's gallonage and pressure recommendations for preplant surface or preemergence applications

Postemergence Application Use sufficient spray volume and pressure to ensure complete coverage of the target weed A spray volume of 10 20 gallons per acre and 30 60 psi at the nozzle tip is recommended On large weeds and/or dense foliage, use 60 psi and a minimum of 20 gallons per acre to ensure coverage of weed foliage

Ground Application

~~Use sufficient spray volume and pressure to ensure complete coverage of the target A minimum spray volume of 15 gals /A and 30-60 psi at the nozzle tip is recommended On large weeds and/or dense foliage use 60 psi and a minimum of 20 gals /A to ensure coverage of weed foliage~~

The use of flat fan nozzles will result in the most effective postemergence application of FOMESAFEN 22 1% SC The sprayer must be calibrated to provide the proper volume and rate per acre In addition the boom and nozzle height must be adjusted to provide complete coverage of target weeds

DO NOT USE FLOOD TYPE OR OTHER SPRAY NOZZLES WHICH DELIVER COARSE LARGE DROPLET SPRAYS

BAND APPLICATIONS

Thorough weed coverage is important for postemergent control Best coverage is obtained with a minimum of two nozzles one directed to each side of the planted row Application with a single nozzle directed over the top of the row is not recommended for postemergence applications but is suitable for preemergence applications Cultivation of untreated areas may be needed following band applications When making postemergence band applications and cultivating in the same operation position nozzles in advance of the cultivation device This will reduce dust in the spray area Dust can intercept spray reducing weed coverage resulting in less than adequate weed control

Calculate the amount of herbicide and water volume needed for postemergence band treatment by the following formulas

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{broadcast rate per acre} = \text{Band herbicide rate per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{broadcast volume per acre} = \text{Band water volume per acre}$$

AERIAL APPLICATION

Use sufficient spray volume and pressure to ensure complete coverage of the target. A minimum of 5 gals /A of spray mixture should be applied with a maximum of 40 PSI pressure. When broadleaf weed foliage is dense use a minimum of 10 gals /A to ensure coverage of weed foliage.

Cultivation

Cultivation prior to application is not recommended. Cultivation may put weeds under stress reducing weed control. Timely cultivation 1-3 weeks after applying FOMESAFEN 22 1% SC may assist weed control.

Rainfastness

FOMESAFEN 22 1% SC requires a 1-hour rain-free period for best results when applied postemergence.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM EXCEPT CENTER PIVOT SYSTEMS

CENTER PIVOT IRRIGATION APPLICATION

FOMESAFEN 22 1% SC alone or in tank mixture with other herbicides on this label, which are registered for center pivot application, may be applied in irrigation water preemergence (after planting but before weeds or crop emerge) at rates recommended on this label. FOMESAFEN 22 1% SC also may be applied postemergence to the crop and preemergence to weeds in crops where postemergence applications are allowed on this label. Follow all restrictions (height, timing, rate, etc) to avoid illegal residues. Apply this product only through a center pivot irrigation system. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension specialists, equipment manufacturers, or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Operating Instructions

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distributions adversely affected.

- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump or piston pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock
- Do not apply when wind speed favors drift beyond the area intended for treatment
- Prepare a mixture with a minimum of 1 part water to 1 part herbicide(s) and inject this mixture into the center pivot system. Injecting a larger volume of a more dilute mixture per hour will usually provide more accurate calibration of equipment. Maintain sufficient agitation to keep the herbicide in suspension.
- Meter into irrigation water during entire period of water application
- Apply in 1/2 - 1 inch of water. Use the lower water volume (1/2 inch) on coarser soils and the higher volume (1 inch) on fine textured soils. More than 1 inch of water at application may reduce weed control by moving the herbicide below the effective zone in the soil.

Precaution for center pivot applications. Where sprinkler distribution patterns do not overlap sufficiently unacceptable weed control may result. Where sprinkler distribution patterns overlap excessively, crop injury may result.

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in patient clinics, nursing homes or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses.

Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive area. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other locations affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words 'KEEP OUT', followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

Posting required for chemigation does not replace other posting and reentry interval requirements for farm worker safety.

Specific Instructions for Public Water Systems

- 1 Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- 2 Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- 3 The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

- 4 The pesticide injection pipeline must contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down
- 5 The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected
- 6 Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock
- 7 Do not apply when wind speed favors drift beyond the area intended for treatment

PRECAUTIONS & RESTRICTIONS

- A maximum of 1.6 pts of FOMESAFEN 22.1% SC (or a maximum of 0.375 lbs a /A of fomesafen from any product containing fomesafen) may be applied per acre per year in Region 1 (see Regional Use Map)
- A maximum of 1.6 pts of FOMESAFEN 22.1% SC (or a maximum of 0.375 lbs a /A of fomesafen from any product containing fomesafen) may be applied per acre in alternate years in Region 2 (see Regional Map)
- A maximum of 1.3 pts of FOMESAFEN 22.1% SC (or a maximum of 0.313 lbs a /A of fomesafen from any product containing fomesafen) may be applied per acre in alternate years in Region 3 (see Regional Map)
- A maximum of 1 pt of FOMESAFEN 22.1% SC (or a maximum of 0.25 lbs a /A of fomesafen from any product containing fomesafen) may be applied per acre in alternate years in Region 4 (see Regional Map)
- A maximum of 0.75 pt of FOMESAFEN 22.1% SC (or a maximum of 0.1875 lbs a /A of fomesafen from any product containing fomesafen) may be applied per acre in alternate years in Region 5 (see Regional Map)
- Thoroughly clean the spray system with water and a commercial tank cleaner before and after each use
- Tank mixes of FOMESAFEN 22.1% SC with other pesticides, fertilizers or any other additives except as specified on this label or other approved Albaugh Inc supplemental labels may result in tank mix incompatibility, unsatisfactory performance and/or unsatisfactory crop injury
- Apply postemergence to actively growing weeds. Avoid applying FOMESAFEN 22.1% SC to weeds or soybeans which are under stress from moisture, temperature, low soil fertility, mechanical or chemical injury, as reduced weed control and/or increased crop injury may result
- Avoid overlapping spray swaths, as injury may occur to rotational crops
- To provide adequate spray coverage, ground speed must not exceed 10 MPH during application
- Do not graze treated areas or harvest for forage or hay
- Do not apply within 45 days of soybean harvest

Replanting

If replanting is necessary in fields previously treated with FOMESAFEN 22 1% SC the field may be replanted to cotton dry beans **potatoes** snap beans or soybeans **During replanting, a minimum of tillage is recommended to preserve the herbicide barrier for effective weed control** Do not apply a second application of FOMESAFEN 22 1% SC or other fomesafen containing product as crop injury or illegal residues may occur in harvested crops If tank mix combinations were used refer to product labels for any additional replanting instructions

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying FOMESAFEN 22 1% SC at recommended rates in soybeans

Crops To Be Planted	Minimum Rotation Interval (Months After Last FOMESAFEN 22 1% SC Application)
Cotton dry beans potatoes snap beans and soybeans	0
Small grains such as wheat barley rye, peppers (transplanted), tomatoes (transplanted)	4
Corn peanuts peas rice seed corn	10
To avoid injury do not plant alfalfa sunflowers sugar beets sorghum or any other crop within	18

Use 12 month minimum rotation interval for popcorn in the states of Ohio Kentucky Illinois Indiana Iowa and Region 4 when applied at a rate of 1.0 pt/A or more

Use 18 month minimum rotation interval for sweet corn in the states of Connecticut Maine Massachusetts New Hampshire New York Rhode Island Vermont and Region 5


Sorghum may be planted back after 10 months in Region 1

Do not graze rotated small grain crops or harvest forage or straw for livestock

FOMESAFEN 22 1% SC USE RATES AND WEEDS CONTROLLED


REFER TO MAP FOR DEFINITION OF SPECIFIED
GEOGRAPHIC REGIONS

FOMESAFEN 22 1% SC REGIONAL USE MAP



REGION 1
(Maximum Rate 1.6 pts /A per year)

REGION 1 Includes the following states or portion of states where FOMESAFEN 22.1% SC may be applied: Alabama, Arkansas, Florida (except Miami Dade County), Georgia, Louisiana, Mississippi, Missouri (Counties of Bollinger, Butler, Cape Girardeau, Dunklin, Madison, Mississippi, New Madrid, Pemiscot, Perry, Ripley, Scott, Stoddard and Wayne), North Carolina, Oklahoma (East of U.S. Highway 75 and East of Indian Nation Parkway), South Carolina, Tennessee and Texas (all areas East of U.S. Highway 77 to State Road 239 including all of Calhoun County).




REGION 2
(Maximum Rate 1.6 pts /A alternate years)

REGION 2 Includes the following states or portion of states where FOMESAFEN 22.1% SC may be applied: Delaware, Kentucky, Maryland, Virginia, and West Virginia; South of Interstate 70 in the following states: Illinois, Indiana, and Ohio; and in Pennsylvania (all areas South of Interstate 80 to the intersection of U.S. Highway 15 and East of U.S. Highway 15 and U.S. Highway 522).




REGION 3
(Maximum Rate 1.3 pts /A alternate years)

REGION 3 Includes the following states or portion of states where FOMESAFEN 22.1% SC may be applied Connecticut Iowa Maine Massachusetts Missouri (all counties except for those listed in Region 1) New Hampshire New Jersey New York Pennsylvania (all areas except those listed in Region 2) Rhode Island Vermont Wisconsin (South of U.S. Highway 18 between Prairie Du Chien and Madison and South of Interstate 94 between Madison and Milwaukee) and North of Interstate 70 in the following states Illinois Indiana and Ohio



REGION 4
(Maximum Rate 1 pt /A alternate years)

REGION 4 Includes the following states or portion of states where FOMESAFEN 22 1% SC may be applied Kansas (all counties East of or intersected by U S Highway 281) Michigan (Southern Peninsula) Minnesota (all areas South of Interstate 94) Nebraska (all counties East of or intersected by U S Highway 281) and Wisconsin (all areas except those in Region 3 South of Interstate 94 from Minnesota state line to Eau Claire and South of U S Highway 29 from Eau Claire to Green Bay plus Barron Chippewa Clark Door Dunn Eau Claire Kewaunee Marathon Menominee Oconto Polk Shawano and St Croix counties) The following counties are excluded Adams Marquette Portage Waupaca Waushara and Wood) North Dakota (all areas East of Interstate 29 from Fargo South to the South Dakota state line) South Dakota (all areas East of Interstate 29 from the North Dakota state line to Watertown all areas East of Highway 81 from Watertown to Madison and all areas East and South of State Road 34 and U S Highway 281 to the Nebraska state line)



REGION 5
(Maximum Rate 0.75 pts /A alternate years)

REGION 5 Includes the following states or portion of states where FOMESAFEN 22.1% SC may be applied North Dakota (all areas East of U S Highway 281 except those areas in Region 4) South Dakota (all areas East of U S Highway 281 except those areas in Region 4) and Minnesota (all areas South of U S Highway 2 except those areas in Region 4)



WEEDS CONTROLLED

Table 1 Weeds controlled or partially controlled* by preemergence activity of FOMESAFEN 22 1% SC at 1 to 1 6 pints/A¹

<u>Broadleaf Weeds Controlled</u>	<u>Soil Texture</u>	<u>Organic Matter</u>
<u>Amaranth Palmer</u>	<u>All soil types</u>	<u>Up to 5%</u>
<u>Croton, topic²</u>		
<u>Eclipta</u>		
<u>Galinsoga spp</u>		
<u>Lambsquarter, common</u>		
<u>Morningglory, smallflower</u>		
<u>Nightshade, black</u>		
<u>Nightshade, Eastern black</u>		
<u>Pigweed, redroot</u>		
<u>Pigweed, smooth</u>		
<u>Poinsetti, wild</u>		
<u>Purslane, common</u>		
<u>Ragweed, common²</u>		
<u>Sida, prickly²</u>		
<u>Starbur, bristly</u>		
<u>Broadleaf Weeds Partially Controlled*</u>		
<u>Anoda, spurred</u>		
<u>Cocklebur, common</u>		
<u>Morningglory, entireleaf</u>		
<u>Morningglory, ivyleaf</u>		
<u>Morningglory, pitted</u>		
<u>Morningglory, red/scarlet</u>		
<u>Morningglory, tall</u>		
<u>Nightshade, hairy</u>		
<u>Ragweed, giant</u>		
<u>Waterhemp, common</u>		
<u>Sedges Partially Controlled*</u>		
<u>Nutsedge, yellow</u>		

***Partial control means significant activity but not always at a level considered acceptable for commercial weed control**

¹Use the higher end of the rate range when heavy weed populations are anticipated

²Rates less than 1 6 pints/A will provide only partial control of this weed

Table 2 Application Rates For Weed Growth Stages

Weed	FOMESAFEN 22 1% SC (pt /A)			
	Maximum Growth Stage Controlled At			
	0.75 pt /A No of True Leaves	1 pt /A No of True Leaves	1.3 pts /A No of True Leaves	1.6 pts /A No of True Leaves
Anoda, Spurred				2
Balloonvine			2 ^c	2
Carpetweed		6 Diameter Size	Multi leaf 6 diameter	Unlimited Size
Citron (Wild Watermelon)		2	2	4
Cocklebur, Common ^{a b}			2	4
Copperleaf, Hophornbeam		2	2	4
Copperleaf, Virginia		2	2	4
Crotalaria, Showy		4	4	6
Croton, Tropic		2	2	4
Cucumber, Volunteer		4	4	6
Eclipta		2	2	4
Groundcherry, Cutleaf		4	4	6
Hemp ^b			4	6
Horsenettle ^b		2 ^c	3 ^c	4 ^c
Jimsonweed	2	4	6	8
Ladysthumb		2	2	4
Lambsquarters, Common ^c		2	2	2
Mexicanweed		2 ^c	2 ^c	2
Morningglory				
Cypressvine		4	4	6
Entireleaf var	2 ^c	2	2	4
Ivyleaf	2 ^c	2	2	4
Purple Moonflower		2	4	4
Red (Scarlet)		2	2	4
Smallflower		2	2	4
Pitted (Smallwhite)		4	4	4
Tall (Common)	2 ^c	2	2	3
Palmleaf (Willowleaf)		2	2	4
Mustard, Wild	2	4	6	8
Nightshade, Black	2	4	4	4
Nutsedge, Yellow				Suppression Only
Pigweed, spp				
Amaranth, Palmer	2 ^c	4	4	6
Amaranth, Spiny	2 ^c	2	2	4
Redroot	2 ^c	4	6	6
Smooth	2 ^c	4	4	6
Poinsettia, Wild				3
Purslane, Common		Multi Leaf 6 Diameter	Multi Leaf 6 Diameter	Multi Leaf 8 Diameter

Weed	FOMESAFEN 22 1% SC (pt /A)			
	Maximum Growth Stage Controlled At			
	0.75 pt /A No of True Leaves	1 pt /A No of True Leaves	1.3 pts /A No of True Leaves	1.6 pts /A No of True Leaves
Pusley, Florida		2	2	4
Ragweed, Common	2	4	4	6
Ragweed, Giant ^b			4	4
Redweed				3 ^c
Sesbania, Hemp		6	6	12
Sicklepod				Cotyledon ^c
Sida, Prickly				Cotyledon ^c
Smartweed, Pennsylvania	2 ^c	4	4	6
Smellmelon				2
Spurge, Prostrate				1 Diameter
Spurge, Spotted				2 ^c
Starbur, Bristly		2	2	4
Sunflower, Common				2
Velvetleaf ^b			2	4
Venice Mallow	2	4	4	6
Witchweed		Multi Leaf Up to 7	Multi Leaf Up to 7	Multi Leaf Up to 10
Waterhemp, common	2 ^c	2	2	6
Waterhemp, Tall	2 ^c	2	2	4
Yellow Rocket	2	4	6	6

*Partial control means significant activity but not always at a level considered acceptable for commercial weed control

Do not apply in cotyledon stage

^b For effective control of this weed it is necessary to use 1% MSO and 2.5% UAN v/v as adjuvant in Regions 2 and 3 (soybeans only)

Partial control

SPECIAL USE DIRECTIONS FOR ADDITIONAL WEED PROBLEMS

Suppression **Partial Control*** of Annual Grasses

The grasses listed below may be suppressed by postemergence applications and controlled or suppressed by preemergence applications of FOMESAFEN 22 1% SC at 1 – 1.6 pts /A. Consult Use Rate Table for maximum rate in each region. For full season broad spectrum annual grass control Fusilade® DX or Fusion® herbicide should be used alone or in tank mix with FOMESAFEN 22 1% SC. Consult tank mix section.

- Barnyardgrass
- Broadleaf Signalgrass
- Crabgrass
- Foxtail
 - Giant
 - Green
 - Yellow
- Goosegrass
- Johnsongrass Seedling
- Panicum Fall
- Panicum Texas

Suppression **Partial Control*** of Perennial Weeds

Use of FOMESAFEN 22 1% SC at postemergence rates of 1 - 1.6 pts /A will aid in suppressing the above ground portions of the weeds listed below until crop canopy can assist in suppression. Perennial weeds continue to regrow from underground rootstocks even if above ground foliage is temporarily controlled or retarded. Even though FOMESAFEN 22 1% SC and crop competition can suppress perennial weeds for a growing season, the rootstocks will continue to live and reestablishment will occur in subsequent years.

- Milkweed Climbing
- Milkweed Honeyvine
- Bindweed Field
- Bindweed Hedge
- Trumpet creeper

***Partial control means significant activity but not always at a level considered acceptable for commercial weed control**

COTTON USE DIRECTIONS

Preemergence Application

Apply Fomesafen 22 1% SC pre emergence at 1 - 1.6 pts /A in cotton in Region 1 for control or partial control of the weeds listed in Table 1 of the container label. Apply as a pre emergence treatment only to coarse textured soils (sandy loam, loamy sand, sandy clay loam).

Do not apply as a pre emergence treatment to medium or fine textured soils as crop injury will likely occur.

To broaden the weed control spectrum, Fomesafen 22 1% SC may be tank mixed with other preemergence herbicides such as Caparol®, Cotoran®, Direx®, Karmex®, Solicam® or Staple®.

For control of emerged weeds, Fomesafen 22 1% SC may be tank mixed with a burndown herbicide such as Gramoxone Inteon™ or glyphosate brands (such as Touchdown®, Roundup®) labeled in cotton. In reduced tillage plantings, Fomesafen 22 1% SC can be applied up to 14 days prior to planting or at planting with a burndown herbicide. Refer to the tank mix partner label for use directions, restrictions and limitations. The most restrictive product labeling applies.

Cotton plants are tolerant to pre emergence applications of Fomesafen 22 1% SC when applied at recommended rates and to coarse textured soil types. Some crinkling or spotting of cotton foliage or stunting may occur, especially if heavy rainfall occurs during or soon after cotton emergence, but cotton plants normally outgrow these effects and develop normally.

Cotton foliage is not tolerant to Fomesafen 22 1% SC. Do not apply Fomesafen 22 1% SC over the top of emerged cotton as unacceptable cotton injury will occur.

Post Directed Application

Apply Fomesafen 22 1% SC in emerged cotton as a post directed treatment using precision post directed, hooded or shielded application equipment to provide complete coverage of emerged weeds. Apply Fomesafen 22 1% SC at 1 - 1.6 pints per acre in a minimum of 10 gallons spray solution per acre.

Post directed applications of Fomesafen 22 1% SC will provide contact control of labeled emerged weeds and residual pre emergence control of labeled weeds (once activated by rainfall or irrigation) See container label sections for a list of weeds controlled recommended application rates weed growth stages and application directions

Fomesafen 22 1% SC should be applied with a non ionic surfactant at 0.25 to 0.5% v/v or crop oil concentrate at 1% v/v to emerged weeds Do not add liquid nitrogen (28% or similar) to Fomesafen 22 1% SC or Fomesafen 22 1% SC tank mixes in cotton

To broaden the weed control spectrum post directed applications of Fomesafen 22 1% SC may be tank mixed with other labeled post directed herbicides such as Caparol Direx Dual MAGNUM® Envoke® Karmex Layby™ Pro Sequence® or Suprend® When applied with hooded or shielded sprayers Fomesafen 22 1% SC and Fomesafen 22 1% SC tank mixes may be applied with burndown products such as Gramoxone Inteon Sequence or glyphosate brands (such as Touchdown Roundup) labeled for in crop application in cotton Refer to the tank mix partner label for use directions restrictions and limitations The most restrictive product labeling applies

Cotton foliage is not tolerant to Fomesafen 22 1% SC applications Avoid contact to cotton foliage as unacceptable injury will occur Application equipment should be calibrated (spray pressure nozzle type and configuration and orifice size) to avoid fine spray droplets contacting green cotton stems and foliage

Post Directed Application Timing In Cotton

Fomesafen 22 1% SC may be applied to cotton at least 6 inches in height through lay by as a post directed application All post directed applications should avoid spray contact with any green non barked parts of the cotton plant or foliage as unacceptable injury will occur Follow the application timing recommendations below for post directed applications in cotton

Shield And Hooded Applications

Make a precision post directed Fomesafen 22 1% SC application to the base of the cotton plant avoiding contact with the cotton stem or foliage when cotton is at least 6 inches in height to avoid cotton injury Use only hooded or shielded spray equipment to apply Fomesafen 22 1% SC in cotton that is 6 inches to 12 inches in height Adjust nozzles to provide full coverage of emerged target weeds

Layby Applications

Make a post directed FOMESAFEN 22 1% SC application to the base of the cotton plant avoiding contact with any non barked portion of the cotton plant or foliage Use precision post directed equipment or hooded or shielded sprayers on cotton that has developed a minimum of 4 inches of brown bark through layby Application equipment should be configured to provide full coverage of emerged target weeds

RESTRICTIONS FOR USE ON COTTON

- Do not apply Fomesafen 22 1% SC later than 70 days before harvest
- Do not apply more than 1.6 pints per acre of Fomesafen 22 1% SC in any year
- Do not make more than one application of Fomesafen 22 1% SC per year
 - If two consecutive year applications are made allow a 2 year interval before another application

Special Use Directions for the Suppression of Woollyleaf Bursage (Lakeweed) Ambrosia gray/ in Texas

Apply Fomesafen 22 1% SC to cultivated areas of cropland in the fall or spring as a spot treatment at a rate of 1.6 pints per acre and incorporate to a depth of 2-3 inches for suppression of woollyleaf bursage Applications should be made with ground equipment

The use of adjuvants as specified under the Spray Additives section will significantly improve the initial burndown of any emerged woollyleaf bursage but this effect is only temporary Therefore an adjuvant may be used if desired but is not necessary

Significant suppression may not be seen until 6 8 months after application but should then continue for at least 2 years after application Cotton or soybeans may be planted in treated areas Under certain conditions significant damage may occur to cotton planted within 18 months of application A 3 year interval from last application to planting is required for all other crops

DRY BEANS AND SNAP BEANS

Preplant Surface and Preemergence Application

Apply FOMESAFEN 22 1% SC as a preplant surface or preemergence application in Regions 1 2 3 and 4 only for control or partial control of the weeds listed in Table 1 FOMESAFEN 22 1% SC can be applied alone or tank mixed or followed sequentially with other labeled dry bean or snap bean herbicides to broaden the weed control spectrum or control newly emerged weeds Refer to the Tank Mix and Sequential Application section for additional information

NOTE Treated soil that is splashed onto newly emerged seedings may result in temporary crop injury but plants normally outgrow these effects and develop normally

Postemergence Application

Apply FOMESAFEN 22 1% SC as a postemergent broadcast application in Regions 1 2 3 4 and 5 for control or partial control of the weeds listed in Table 2 and in the Special Use Directions for Additional Weed Problems section Application rate depends on weed species and growth stage Two applications may be made if necessary but not to exceed the maximum rate specified per geographic region (Refer to map for definition of specified geographic regions) Refer to the Spray Additive section for recommended spray additives Use of crop oil concentrate can improve weed control but may slightly reduce crop tolerance Do not use UAN (28% or similar) or ammonium sulfate on dry beans or snap beans as severe crop injury may occur Apply when dry beans or snap beans have at least one fully expanded trifoliate leaf

FOMESAFEN 22 1% SC can be applied alone or in tank mix with other labeled dry bean or snap bean postemergence herbicides to broaden the weed control spectrum Refer to the Tank Mix and Sequential Application section

Some bronzing crinkling or spotting of dry bean or snap bean leaves may occur following postemergent applications but dry beans and snap beans soon outgrow these effects and develop normally

Tank Mix and Sequential Applications for Dry Beans and Snap Beans

FOMESAFEN 22 1% SC can be used sequentially or in tank mix with the following products

Dry Beans and Snap Beans	Dry Beans Only
Assure II®	Frontier®
Basagran®	Clethodim 2E
Dual MAGNUM	Sonalan®
Eptam®	
Poast®	
Prowl®	
Imazethapyr 2SC	
Raptor®	
Trifluralin 4E	

Under certain conditions the mixture of FOMESAFEN 22 1% SC with one or more of the above mentioned broadleaf herbicides may cause a reduction in activity of any post emergence grass herbicide in the mixture

For sequential applications allow 2 3 days after the application of the post emergence grass herbicide before applying FOMESAFEN 22 1% SC or FOMESAFEN 22 1% SC mixtures Where FOMESAFEN 22 1% SC or the FOMESAFEN 22 1% SC mixture is applied first apply the grass herbicide when the grass weeds begin to develop new leaves (generally around 7 days)

NOTE Tank mix applications can result in increased crop injury as compared to either product used alone

Always read and follow the recommendations restrictions and limitations for all products whether used alone sequentially or in a tank mix The most restrictive labeling of any product used applies

Restrictions Dry Beans and Snap Beans

- Refer to FOMESAFEN 22 1% SC Regional Use Map for the maximum rate of FOMESAFEN 22 1% SC (or other fomesafen containing products) that may be applied in each geographic region
- Do not apply to any field in Regions 2 3 4 or 5 more than once every two years
- For snap beans Do not exceed 1 6 pints of FOMESAFEN 22 1% SC per acre in any one year and also adhere to the maximum rate that may be applied in each geographic region (refer to the FOMESAFEN 22 1% SC Regional Use Map) Do not graze treated areas or harvest for forage or hay Do not utilize hay or straw for animal feed or bedding Do not apply within 30 days of harvest
- For dry beans Do not exceed 1 6 pints of FOMESAFEN 22 1% SC per acre in any one year and also adhere to the maximum rate that may be applied in each geographic region (refer to the FOMESAFEN 22 1% SC Regional Use Map) Do not graze animals on green forage or stubble Do not utilize hay or straw for animal feed or bedding Do not apply within 45 days of harvest

POTATOES

Apply FOMESAFEN 22 1% SC at 1 pt /A as a broadcast preemergence application after planting but before potato emergence for control or partial control of weeds listed in Table 1 Effectiveness will be reduced if later cultural practices expose untreated soil For application by center pivot irrigation, see the Center Pivot Irrigation Application section of this label

Note Potato varieties may vary in their response to FOMESAFEN 22 1% SC When using FOMESAFEN 22 1% SC for the first time on a particular variety, always determine crop tolerance before using

Tank Mixtures With Other Products Registered for Use in Potatoes

For preemergence applications in potatoes, FOMESAFEN 22 1% SC may be tank mixed with other pesticide products registered for use in this way and timing in potatoes Follow the directions for use, observe the stated precautions, and abide by the limitations and restrictions on the most restrictive of the product labels If you have no previous experience mixing these products under your conditions, perform a compatibility test before attempting large scale mixing (see Tank Mix Compatibility Test section of this label)

Product Use Restrictions – Potatoes

- **Do not exceed 1 pt /A of FOMESAFEN 22 1% SC per season Refer to Regional Use Map for the maximum rate of FOMESAFEN 22 1% SC (or other fomesafen containing products) that may be applied per year or alternate year in each geographic region**
- **Do not harvest potatoes treated with FOMESAFEN 22 1% SC within 70 days of application**
- **Do not apply FOMESAFEN 22 1% SC to sweet potatoes or yams**
- **Do not apply FOMESAFEN 22 1% SC as a preplant incorporated application in potatoes or crop injury may occur**
- **Do not apply to emerged potato plants or severe crop injury will occur**

SOYBEANS

Preplant Surface and Preemergence Application

Apply FOMESAFEN 22 1% SC as a preplant surface or preemergence application in Regions 1, 2, 3, and 4 only for control or partial control of the weeds listed in Table 1 FOMESAFEN 22 1% SC can be applied alone or tank mixed or followed sequentially with other labeled soybean herbicides to broaden the weed control spectrum or control newly emerged weeds Refer to the Tank Mix and Sequential Application section for additional information

For control of emerged weeds, FOMESAFEN 22 1% SC may be tank mixed with a burndown herbicide such as Gramoxone Inteon or glyphosate brands (such as Touchdown or Roundup) labeled in soybeans In reduced tillage plantings, FOMESAFEN 22 1% SC can be applied up to 14 days prior to planting or at planting with a burndown herbicide

Postemergence Application

Apply FOMESAFEN 22 1% SC as a postemergence broadcast application in Regions 1, 2, 3, 4 and 5 for control or partial control of weeds listed in Table 2 and in the Special Use Directions For Additional Weed Problems section Application rate depends on weed species and growth stage Refer to the Spray Additive section for recommended spray additives To enhance postemergence control of susceptible broadleaf weeds (soybeans only) in Regions 2, 3, 4 and 5 (see Regional Use Map), FOMESAFEN 22 1% SC can be used with a minimum of 2 5% liquid nitrogen (28% or similar) or a minimum of 10 pounds ammonium sulfate per 100 gallons of spray volume

FOMESAFEN 22 1% SC can be applied alone or in combination with other labeled soybean postemergence herbicides to broaden the weed control spectrum Refer to the Tank Mix and Sequential Application section

Some bronzing, crinkling or spotting of soybean leaves may occur following postemergent applications, but soybeans soon outgrow these effects and develop normally

TANK MIX AND SEQUENTIAL APPLICATIONS FOR SOYBEANS

FOMESAFEN 22 1% SC can be used sequentially or in tank mix with one or more of the following products Assure II® Basagran® Butyrac® Classic® FirstRate® Fusilade DX Fusion Ignite® Glyphosate (such as Touchdown® Roundup® Glyphomax™) Gramoxone® Inteon Harmony® Foast® Poast Plus® Pursuit® Raptor® Resource® Scepter® Select® and Synchrony® STS®

Under certain conditions the mixture of FOMESAFEN 22 1% SC with one or more of the above mentioned broadleaf herbicides may cause a reduction in activity of any postemergence grass herbicide in the mixture

For sequential applications allow 2-3 days after the application of the grass herbicide before applying FOMESAFEN 22 1% SC or FOMESAFEN 22 1% SC mixtures. Where FOMESAFEN 22 1% SC or the FOMESAFEN 22 1% SC mixture is applied first, apply the grass herbicide when grass weeds begin to develop new leaves (generally around 7 days).

- Tank mix applications can result in increased crop injury as compared to either product used alone
- Do not exceed 1 fl. oz. of Butyrac per acre in mixture with FOMESAFEN 22 1% SC
- Do not exceed 0.25 oz./A of Synchrony STS herbicide in the tank with labeled rates of FOMESAFEN 22 1% SC on non STS varieties. This tank mix can be applied postemergence to any soybean variety for additional broadleaf weed control. Refer to the Synchrony STS label for more information and crop rotation restrictions.
- Always read and follow the recommendations, restrictions and limitations for all products whether used alone, sequentially or in a tank mix. The most restrictive labeling of any product used applies.

GLYPHOSATE TOLERANT SOYBEAN TANK MIXES

FOMESAFEN 22 1% SC at 6-12 oz./A can be tank mixed with glyphosate products (such as Touchdown or Roundup) that are labeled for glyphosate tolerant soybeans for improved postemergence control of many weeds such as morning glory spp., hemp sesbania, waterhemp, and black nightshade which are known to have tolerance to glyphosate but are susceptible to FOMESAFEN 22 1% SC.

FOLLOW THE RECOMMENDATIONS ON THE GLYPHOSATE PRODUCT LABEL FOR THE USE OF SPRAY ADDITIVES IN THIS TANK MIX.

Do not allow this tank mix to move off target as contact by even minute quantities can cause severe damage or death to any non-target vegetation.

Note: Postemergence application of this tank mix on soybean varieties which do not contain the glyphosate tolerant gene will result in severe crop injury or death of the soybean crop. Always read and follow the recommendations, restrictions and limitations for all products used. The most restrictive labeling of any product applies.

Product Use Restrictions – Soybeans

Refer to Regional Use Map for the maximum rate of FOMESAFEN 22 1% SC (or other fomesafen containing products) that may be applied in each geographic region. Do not apply to any field in Region 2, 3, 4 or 5 more than once every two years.

Do not exceed 1.6 pints of FOMESAFEN 22 1% SC per acre in one year and also adhere to the maximum rate that may be applied in each geographic region (refer to the Regional Use Map).

Do not graze treated areas or harvest for forage or hay.

Do not apply within 45 days of harvest.

AERIAL SPRAY DRIFT MANAGEMENT ADVISORY

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations they should be observed.

The applicator should be familiar with and take into account the information covered in the AERIAL DRIFT REDUCTION ADVISORY.

AERIAL DRIFT REDUCTION ADVISORY INFORMATION

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (See Wind, Temperature and Humidity and Temperature Inversion sections of this label).

CONTROLLING DROPLET SIZE

- Volume: Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure: Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of Nozzles: Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation: Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type: Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 ft above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator should compensate for this displacement by adjusting

the path of the aircraft upwind Swath adjustment distance should increase with increasing drift potential (higher wind smaller drops etc)

WIND

Drift potential is lowest between winds speeds of 2 10 mph However many factors including droplet size and equipment type determine drift potential at any given speed Application should be avoided below 2 mph due to variable wind direction and high inversion potential NOTE Local terrain can influence wind patterns Every applicator should be familiar with local wind patterns and how they affect spray drift

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity set up equipment to produce larger droplets to compensate for evaporation Droplet evaporation is most severe when conditions are both hot and dry

TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high Temperature inversions restrict vertical air mixing which causes small suspended droplets to remain in a concentrated cloud This cloud can move in unpredictable directions due to the light variable winds common during inversions Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind They begin to form as the sun sets and often continue into the morning Their presence can be indicated by ground fog however if fog is not present inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion while smoke that moves upward and rapidly dissipates indicates good vertical air mixing

SENSITIVE AREAS

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e g residential areas bodies of water known habitat for threatened or endangered species nontarget crops) is minimal (e g when wind is blowing away from the sensitive areas)

APPENDIX

Scientific names are listed for those weeds referred to in the FOMESAFEN 22 1% SC label

COMMON NAME	SCIENTIFIC NAME
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, Spiny	<i>Amaranthus spinosus</i>
Anoda, Spurred	<i>Anoda cristata</i>
Balloonvine	<i>Cadiospermum halicacabum</i>
Barnyardgrass	<i>Echinochloa crus galli</i>
Bindweed, Field	<i>Convolvulus arvensis</i>
Bindweed, Hedge	<i>Calystegia sepium</i>
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>
Carpetweed	<i>Mollugo verticillata</i>
Citron (Wild Watermelon)	<i>Citrullus vulgaris</i>
Cocklebur, Common	<i>Xanthium strumarium</i>
Copperleaf, Hophornbeam	<i>Acalypha ostryifolia</i>
Copperleaf, Virginia	<i>Acalypha virginica</i>
Crabgrass	<i>Digitaria spp</i>
Crotalaria, Showy	<i>Crotalaria spectabilis</i>
Croton, Tropic	<i>Croton glandulosus</i>
Cucumber, Volunteer	<i>Cucumis sativas</i>

COMMON NAME	SCIENTIFIC NAME
Eclipta	Eclipta prostrate
Foxtail, Giant	Setaria faberi
Foxtail, Green	Setaria viridis
Foxtail, Yellow	Setaria glauca
Goosegrass	Eleusine indica
Groundcherry, Cutleaf	Physalis angulata
Hemp	Cannabis sativa
Horsenettle	Solanum carolinense
Jimsonweed	Datura stramonium
Johnsongrass, Seedling	Sorghum halepense
Ladysthumb	Polygonum persicaria
Lambsquarters, Common	Chenopodium album
Mexicanweed	Caperonia castanifolia
Milkweed, Climbing	Sarcostemma cyanchoides
Milkweed, Honeyvine	Ampelamus albidus
Morningglory, Cypressvine	Ipomoeaquamoclit
Entireleaf	Ipomoea hederacea var integriuscula
Ivyleaf	Ipomoea hederacea var hederacea
Purple Moonflower	Ipomoea turbinata
Red (Scarlet)	Ipomoea coccinea
Smallflower	Jacquemontia tamnifolia
Pitted (Smallwhite)	Ipomoea lacunose
Tall (Common)	Ipomoea purpurea
Palmleaf (Willowleaf)	Ipomoea wrightii
Mustard, Wild	Brassica kaber
Nightshade, Black	Solanum nigrum
Nutsedge, Yellow	Cyperus esculentus
Panicum, Fall	Panicum dichotomiflorum
Panicum, Texas	Panicum texanum
Pigweed, Redroot	Amaranthus retroflexus
Pigweed, Smooth	Amaranthus hybridus
Poinsettia, Wild	Euphorbia heterophylla
Purslane, Common	Portulaca oleracea
Pusley, Florida	Richardia scabra
Ragweed, Common	Ambrosia artemisiifolia
Ragweed, Giant	Ambrosia trifida
Redweed	Melochia corchorifolia
Sesbania, Hemp	Sesbania exaltata
Sicklepod	Cassia obtusifolia
Sida, Prickly	Sida spinosa
Smartweed, Pennsylvania	Polygonum pennsylvanicum
Smellmelon	Cucumis melo
Spurge, Prostrate	Euphorbia humistrata
Spurge, Spotted	Euphorbia maculate
Starbur, Bristly	Acanthospermum hispidum
Sunflower, Common	Helianthus annuus
Trumpetcreeper	Campsis radicans
Velvetleaf	Abutilon theophrasti
Venice Mallow	Hibiscus trionum
Waterhemp, Common	Amaranthus rudis

COMMON NAME	SCIENTIFIC NAME
Waterhemp, Tall	Amaranthus tuberculatos
Witchweed	Striga asiatica
Yellow Rocket	Barbarea vulgaris

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

Prohibitions Open dumping is prohibited Do not reuse empty container

PESTICIDE STORAGE Store above 32 F in original containers only If product solidifies return to room temperature and agitate to reconstitute Keep container closed when not in use Do not store near food or feed In case of spill or leak on floor or paved surfaces soak up with sand earth or synthetic absorbent Remove to chemical waste area

PESTICIDE DISPOSAL Pesticide wastes are toxic Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance

Container Handling [Less Than 5 Gallons]

Non refillable container Do not reuse or refill this container Offer for recycling if available Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or mix tank Drain for 10 seconds after the flow begins to drip Fill the container % full with water and recap Shake for 10 seconds Pour rinsate into application equipment or mix tank or store rinsate for later use and disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

Container Handling [For Bulk and Mini Bulk Containers]

Refillable container Refill this container with pesticide only Do not use this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the person refilling To clean container before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allowed by state and local authorities If the container is damaged leaking or obsolete contact Albaugh Inc at 1 800 247 8013

CONTAINER IS NOT SAFE FOR FOOD FEED OR DRINKING WATER

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product If the terms are not acceptable return the product at once unopened and the purchase price will be refunded

The Directions for Use of this product must be followed carefully It is impossible to eliminate all risks inherently associated with the use of this product Crop injury ineffectiveness or other unintended consequences may result because of such factors as manner of use or application weather or crop conditions presence of other materials or

other influencing factors in the use of the product which are beyond the control of ALBAUGH INC or Seller To the extent permitted by applicable law Buyer and User agree to hold ALBAUGH INC and Seller harmless for any claims relating to such factors

ALBAUGH INC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use subject to the inherent risks referred to above when used in accordance with directions under normal use conditions To the extent permitted by applicable law (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or ALBAUGH INC and (2) Buyer and User assume the risk of any such use TO THE EXTENT PERMITTED BY APPLICABLE LAW ALBAUGH INC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL

To the extent permitted by applicable law in no event shall ALBAUGH INC be liable for any incidental consequential or special damages resulting from the use or handling of this product TO THE EXTENT PERMITTED BY APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF ALBAUGH INC AND SELLER FOR ANY AND ALL CLAIMS LOSSES INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY CONTRACT NEGLIGENCE TORT STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR AT THE ELECTION OF ALBAUGH INC OR SELLER THE REPLACEMENT OF THE PRODUCT

ALBAUGH INC and Seller offer this product and Buyer and User accept it subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of ALBAUGH INC

Fusilade® Fusion® Gramoxone® Touchdown® are trademarks of a Syngenta Group Company
Basagran® Poast® Poast Plus® Pursuit® Raptor® and Scepter® trademarks of BASF Ag Products
Assure II® Classic® Harmony® Synchrony® STS® and Viton trademarks of E I DuPont de Nemours & Co Inc
Select® and Resource® trademarks of Valent Chemical Co
Butyrac® trademark of Albaugh Inc
Roundup® trademark of Monsanto Company
FirstRate® and Glyphomax trademark of Dow Agro Sciences
Ignite® trademark of Bayer CropScience

c c
c c