



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave., NW
 Washington, D.C. 20460

EPA Reg. Number: 42750-233
 Date of Issuance: OCT - 3 2011

NOTICE OF PESTICIDE:
 Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance: Conditional

Name of Pesticide Product: 91% MCPA ESTER

Name and Address of Registrant (include ZIP Code):
 Albaugh, Inc.
 PO Box 2127
 Valdosta, GA 31604-2127

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/ reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Add the phrase "EPA Registration No. 42750-233" to your label before you release the product for shipment.

Signature of Approving Official:

 Kathryn V. Montague
 Product Manager (23)
 Herbicide Branch
 Registration Division (7505P)

Date: OCT - 3 2011

page 2

EPA Reg. No. 42750-2339

3. Submit one (1) copy of your final printed labeling before you release the product for shipment.

4. Submit the results of one year storage stability and corrosion characteristic studies to EPA within one year of this date of registration. Submit a hard copy and an electronic copy also.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

91% MCPA ESTER
Selective Herbicide

FOR THE SELECTIVE CONTROL OF BROADLEAF WEEDS IN BARLEY, OATS, RYE, WHEAT, FLAX,
GRASSES, RANGELANDS, PASTURES, AND NON-CROP AREAS.

ACTIVE INGREDIENT:

Isooctyl (2-ethylhexyl) Ester of 2-methyl-4-Chlorophenoxyacetic Acid* 91.0%

OTHER INGREDIENTS: 9.0%

TOTAL: 100.0%

Isomer Specific AOAC Method, Equivalent to: *58.4% 2-Methyl-4-Chlorophenoxyacetic Acid (5.2 lbs. per gal.)

KEEP OUT OF REACH OF CHILDREN

WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a listed en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	


SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 42750-EGG

EPA Est. No. 42750-MO-001

NET CONTENTS: _____

MANUFACTURED BY:
Albaugh, Inc.
Ankeny, IA 50021



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING / AVISO

May be fatal if absorbed through skin. Harmful if swallowed. Do not get in eyes, on skin, or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an ERA chemical-resistance category selection chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant shoes plus socks and,
- Chemical-resistant gloves when mixing, loading, or using any hand-held equipment.

Additional PPE requirements for mixers and loaders supporting aerial application to rangelands, pasture lands, or non-cropland.

These mixers/loaders also must wear: A chemical-resistant apron and a NIOSH-approved dust/mist filter equipped with N, R, P, or HE class filter media. The respirator should have a NIOSH approval number prefix TC-84A. It is recommended that you require that the respirator wearer be fit tested, and trained in use, maintenance, and limitations of the respirator.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)]. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the WPS (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls,
- shoes plus socks,
- chemical-resistant gloves made of any waterproof material, and
- protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people to enter the treated area until sprays have dried.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2 to 10 mph at the application site.

Additional Requirements for Aerial Applications

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications into temperature inversions.

Additional Requirements for Ground Boom Applications.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

EFFECTIVENESS ON WEEDS: This product is effective on a large number of broadleaf weeds. On Mustard, Thistle, and White top or Hoary cress, it has superiority over 2,4-D. The following are some of the weeds killed: Beggar tick, Burcucumber, Burdock, Buttercup, Canada thistle, Carpetweed, Cats ear, Chervil, Cocklebur, Croton, Daisy, Dandelion, Dock, Dragonhead mint, Dwarf nettle, Fanweed, Fat hen, Fennel, Field bindweed, Galinsoga, Goatsbeard, Goldenrod, Halbertleaved salt brush, Hemp nettle, Hoary cress, Honeysuckle, Jimsonweed, Knotweed, Kochia, Lambsquarter (Goosefoot), Marshelder, Mexican poppy, Miner's lettuce, Narrow leaf plantain, Pennycress, Pepperweed, Perennial morningglory, Perennial ragweed, Pigweed, Poison hemlock, Prickly lettuce, Puncturevine, Purslane, Red root, Redstem, Shepherds purse, Sicklepod, Small plantain, Sneezeweed, Sowthistle, Spanish needle, Stinging nettle, Stinkweed, Sunflower, Thornapple, Tree-of-heaven, Vetch, Whitebrush, Wild carrot, Wild gooseberry, Wild jute, Wild marigold, Wild petunia, Wild radish, Wild sage, Witchweed, Yellow charlock-mustard, Yellow daisy, Yellow rocket.

SELECTIVE SPRAYING

NOTE: When using on grains and flax - do not forage or graze dairy and meat animals on treated areas within seven days of slaughter. Also, except for small underseeded grains, use at least 8 to 10 gallons of water per acre for ground application and at least 1 to 5 gallons of water per acre for aerial application.

WHEAT, BARLEY, RYE underseeded with legumes:

Spring planted - not seeded to legumes.

Apply as a water mix spray by ground sprayer or airplane. Use 1/3 to 1 pint of this product per acre for the more susceptible weeds and up to 1.15 pints per acre for less susceptible weeds. Treat when fully tillered grains have reached 3 to 4 leaf stage (8 to 10 inches tall). Do not spray grain in the boot to dough stage.

Fall Planted - Fall application - not seeded to legumes.

The safest time to treat is when Fall planted small grains are fully tillered and have reached 3 to 4 leaf to the boot stage (8 to 10 inches tall). Infestations of broadleaf weeds germinating in Fall and Winter may, however, justify treatment at an earlier stage. Fall planted wheat, barley, or rye can be sprayed with 1/3 to 3/4 pint of this product per acre from the 3 to 4 leaf stage to the boot stage.

Spring application -

Apply in early Spring when annual broadleaf weeds are small and grain is fully tillered but before grain is in the jointed stage.

WHEAT, BARLEY, RYE, OATS

Underseeded with Alfalfa, Birdsfoot Trefoil, Lespedeza, Red and White Clover

For emergency control of serious infestations of Mustard, Yellow rocket, and other susceptible broadleaf weeds, apply 3 to 6 ounces in not more than 1 to 10 gallons of water per acre. Make application after cereal is well tillered (4 to 8 inches tall) and before reaching boot stage. The nurse crop and weeds should provide a protecting canopy which, together with the use of low gallonage applied at low pressure, will reduce the risk of damage to the legumes. Do not apply to small grains underseeded with Vetch or Sweet clover, which are very susceptible. There is also some risk to other legumes if only thinly protected by a canopy. Only one application is permitted per crop season.

OATS:

Spray after oats are fully tillered (4 to 6 inches tall) but before they are in the boot to dough stage. Apply the lower rate of 6 ounces where annual weeds are the problem, the higher rate of 3/4 pint for perennial weeds.

RESTRICTIONS AND LIMITATIONS FOR USE ON BARLEY, OATS, RYE AND WHEAT:

Do not apply more than 1.15 pints/acre per year.

FLAX

Use 3 to 6 ounces per acre. Use lower rate where susceptible weeds such as Mustard are the main problem. Use higher rate when weeds such as Pigweed, Lambsquarters, Stinkweed, Ragweed, Cocklebur are the problem. Apply only when weeds are up and when flax is 2 to 8 inches high and before it comes into bud. Treatment after early bud stage may result in severe damage. If Canada thistle is present, it may be necessary to go as high as 1/2 pint per acre to prevent seed head production. Some injury to flax may result.

RESTRICTIONS AND LIMITATIONS FOR USE ON FLAX:

Do not apply more than 0.375 pints/acre per year.

RANGE AND TIMBER LANDS

For control of Whitebrush - Use 1 quart of product in sufficient water to make 8 gallons of solution per acre.

Apply in Spring or Fall under good moisture conditions, full leaf, before blossoms begin to fall.

GRASSES

Established Lawns, Golf Courses and similar Turf Grasses - Use 3/4 to 2 pints per acre in 10 to 100 gallons of water to give thorough coverage. Use higher rate for White top and Canada thistle. On smaller areas use 3/4 fluid ounce (3 tablespoonfuls) mixed in 3 to 5 gallons of water and apply uniformly over 2,000 square feet. Do not mow within 2 days before or after application. Spring and Fall are best time to treat. Do not use on lawns of creeping grasses, such as bent, except for spot spraying, nor on freshly-seeded turf until the grass has become well established.

RESTRICTIONS AND LIMITATIONS FOR USE ON GOLF COURSES:

Do not apply more than 2.25 pints/acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

ESTABLISHED GRASSLAND AND PASTURES

Use 3/4 to 2 pints per acre in sufficient water (2 to 100 gallons of mixture) in airplane or ground spray application and give thorough coverage. Use higher rate for White top, Canada thistle and other less easy-to-kill weeds; spray perennials in early bud to full bloom stage and regrowth in Fall. Other weeds in Spring or Fall.

RESTRICTIONS FOR ESTABLISHED GRASSLAND AND PASTURES:

Do not apply more than 1.5 lbs ae/acre/per year (2.3 pints of product/acre/year). Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

GRASSES GROWN FOR SEED

Use 3/4 to 1-1/2 pints per acre in 1 to 50 gallons of water by air or ground sprayer application. Use higher rate where weed stands are heavy. In established grasses, apply in Spring before head comes into boot stage and on seedling grass after grass has tillered.

RESTRICTIONS AND LIMITATIONS FOR USE ON GRASSES GROWN FOR SEED:

Do not apply more than 2.25 pints/per acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

NOTE: For weed control in grasses, repeat treatment may be needed for less susceptible weeds. White clover and other legumes may be temporarily injured or killed. In some areas, bent, buffalo, carpet, centipede, dichondra and St. Augustine may also be injured by the treatment.

NON-CROP SPRAYING

Canada thistle, White top and Meadow buttercup. (In Non-Crop Areas such as Roadsides, Fence rows, Rights-of-way and similar places): For spot treatment, use 3 ounces to 3 to 4 gallons of water or 1/2 gallon per acre in 12 to 20 gallons of water to give coverage for most extensive areas. Spray to wet weeds thoroughly when in bud to early bloom and again on Fall regrowth.

Local conditions may affect the use of this chemical. Consult State Agricultural Extension or Experiment Station weed specialists for specific recommendations for local weed problems and for information on possible lower dosages.

RESTRICTIONS AND LIMITATIONS FOR NON-CROP SPRAYING:

Do not apply more than 1.5 lbs ae/acre/per year (2.3 pints of product/ acre/ year). Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days. For spot treatment do not apply more than 3.0 lbs of ae/acre/per year (4.6 pints of product / acre / per year).

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixing with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of the material not in strict accordance with directions given herewith.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always store pesticides in a secured warehouse or storage building, Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal.

CONTAINER DISPOSAL:

Non-refillable Containers 5 Gallons or Less: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Non-refillable containers larger than 5 gallons: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refutable containers larger than 5 gallons: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.