



42750-256

4/8/2014

1/9

 U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7504P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460 (under FIFRA, as amended)	EPA Reg. Number: 42750-256	Date of Issuance: APR - 8 2014
	Unconditional	
	Name of Pesticide Product: Difenoconazole 3L ST	
Name and Address of Registrant (include ZIP Code): Albaugh, Inc. P.O. Box 2127 Valdosta, GA 31604 Attn.: Morris Gaskins		
Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA section 3(c)(5), provided that you:</p> <ol style="list-style-type: none">1. Submit and/or cite all data required for registration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.		
Signature of Approving Official:  Product Manager Name and Number Fungicide Branch, Registration Division (7504P)		Date: APR - 8 2014

Notice of Pesticide Registration
Product Name: Difenoconazole 3L ST
EPA Reg. No. 42750-256
Page 2 of 2

2. Make the following change to the label:
 - a. Change the product registration number to "EPA Reg. No. "42750-256".
3. Submit one copy of the revised final printed label before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

The basic CSF dated 5/7/13 is acceptable.

A copy of the label stamped "Accepted" is enclosed for your records.

Enclosure:

Label stamped "Accepted"
Product Chemistry Review DP 412781

49

EDITOR'S NOTE: 4/8/14 3:34PM revised draft label to remove any ag/turf/ornamental foliar uses for 42750-ELA and retain only seed treatment uses.

DIFENOCONAZOLE 3L ST
Fungicide

A seed treatment for control of diseases of barley, canola, cotton, oats, rye, sweet corn, wheat and triticale

ACTIVE INGREDIENT:

*Difenoconazole: 32.8%
OTHER INGREDIENTS: 67.2%
TOTAL: 100.0%

*CAS No. 119446-68-3
Contains 3.1 lbs. active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
NOTE TO PHYSICIAN: If ingested, induce emesis or lavage stomach. Treat symptomatically. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	


HOT LINE NUMBER For 24 Hour Medical or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), call CHEMTREC toll free at 1-800-424-9300.

EPA Reg. No. 42750-~~ELA~~ 256

EPA Est. No. xxxxxx-xx-xxx

NET CONTENTS: _____

Manufactured By:
ALBAUGH, INC.
ANKENY, IA 50021



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if inhaled or absorbed through skin. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and other aquatic invertebrates. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion.

PHYSICAL OR CHEMICAL HAZARDS

Do not use with or store near any oxidizing or reducing agents.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

STORAGE: Keep container tightly closed when not in use. Store out of the reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty

5/19

the remaining contents into application equipment or a mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not formulate into pesticides used for foliar applications to crops.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

GENERAL INFORMATION

DIFENOCONAZOLE 3L ST is a systemic seed dressing which controls or suppresses certain seed-borne, soil-borne, and early season foliar diseases of wheat, barley, oats, rye, and triticale, and certain seed and seedling diseases of canola, sweet corn, and cotton. Use an EPA-approved dye or colorant that imparts an unnatural color to the seed. For commercial seed treatment only. Do not use for at-plant applications involving hopper boxes, seed boxes, etc.

Mixing Procedures

Apply DIFENOCONAZOLE 3L ST as a water-based slurry through standard slurry or mist-type seed treatment equipment that accurately meters and mixes a liquid flowable seed treatment. The mix tank should be equipped with a mechanical or paddle agitator to ensure uniform blending of the slurry. The equipment must provide uniform coverage of DIFENOCONAZOLE 3L ST on the seed. Uneven seed coverage may not give the desired level of disease control. Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on the operation and calibration of the equipment.

Fill the mix tank 1/4- 1/2 full with clean water (see maximum recommended slurry volume table below). Start agitation of the mix tank. Next, open the container and add the recommended amount of DIFENOCONAZOLE 3L ST into the mix tank, and maintain agitation. Thoroughly blend the mixture while adding the remainder of the required water.

After the mixture appears completely uniform, begin application to the seed. Mix the slurry thoroughly with the seed to provide uniform coverage.

The following table lists the maximum recommended total slurry volumes by crop.

Crop	Maximum fl. oz. slurry per 100 lbs. seed
Wheat, Oats, Rye and Barley	16
Canola	23

6/9

Sweet Corn	8 to 16
Cotton	30

Confirm physical compatibility, seed safety, seed safety in stored seed, and biological efficacy before tank mixing with other seed treatment products.

Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing.

For commercial seed treatment only. Do not use for at-plant applicators involving hopper boxes, seed boxes, etc.

SEED BAG LABEL REQUIREMENT

Federal law requires that bags containing treated seeds shall be labeled with the following information:

"This seed has been treated with difenoconazole fungicide. Do not use for feed, food, or oil purposes. Store away from feeds and foodstuffs. Use with an EPA-approved dye or colorant that imparts an unnatural color to the seed. Green wheat, oats, rye or triticale forage may not be grazed until 55 days after planting. Green barley forage may not be grazed until 30 days after planting. Do not plant any crop other than those on this label within 30 days to fields in which treated seeds were planted. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues remain in ethanol by-products that are used in agronomic practice."

BARLEY

Apply DIFENOCONAZOLE 3L ST to barley seed at the rates given in the following table:

Rate/100 lbs. seed	Diseases Controlled	Diseases Partially Controlled ¹
0.5 fl. oz.	Barley Stripe General Seed Rots ² Covered Smut Pythium Damping Off	Common Root Rot (<i>Cochliobolus</i> spp.) Fusarium Seed Scab Rhizoctonia Root Rot
1.0 fl. oz.	Barley Stripe General Seed Rots ² Fusarium Seed Scab Covered Smut Pythium Damping Off	Take-all Common Root Rot (<i>Cochliobolus</i> spp.) Fusarium Root Rot Fusarium Crown Rot Rhizoctonia Root Rot

¹ Partial control can either mean erratic control from good to poor, or consistent control at a level below that is generally considered acceptable for commercial disease control.

² General seed rots controlled include those caused by saprophytic organisms such as *Penicillium* spp. and *Aspergillus* spp.

CANOLA

Apply DIFENOCONAZOLE 3L ST at a rate of 1.0 fl. oz. of product per 100 lbs. of seed for protection against seed-borne blackleg caused by *Leptosphaeria maculans*. Tank mix DIFENOCONAZOLE 3L ST with seed treatment fungicide Maxim[®] 4FS for broad spectrum protection against other diseases such as Fusarium and Rhizoctonia.

COTTON

Apply DIFENOCONAZOLE 3L ST at a rate of 0.5 to 1.45 fl. oz. of product per 100 lbs. of seed for protection against preemergent and postemergent damping off seedling diseases caused by *Rhizoctonia* spp. and *Fusarium* spp. Use the higher rate where disease is expected to be severe. Always apply in combination with metalaxyl for activity against seed and seedling diseases caused by *Pythium* spp. Follow application and other instructions found on the metalaxyl product label.

SWEET CORN

Apply DIFENOCONAZOLE 3L ST at a rate of 0.5 to 1.24 fl. oz. of product per 100 lbs. of seed for control of die-back in sweet corn caused by *Penicillium* spp., *Fusarium* spp. and *Aspergillus* spp. Confirm physical compatibility, seed safety, seed safety in stored seed, and biological efficacy before tank mixing with other seed treatment products.

WHEAT, OATS, RYE AND TRITICALE

Apply DIFENOCONAZOLE 3L ST to wheat or triticale seed at the rates given in the following table.

For control of damping-off of wheat caused by *Pythium* spp., the recommended rate of DIFENOCONAZOLE 3L ST must be tank mixed with metalaxyl in accordance with the most restrictive label limitations and precautions. Labeled dosage rates must not be exceeded.

Use metalaxyl at labeled rates with the specified rate of DIFENOCONAZOLE 3L ST. Use the higher listed rates of metalaxyl in combination with DIFENOCONAZOLE 3L ST if disease pressure is expected to be severe. Follow application and other instructions found on the metalaxyl label.

Rate/100 lbs. seed	Diseases Controlled	Fall Season Foliar Diseases Controlled ¹	Early Season Control
0.5 fl. oz.	Common Bunt ¹ , Dwarf Bunt, Flag Smut, Seed-borne Septoria, Loose Smut, General Seed Rots ³ , Fusarium Seed Scab		Common Root Rot (<i>Cochliobolus</i> spp.), Rhizoctonia Root Rot
1 fl. oz.	Common Bunt ² , Dwarf Bunt, Loose Smut, Flag Smut, Seed-borne Septoria, General Seed Rots ³ , Fusarium Seed Scab	Fall Season Powdery Mildew, Fall Season Leaf Rust, Fall Season Septoria Leaf Blotch	Common Root Rot (<i>Cochliobolus</i> spp.), Fusarium Root Rot, Fusarium Crown Rot, Take-All, Rhizoctonia Root Rot

¹DIFENOCONAZOLE 3L ST provides control of fall season powdery mildew, fall season leaf rust, and fall season Septoria leaf blotch in winter wheat for the first 6 weeks after planting. For full season control of these foliar diseases, use Tilt® fungicide according to label instructions.

²DIFENOCONAZOLE 3L ST controls both seed-borne and soil-borne common bunt.

³General seed rots controlled include those caused by saprophytic organisms such as *Penicillium* spp. and *Aspergillus* spp.

SPRING WHEAT AND TRITICALE

Rate/100 lbs. seed	Diseases Controlled	Early Season Control
0.5 fl. oz.	Common Bunt ¹ Seed-borne Septoria Loose Smut	Common Root Rot (<i>Cochliobolus</i> spp.) Rhizoctonia Root Rot

8/9

	Flag Smut General Seed Rots ² Fusarium Seed Scab	
1 fl. oz.	Common Bunt ¹ Loose Smut Flag Smut Seed-borne Septoria General Seed Rots ² Fusarium Seed Scab	Common Root Rot (<i>Cochliobolus</i> spp.) Fusarium Root Rot Fusarium Crown Rot Take-All Rhizoctonia Root Rot

¹ DIFENOCONAZOLE 3L ST controls both seed-borne and soil-borne common bunt.

² General seed rots controlled include those caused by saprophytic organisms such as *Penicillium* spp. and *Aspergillus* spp.

OATS

Rate/100 lbs. seed	Diseases Controlled
0.5 fl. oz.-1 fl. oz.	General Seed Rots ¹ Seedling Blight caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> Seedling Root Rot caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> <i>Pythium</i> and <i>Fusarium</i> Damping-Off Covered Smut Loose Smut Also suppression of Common Root Rot (<i>Cochliobolus</i> spp.)

¹ General seed rots controlled include those caused by saprophytic organisms such as *Penicillium* and *Aspergillus*.

RYE

Rate/100 lbs. seed	Diseases Controlled	Diseases Suppressed ²
0.5 fl. oz.-1 fl. oz.	General Seed Rots ¹ Seedling Blight caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> Seedling Root Rot caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> Seed-borne <i>Septoria</i> <i>Pythium</i> and <i>Fusarium</i> Damping-Off Common Bunt Dwarf Bunt	Common Root Rot (<i>Cochliobolus</i> spp.) <i>Fusarium</i> Crown and Foot Rot Take-All

¹ General seed rots controlled include those caused by saprophytic organisms such as *Penicillium* and *Aspergillus*.

² Suppression means consistent control at a level which is not optimal but is still of commercial benefit.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ALBAUGH CROP PROTECTION, Inc. or Seller. To the extent consistent with

applicable law. Buyer and User agree to hold ALBAUGH and Seller harmless for any claims relating to such factors.

ALBAUGH warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or ALBAUGH, and, (2) Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALBAUGH MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent consistent with applicable law, in no event shall ALBAUGH be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ALBAUGH AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ALBAUGH OR SELLER, THE REPLACEMENT OF THE PRODUCT.

ALBAUGH and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of ALBAUGH.