



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

January 28, 2016

Morris Gaskins
Albaugh LLC
P.O. Box 2127
Valdosta, GA 31604

Subject: Label Amendment – Addition of ornamental me-too use
Product Name: AZOXY 11% + TEB 26% FL
EPA Registration Number: 42750-292
Application Date: 07/10/2015
Decision Number: 507147

Dear Mr. Gaskins:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, you may contact Maryam K. Muhammad at 703-347-0301 or via email at Muhammad.maryam@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to be 'HJ' with a large loop at the bottom.

Hope Johnson, Product Manager 21
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

Jan 28, 2016

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 42750-292

AZOXY 11% + TEB 26% FL

Broad-spectrum fungicide for control of specified turf diseases on golf courses, and specified diseases on ornamentals in field, nursery and container ornamentals and commercial and residential landscapes

ACTIVE INGREDIENTS: % BY WT

Azoxystrobin: methyl (E)-2-[[6-(2-cyanophenoxy)-4-pyrimidinyl]oxy]alpha-methoxymethylene) benzeneacetate	11.9%
Tebuconazole: (+)-alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1 H-1,2,4-triazole-1-ethanol	26.9%
OTHER INGREDIENTS:	61.2%
TOTAL	100.00%

A suspension concentrate fungicide containing 2.34 lb. Tebuconazole and 1.04 lb. Azoxystrobin per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements and directions for use inside booklet. Reformulation is prohibited, See individual container labels for repackaging limitations.

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told by the poison control center or doctor. • Do not give anything to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOTLINE NUMBER - FOR MEDICAL OR TRANSPORT EMERGENCIES INVOLVING THIS PRODUCT, CALL CHEMTREC TOLL FREE AT 1-800-424-9300.	

EPA Reg. No. 42750-292

EPA Est. No. 42750-MO-1

NET CONTENTS: _____ Gal.

MANUFACTURED BY:
Albaugh, LLC
Ankeny, IA 50021

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION. HARMFUL IF SWALLOWED. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemically resistant to this product are listed below.

Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants
2. Chemical-resistant gloves made of any waterproof material (polyvinyl chloride, nitrile rubber or butyl rubber)
3. Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separate from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing;
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Azoxystrobin and tebuconazole are toxic to freshwater and estuarine/marine fish; and azoxystrobin is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

GROUND WATER ADVISORY

Azoxystrobin, can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

Surface Water Advisory:

This product may contaminate water through drift or spray in wind. This product has a high potential for runoff into surface waters for several days to weeks after application. Poorly draining soils, wet soils with readily visible slopes toward adjacent surface waters or soil with shallow water tables are more prone to runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which

this product is applied and surface water features such as ponds, streams and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours. Notify state and/or Federal authorities immediately if you observe any adverse environmental effects due to use of this product.

Notify state and/or Federal authorities and ALBAUGH immediately if you observe any adverse environmental effects due to use of this product.

PHYSICAL HAZARDS

Do not mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. This label must be in the possession of the user at the time of pesticide application.

FAILURE TO FOLLOW THE USE DIRECTIONS AND RESTRICTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Chemigation - Do not apply this product through any irrigation/chemigation system.

PRODUCT INFORMATION

AZOXY 11% + TEB 26% FL is a combination of two broad spectrum, preventative fungicides with systemic properties for the control of many important turfgrass diseases on golf course turf only. AZOXY 11% + TEB 26% FL may be applied as a foliar spray in alternating spray programs or in tank mixes with other registered, turf and ornamental protection products. All applications must be made according to the use directions that follow. This product is not for residential use.

PRODUCT TANK MIX USE PRECAUTIONS

AZOXY 11% + TEB 26% FL has demonstrated some phytotoxic effects when mixed with products that are formulated as EC's. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone have also contributed to phytotoxicity.

INTEGRATED PEST (DISEASE) MANAGEMENT

AZOXY 11% + TEB 26% FL should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. The USE DIRECTIONS section in this label identifies specific IPM recommendations. Consult your local turf authority for additional IPM strategies established for your area. AZOXY 11% + TEB 26% FL may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

AZOXY 11% + TEB 26% FL is a mixture of a Group 3 (tebuconazole) and a Group 11 (azoxystrobin) fungicide. AZOXY 11% + TEB 26% FL has two modes of action: (1) DMI (Demethylation Inhibitor of sterol biosynthesis) [Group 3], and (2) inhibitor of the Qo (quinone outside) site within the electron transport system (Qol) as well as disrupting membrane synthesis by blocking demethylation [Group 11]. Fungal pathogens can develop resistance to products with the same mode of action when, used repeatedly. AZOXY 11% + TEB 26% FL is not cross-resistant with other classes of fungicides which have different modes of action. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for turf in your area. Consult your local or State turf authorities for resistance management strategies that are complementary to those in this label.

Resistance management strategies may include rotating and/or tank-mixing with products having different modes of action or limiting the total number of applications per season. Avoid alternation with other Qols, such as pyraclostrobin and trifloxystrobin or other DMI's such as triadimefon, fenarimol, and myclobutanil. AZOXY 11% + TEB 26% FL should not be alternated or tank mixed with fungicides to which resistance has already developed. Continual use of AZOXY 11% + TEB 26% FL may allow less sensitive strains of pathogens to increase in the population and reduce the efficacy of AZOXY 11% + TEB 26% FL. ALBAUGH encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

Use of a multiple-site contact fungicide such as Daconil as an alternation or tank-mix partner can help alleviate selection pressure on pathogens and delay the onset of resistance.

Since pathogens differ in their potential to develop resistance to fungicides use the USE DIRECTIONS section in this label provides resistance management strategies specific for each disease.

SPRAYING/MIXING

Do not apply AZOXY 11% + TEB 26% FL to turf or ornamentals by air.

AZOXY 11% + TEB 26% FL may be applied with all types of spray equipment commonly used for making ground applications. Do not apply AZOXY 11% + TEB 26% FL through any type of ultra low volume (ULV) spray system (less than 3 gals/A). The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, highly susceptible varieties, or when disease conducive environmental conditions exist

Apply AZOXY 11% + TEB 26% FL in sufficient water volume for adequate coverage.

SPRAY DRIFT MANAGEMENT

ATTENTION

Bermudagrass can be sensitive to AZOXY 11% + TEB 26% FL.

AZOXY 11% + TEB 26% FL is extremely phytotoxic to certain apple varieties. AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

RESTRICTIONS

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

- Do not apply by ground within 100 feet of aquatic areas listed above
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.
- Apply only during alternate years in fields adjacent to aquatic areas listed above.
- DO NOT spray AZOXY 11% + TEB 26% FL where spray drift may reach apple trees,
- Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat.
- DO NOT spray when conditions favor drift beyond area intended for application.
- DO NOT use spray equipment which has been previously used to apply AZOXY 11% + TEB 26% FL to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.
- Do not apply AZOXY 11% + TEB 26% FL to turf or ornamentals by air.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR

APPLICATION INSTRUCTIONS

Apply AZOXY 11% + TEB 26% FL at rates and timings as described in this label.

MIXING INSTRUCTIONS

To prepare spray solution, partially fill the spray tank with clean water and begin agitation. Add the specified amount of AZOXY 11% + TEB 26% FL to the tank, allowing time for good dispersion, then add an adjuvant, if recommended. If tank mixes are required, product should be added to the spray tank in the following order; AZOXY 11% + TEB 26% FL, wettable granular or dry flowable formulations, wettable powders and flowable (aqueous suspensions) products. Finish filling the tank to the desired volume to obtain the proper spray concentration. Maintain agitation throughout the spraying operation.

Do not allow spray mixture to stand overnight or for prolonged periods. Make up only the amount of spray required for immediate use. Sprayers should be thoroughly cleaned immediately after application.

Do not use silicone-based products with AZOXY 11% + TEB 26% FL due to possible phytotoxicity.

If spray-tank mixture is unsprayed for more than 18 hours (overnight), re-suspend product with agitation for 20 minutes.

AZOXY 11% + TEB 26% FL is compatible with many commonly used fungicides, liquid fertilizers, herbicides, insecticides and biological control products, if tank mixes are desired, observe all directions, precautions* and limitations on labeling of all products used.

Consult compatibility charts or your local or State turf authority for compatibility information. Do not combine AZOXY 11% + TEB 26% FL in the spray tank with pesticides, surfactants, or fertilizers, unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective, and non-injurious under your conditions of use. If physical compatibility is unknown, the following procedure should be followed; Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least twenty (20) minutes, if the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

TURF USE DIRECTIONS

AZOXY 11% + TEB 26% FL is for control of certain pathogens causing foliar, stem, and root diseases, including leaf and stem blights, leaf spots, patch diseases, mildews, anthracnose, fairy rings, molds, and rusts of turfgrass in golf courses. This product is not for residential use. Do not apply this product by aerial application.

Integrated Pest (Disease) Management (IPM):

Sound turf management resulting in healthy, vigorous turf is the foundation of a good IPM program. Cultural practices such as proper choice of turf variety, nutrient management, proper cutting height, thatch management, and proper watering, drainage, and moisture stress management should be integrated with the use of fungicides to increase turf vigor and reduce the susceptibility to disease, immunoassay detection kits and extension service diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

Application Directions:

AZOXY 11% + TEB 26% FL should be applied prior to disease development. Mix AZOXY 11% + TEB 26% FL with the required amount of water and apply as a dilute spray application in 2-4 gals of water per 1000 sq. ft. (87-174 gals./A).

Repeat applications at specified intervals for as long as required, not to exceed the maximum specified use rate per season. For spot treatments, use 0.75 fl. oz. AZOXY 11% + TEB 26% FL per 1 to 2 gals. of water.

For use with soil injection applications:

AZOXY 11% + TEB 26% FL may be applied through a liquid fungicide injector for the control of ectotrophic root diseases such as summer patch and take-ail patch. Use AZOXY 11% + TEB 26% FL only in liquid injection equipment specifically designated for pesticide use.

For use with soil injection applications:

Apply AZOXY 11% + TEB 26% FL at 0.75 - 1.7 fl. oz. per 1000 sq. ft. Spray carrier volume should fall within 30-150 gals, of water per 1000 sq. ft. Injection hole spacing of 1 inch by 1 inch is recommended for optimum control. Injection depth should be no greater than 2 inches. One inch depth is recommended for optimum results. Application should follow disease control strategies used for normal broadcast spray programs.

For use in the establishment of turfgrass from seed or in overseeding of dormant turfgrass:

AZOXY 11% + TEB 26% FL may be used for control of certain turfgrass diseases associated with turfgrass establishment from seed. AZOXY 11% + TEB 26% FL may also be used during overseeding of dormant turfgrass.

AZOXY 11% + TEB 26% FL may be safely applied before or after seeding or at seedling germination and emergence to ryegrass, bentgrass, bluegrass, and fescue turfgrass types. Optimum application timing is during seeding. See Application Directions section.

Rate Ranges:

Use the shorter specified application interval and/or use the higher specified rate when prolonged favorable disease conditions exist.

TURF USE RESTRICTIONS

- For use on golf course turf only
- Not for residential use
- Not for use on turf being grown for sale or commercial use as sod.
- Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
- Do not graze or feed clippings from treated turf areas to animals.
- Do not apply AZOXY 11% + TEB 26% FL to turf by air. Applications must be made by ground only.
- Do not exceed 1.5 fl. oz./1,000 sq. ft. every 30 days on any variety of bermudagrass.
- In FL, do not apply AZOXY 11% + TEB 26% FL to bermudagrass golf course greens when temperatures exceed 90°F.
- Do not exceed 9.0 fl. oz. of AZOXY 11% + TEB 26% FL per 1,000 sq. ft. per year
- Do not apply more than 6 applications per year.
- In the state of New York, do not exceed 2.7 fl. oz. of AZOXY 11% + TEB 26% FL per 1,000 sq. ft. (2.2 lb a.i./Acre) per year

Target Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Application Interval (days)	Remarks*
Anthracnose (<i>Colletotrichum cereale</i>) (formerly known as <i>C. graminicola</i>)	0.75 - 1.5	14 - 28	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development. The maximum application interval for greens is 14 days.
Bentgrass Dead Spot (<i>Ophiosphaerella Agrostis</i>)	0.75 - 1.5	14	Begin applications prior to disease development.
Brown Patch (<i>Rhizoctonia solani</i>)	0.5 - 0.75	14	For higher cut cool season turfgrass, St, Augustine grass and centipedegrass (above 0.375 inch in height). Apply when conditions are favorable for disease development
Brown Patch (<i>Rhizoctonia solani</i>) and other Rhizoctonia (<i>R. spp.</i>) diseases such as Large Patch, Zoysia Patch, Leaf & Sheath Spot (<i>R. zea</i>)	0.75 - 1.5	14 - 28	For Brown Patch, apply when conditions are favorable for disease development. For Large Patch of all warm season turfgrasses, make 1 or 2 applications in the fall prior to infection or when conditions are favorable for infection. For Zoysia Patch, make 1 or 2 applications approximately one month prior to zoysiagrass dormancy. Reapply 14 to 28 days later. For Leaf & Sheath Spot (<i>R. zea</i>), apply when conditions are favorable for infection such as sequential or periods of temperatures at or above 90 degrees Fahrenheit. Curative control may necessitate several applications. Target spray applications at crown of turfgrass.
Cool Weather Brown Patch, Yellow Patch (<i>Rhizoctonia cerealis</i>)	1.5	28	Make 1 or 2 applications in fall or when conditions are favorable for disease development.
Dichondra Rust	1.5	14 - 21	Apply when conditions are favorable for disease

Target Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Application Interval (days)	Remarks*
<i>(Puccinia dichondrae)</i>			development.
Dollar Spot <i>(Sclerotinia homoeocarpa)</i>	0.75 - 1.5	7 - 14	Apply preventively on higher cut cool season turfgrass (fairways, lawns, etc.) mowed at or above 0.375 inch in height during periods of low to moderate disease pressure.
Dollar Spot <i>(Sclerotinia homoeocarpa)</i>	0.75 - 1.5	14 - 28	Apply preventively when conditions are favorable for disease development. For turf with active dollar spot, use the higher rate and combine with Daconil.
Fairy Ring <i>(Lycoperdon spp., Agrocybe pediades, and Bovista plumbea)</i>	0.75 - 1.5	14 - 28	Apply as soon as possible after fairy ring symptoms develop. Apply in 2-4 gals, of water per 1000 sq. ft. Add the recommended rate of a wetting agent to the final spray. Severely damaged or thin turf may require reseeding. Fairy ring symptoms may take 2 to 3 weeks to disappear following application. If area is hydrophobic, use wetting agents and irrigate prior to application(s) of AZOXY11% + TEB 26% FL. Reapplication after 28 days may be required in some cases.
Gray Leaf Spot <i>(Pyricularia grisea)</i>	0.75 - 1.5	14 - 28	Use AZOXY11% + TEB 26% FL in a preventive disease control program. Begin applications before disease is present and alternate with other fungicide chemistries that control gray leaf spot.
Gray Snow Mold, Typhula Blight <i>(Typhula incarnata)</i>	1.5	10 - 28	Make two applications of 1.5 fl. oz. spaced 10-28 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide, such as Daconil and Medallion, may enhance control under severe disease pressure.
Leaf Rust, Stem Rust, Stripe Rust <i>(Puccinia spp.)</i>	0.75 - 1.5	14 - 28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Leaf Spot <i>(Bipolaris sorokiniana)</i>	0.75 - 1.5	14 - 21	Apply when conditions are favorable for disease development.
Melting Out <i>Drechslera poae)</i>	0.75 - 1.5	14 - 21	Apply when conditions are favorable for disease development.
Microdochium Patch (formerly known as Fusarium Patch) <i>(Microdochium nivale)</i>	0.75 - 1.5	14 - 28	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Necrotic Ring Spot <i>(Ophiosphaerella korrae)</i>	1.5	14 - 28	Apply when conditions are favorable for disease development.
Pink Patch <i>(Limonomyses roseipellis}</i>	0.75 - 1.5	14 - 28	Apply when conditions are favorable for disease development.
Pink Snow Mold <i>(Microdochium nivale)</i>	1.5	10 - 28	Make two applications of 1.5 fl. oz. spaced 10-28 days apart in- late fall just before snow cover. Tank mixing with another snow mold fungicide, such as Daconil .and Medallion, may enhance control under severe disease pressure.
Powdery Mildew <i>(Erysiphe graminis)</i>	0.75 - 1.5	14 - 28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Pythium Blight, Pythium Root Rot <i>(Pythium aphanidermatum,</i>	1.5	10 - 14	Use preventatively. Begin applications when c conditions are favorable for disease infection prior to disease symptom development.

Target Diseases	Use Rate (fl. oz. product per 1000 sq. ft.)	Application Interval (days)	Remarks*
<i>Pythium</i> spp.)			During periods of prolonged favorable conditions, treat on the 10 day application interval. For use on newly seeded well as established turf.
Red Thread (<i>Laetisaria fuciformis</i>)	0.75 - 1.5	14 - 28	Apply when conditions are favorable for disease development.
Southern Blight (<i>Scierotium rotsii</i>)	0.75 - 1.5	14 - 28	Apply when conditions are favorable for disease development.
Spring Dead Spot (<i>Ophiosphaerella korrae</i>) or (<i>Ophiosphaerella narmari</i>) or (<i>Ophiosphaerella herpotricha</i>)	1.5	14 - 28	Apply 1 or 2 applications approximately one month prior to bermudagrass dormancy. 1/4" to 1/2" of irrigation directly after application is recommended. Reapply 14 to 28 days later.
Summer Patch (<i>Magnaporthe poae</i>)	0.75 - 1.5	14 - 28	Apply when conditions are favorable for disease development. Initiate applications when soil temperatures reach 65°F at a 2 inch soil depth.
Take-all Patch (<i>Gaeumannomyces graminis</i> var. <i>venae</i>)	1.5	28	Begin applications when conditions are favorable for disease infection prior to disease symptom development. Make two applications, 28 days apart in the spring and two applications 28 days apart in the fall.

*Do not apply more than two sequential applications of AZOXY 11% + TEB 26% FL for control of Gray Leaf Spot and *Pythium* spp. For all other diseases when Gray Leaf Spot and *Pythium* spp. are not present, do not apply more than three sequential applications of AZOXY 11% + TEB 26% FL.

AZOXY 11% + TEB 26% FL Rate Conversion Chart for Turf

Fl. Oz. Product* Per 1000 Sq. Ft.	Pints Product Per Acre	Gal. Product Per Acre
0.75	2.0	0.26
1.0	2.7	0.34
1.5	4.0	0.51

*One fluid ounce of AZOXY 11% + TEB 26% FL contains 0.2925 ounces (weight) of tebuconazole and 0.1300 ounces (weight) azoxystrobin.

Amount of AZOXY11% + TEB 26% FL for Turf Applications per 100 Gallons

Spray volume: 2 gals./1000 sq. ft.

AZOXY11% + TEB 26% FL Use Rate (fl. oz./1000 sq. ft.)	Pints of product/100 gals.
0.75	2.3
1.0	3.1
1.5	4.7

Spray volume: 3 gals./1000 sq. ft.

AZOXY11% + TEB 26% FL Use Rate (fl. oz./1000 sq. ft.)	Pints of product/100 qals.
0.75	1.5
1.0	2.1
1.5	3.0

Spray volume: 4 gals./1000 sq. ft.

AZOXY11% + TEB 26% FL Use Rate (fl. oz./1000 sq. ft.)	Pints of product/100 gals.
0.75	1.15
1.0	1.55
1.5	2.25

USE DIRECTIONS FOR DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL and RESIDENTIAL LANDSCAPES

ORNAMENTAL USE RESTRICTIONS

- For use on ornamental plants only, not for woodlands or forest management
- Intended for use only by professional applicators
- Do not apply more than 15.0 fl oz per acre in a single application.
- Do not make more than 4 applications per year at highest rate
- Do not apply more than 0.47 gallons (60.0 fl oz) of AZOXY 11% + TEB 26% FL (equal to 1.09 lbs of tebuconazole) per acre per year
- Do not apply to bearing fruit trees or vegetables

INTEGRATED PEST (DISEASE) MANAGEMENT: AZOXY11% + TEB 26% FL should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance, optimum plant populations, proper fertilization, winter and/or spring pruning, plant residue management and proper timing and placement of irrigation. Immunoassay detection kits and diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

RESISTANCE MANAGEMENT: Some ornamental disease pathogens are known to have developed resistance to fungicides used repeatedly for their control. AZOXY11% + TEB 26% FL should be applied in an alternation or tank mix program with other registered fungicides that have a different mode of action and to which pathogen resistance has not developed. Do not make more than three (3) sequential applications of AZOXY11% + TEB 26% FL before alternating with a fungicide of a different mode of action. A sound resistance management program would include blocks of three AZOXY11% + TEB 26% FL applications separated by blocks of two alternate fungicide applications. Do not alternate AZOXY11% + TEB 26% FL with other strobilurin fungicides.

Use only surfactants approved for ornamental plants in combination with AZOXY11% + TEB 26% FL. Do not use silicone based products with AZOXY11% + TEB 26% FL due to possible phytotoxicity. Always test tankmixes on a small group of representative plants prior to broadscale use.

In addition, do not tankmix AZOXY11% + TEB 26% FL with other fungicides, insecticides, herbicides, fertilizers, adjuvants, etc, unless local experience indicates that the tankmix is safe to ornamental plants.

ORNAMENTAL USE PRECAUTIONS

Do not apply AZOXY11% + TEB 26% FL to apple or cherry trees (Flowering, Yoshina variety) due to

possible phytotoxicity. Further, do not use spray equipment that has applied AZOXY11% + TEB 26% FL for use in these sensitive crops due to possible phytotoxicity from residue remaining in the sprayer.

AZOXY11% + TEB 26% FL may be applied to certain varieties of crabapple for control of apple scab. Due to the large number of genera, species, and varieties of crabapple, it is impossible to test every one for tolerance to AZOXY11% + TEB 26% FL. The professional user should conduct small scale testing to insure plant safety prior to broadscale commercial use on plant genera and species not listed on this label.

PLANT SAFETY: Due to the large number of genera, species and varieties of ornamental and nursery plants, it is impossible to test every one for tolerance to AZOXY11% + TEB 26% FL. Neither the manufacturer nor the seller has determined whether or not AZOXY11% + TEB 26% FL can be used safely on genera, species, or varieties of ornamental and nursery plants not specified on this label. The professional user should conduct small scale testing to insure plant safety prior to broadscale commercial use on plant genera and species not listed in this label.

NOTE: The "Directions for Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results.

This product is not recommended for use on African Violets, Begonias, Boston Fern, and Geraniums.

Do not use spray equipment that has applied AZOXY11% + TEB 26% FL for use in these sensitive crops due to possible phytotoxicity from residue remaining in the sprayer.

AZOXY 11% + TEB 26% FL can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when AZOXY 11% + TEB 26% FL is used in conjunction with sound disease management practices

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

Apply AZOXY 11% + TEB 26% FL at rates of 6-15 fl oz per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.

ORNAMENTALS DISEASE CONTROL

PLANTS	DISEASE	APPLICATION	
		To Prevent Diseases	To Treat Existing Disease
Roses	Black Spot Powdery Mildew Rust	Apply every 14 - 21 days during the growing season, starting when leaves first appear	Apply every 14 days for a total of 3 applications beginning at the first sign of disease
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year, 14 - 21 days apart, beginning with Spring bud break Rotation or Tank mixing with barrier protectant fungicides is recommended for resistance management	
Crabapples* (Ornamental), Dogwoods and Other Landscape (Ornamental) Trees	Anthracnose Leaf Spot Powdery Mildew Rust Scab		
Azaleas, Camellias, Rhododendrons and Other Landscape (Ornamental) Shrubs	Anthracnose Black Spot Leaf Spot Petal Blight Powdery Mildew Rust	Petal Blight - Apply 2 - 3 times per week into the flowers as the open and develop color.	
Ground Covers and Vines	Southern Blight		
HOW MUCH TO USE FOR SMALL PLANTINGS: ADD 1.5 TEASPOONS TO 2.5 GALLONS OF WATER			

*Do not apply to Flame, Brandywine or Novamac varieties

Pump Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank
2. Close the sprayer, shake well and pressurize
3. Adjust nozzle to a coarse spray pattern and apply
4. Occasionally re pressurize the sprayer if needed to maintain a good spray pattern

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop up and remove to chemical waste storage area until proper disposal can be made if product cannot be reused according to the label.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance. Open dumping is prohibited.

CONTAINER HANDLING: [less than 5 gallons] - Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Container Handling [greater than 5 gallons] - Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY NOTICE:

Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product, if the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully, it is impossible to eliminate all risks inherently associated with the use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or turf conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ALBAUGH, LLC or Seller. To the extent permitted by applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold

ALBAUGH, LLC and Seiler harmless for any claims relating to such factors, ALBAUGH, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or

under conditions not reasonably foreseeable to or beyond the control of Seller or ALBAUGH, LLC, and, (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, ALBAUGH, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABELSTATED ABOVE.

To the extent permitted by applicable law, in no event shall ALBAUGH, LLC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ALBAUGH, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ALBAUGH, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

ALBAUGH, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of ALBAUGH, LLC.