



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

42750-309

Date of Issuance:

8/1/16

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Azoxystrobin 2L H&G

Name and Address of Registrant (include ZIP Code):

Morris Gaskins  
Albaugh, LLC  
P.O. Box 2127  
Valdosta, GA 31604-2127

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Shaja B. Joyner, Product Manager 20  
Fungicide-Herbicide Branch  
Registration Division 7505P

Date:

8/1/16

EPA Form 8570-6

2. You are required to comply with the data requirements described in the DCI identified below:
  - a. Azoxystrobin GDCI-128810-892

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, “EPA Reg. No. 42750-309.”
4. Submit one copy of the final printed label for the record before you release the product for shipment.


Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 3/9/2016

If you have any questions, please contact Nathan Mellor by phone at 703-347-8562, or via email at [mellor.nathan@epa.gov](mailto:mellor.nathan@epa.gov).

Enclosure



**AZOXYSTROBIN 2L H&G**  
Fungicide Concentrate

ACTIVE INGREDIENT:

Azoxystrobin: methyl (E)-2-{2-[6-(2-cyanophenoxy) pyrimidin-4-yloxy]phenyl}-3-methoxycarylate <sup>1*</sup> . . . . .	22.9%
OTHER INGREDIENTS: . . . . .	<u>77.1%</u>
TOTAL: . . . . .	100.0%

<sup>1</sup>CAS # 131860-33-8 \*IUPAC  
This product contains 2.08 lbs. of active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

FIRST AID

IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF IN EYES	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
In case of medical or transport emergency, contact CHEMTREC toll free at 1-800-424-9300	

EPA Reg. No. 42750-XXX

EPA Est. No.

NET CONTENTS:

MANUFACTURED FOR:  
Albaugh, LLC  
1525 NE 36<sup>th</sup> Street  
Ankeny, IA 50021

## Appendix 1 - Consolidated List of Label Claims

- *Broad spectrum and systemic disease control on ornamentals, flowers, home lawns, vegetables, fruits, trees, and shrubs*
- *Broad spectrum and systemic disease control on ornamentals, flowers, trees, and shrubs*
- *Broad spectrum and systemic disease control on home lawns*
- *For use on home lawns*
- *Broad spectrum and systemic disease control on vegetables and fruits*
- *Controls brown patch, powdery mildew, anthracnose, rust, blights. Scab, and more*
- *Prevents and stops major diseases on vegetables, flowers, lawns, trees, and shrubs*
- *Concentrate - Mix with water and spray*
- *For use in pump-up and hose-end sprayers*
- *Four ounces will treat up to 12,000 sq. ft. of lawn*
- *Eight ounces will treat up to 24,000 sq. ft. of lawn*
- *Sixteen ounces will treat up to 48,000 sq. ft. of lawn*
- *Absorbed into the plant- Can't be washed off by rain or sprinklers after spray has dried*
- *Rainfast shortly after spray has dried*
- *Not affected by rain after spray has dried*
- *Systemic action- Can't be washed off by rain after spray has dried*
- *Broad spectrum disease control on home lawns and ornamentals*
- *Prevents and stops: Anthracnose, Blights, Rust, Brown Patch Powdery Mildew, Leaf Spot and more*
- *Works up to 4 weeks for certain diseases*

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking chewing gum or using tobacco.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear long sleeved shirt and long pants, socks and shoes and chemical resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl-rubber. Wear protective eyewear (if appropriate).

#### Restrictions:

Do not use utensils such as measuring spoons and cups after use with pesticides.

Do not use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

#### Precaution:

*(Walking dog icon)* People and pets should not be allowed in treated areas until spray has dried.

#### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separate from other laundry.

#### ENVIRONMENTAL HAZARDS

Azoxystrobin is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Azoxystrobin can be persistent for several months or longer. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to

intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

#### GROUND WATER ADVISORY

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

#### SURFACE WATER ADVISORY

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

#### PHYSICAL CHEMICAL HAZARDS

Do not mix or allow coming in contact with oxidizing agent. Hazardous Chemical reaction may occur.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For best results read and follow all label directions.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ALBAUGH, LLC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ALBAUGH, LLC and Seller harmless for any claims relating to such factors.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR INJURY TO DESIRABLE PLANTS

#### GENERAL INFORMATION:

This water-based systemic fungicide effectively prevents and controls major diseases on flowers, fruits, lawns, ornamentals, shrubs, trees, and vegetables.

#### USE TIPS

- This product is highly effective when it is used to prevent diseases or applied at the first sign of disease. Begin applications when conditions are favorable for disease infection. This product will not eliminate disease already present.
- When used as directed, this product is safe to plants and controls the diseases listed. Plants not listed may be injured by this product and diseases not listed may not be controlled.

- Do not use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.
- If you need more information about fruit, vegetable, lawn and ornamentals diseases, contact your local County Extension Service.

To control diseases on plants other than turfgrass, apply product to the plant until it begins to run off of the leaves.

- Allow spray to completely dry before watering.

#### MIXING PROCEDURES

Shake well before using.

Note:        ¼ Fl. Oz. = 1½ teaspoons  
                  ½ Fl. Oz. = 1 Tablespoon = 3 teaspoons  
                  1 Fl. Oz. = 2 Tablespoons = 6 teaspoons

**Restrictions:** Do not use utensils such as measuring spoons and cups after use with pesticides.

Note: Move indoor houseplants outdoors before treating. Allow to dry before handling or moving indoors.

#### **Pump-Up Sprayer (for all uses)**

- Fill with ½ required amount of water.
- Add the amount of concentrate specified in the table that follows.
- Fill with remaining water.
- Shake well and pressurize
- Adjust nozzle for medium to coarse spray and apply.
- Occasionally repressurize sprayer if needed to maintain spray pattern.

#### **Concentrate Plus Water Style Hose-End Sprayer (use for application to Lawns, Flowers, Trees, Ornamentals and Shrubs).**

- From the following table, determine the amount of concentrate required.
- Pour concentrate into sprayer jar to the required "fl. oz." level.
- Then add the required amount of water to the "gals." level.
- Replace sprayer top on jar and shake well.
- Attach sprayer to hose and apply.
- For Lawns, apply to the measured square foot area to be sprayed.
- For Flowers, Trees, Ornamentals and Shrubs, apply to obtain sufficient spray coverage without excessive runoff.

#### **Dial-Style Hose-End Sprayer (use for application to Lawns, Flowers, Trees, Ornamentals and Shrubs).**

- Fill the sprayer jar with only enough product to cover the plants or the area to be treated.
- Set Dial on the "fl. oz./gallon" level specified in the following table.
- Close sprayer, attach to hose and apply.
- For Lawns, apply to the measured square foot area to be sprayed.

- For Flowers, Trees, Ornamentals and Shrubs, apply to obtain sufficient spray coverage without excessive runoff.

*(Optional language for product sold in a HOSE-END SPRAYER package)*

**[HOSE-END SPRAYER: Easy to Use (use for application to Lawns, Flowers, Trees, Ornamentals and Shrubs).]**

- Make sure knob control is in the OFF position. Shake well and attach to hose.
- Turn water on at faucet, aim nozzle toward lawn or desired plant and turn control lever to ON to begin spraying.
- When finished turn control lever to OFF position. Discharge residual water pressure by turning ON and OFF again.
- For Lawns, apply to the [xxx) square foot area to be sprayed. [Square feet value to be inserted per specific package size].
- For Flowers, Trees, Ornamentals and Shrubs, apply to obtain sufficient spray coverage without excessive runoff.

#### APPLICATION INSTRUCTIONS

##### Precaution:

*(Walking dog icon)* People and pets should not be allowed in treated areas until spray has dried.

##### Restrictions:

1. Do not apply this product to apples, crabapples, or flowering (ornamental) cherry trees due to possible phytotoxicity.
2. DO NOT use sprayers that have been used to apply this product to spray apple trees, crabapples and flowering (ornamental) cherries. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

Note: Apply sufficient water to obtain coverage without excessive runoff.

FLOWERS (annuals and perennials)

##### General Directions

Do not apply more than two applications before alternating with a fungicide that has a different mode of action.

Do not make more than 4 applications of this product (.75 lbs. azoxystrobin per acre per year).

Plant	Disease	How Much to Apply (per gal. of water)	Comments
Roses	Downy Mildew Leaf Spot Powdery Mildew Rusts	1/4-1/2 tsp.	Thorough coverage needed for best results. Begin application at bud break. Treat every 7-21 days.
Annual and Perennial Flowers-Bedding Plants	Downy Mildew Leaf Spots Leaf Blights	1/8-1/4 tsp.	Apply every 7-14 days prior to infection. Do not apply the 3 teaspoon rate on less than 14



Plant	Disease	How Much to Apply (per gal. of water)	Comments
	Powdery Mildew Rusts Scab		day spray intervals.
Annual and Perennial Flowers/Mature Plants	Downy Mildew Leaf Spots Leaf Blights Powdery Mildew Rusts Scab	1/4-1/2 tsp.	Apply before disease begins.  Apply every 7-28 days.
Geraniums	Flower Blights	1/8-1/2 tsp.	Apply before disease begins.  Apply every 7-21 days.
Petunias Vinca	Shoot/Stem Diseases	1/8-1/4 tsp.	Apply every 7-28 days.
Chrysanthemums Cyclamen Impatiens Sweet alyssum	Soil-Borne Diseases	1/8-1/4 tsp.	Use as a directed spray.  Apply every 7-21 days.

## LAWNS

### General Directions

1. Apply after mowing or allow sprayed area to completely dry before mowing.
2. Measure the length and width of area to be treated before application. Length x width = square feet.  
For example, an area 25 ft. long and 20 ft. wide is 500 sq. ft.
3. Begin applications when conditions are favorable for disease infection, prior to disease symptoms.
4. Can be safely applied after seeding or at seed germination.
5. Do not use more than 5 lbs. azoxystrobin per acre per year.

Disease	How Much to Apply (tsp. per 250 sq. ft.)	Comments
Brown Patch	½ - 1 tsp.	First application should be in May or June before disease is present.  If needed, repeat treatment every 14-28 days.
Anthracnose Fusarium Patch Pink Patch Summer Patch Gray Leaf Spot Leaf Spot Rhizoctonia Leaf Spot Necrotic Ring Spot Leaf Rust Stem Rust Stripe Rust Melting Out Powdery Mildew Southern Blight Red Thread	½ - 1 tsp.	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.  Apply every 14-28 days

Cool Weather Brown Patch Yellow Patch	½ - 1 tsp.	Make 1 or 2 applications in fall (14-28 days interval) or when conditions are favorable for disease development.
Take-All Patch	½ - 1 tsp.	Apply twice in the Fall (Sept. and Oct.) and twice in the Spring (April and May).  Southern lawns also need to be treated twice in the summer (June-August).
Snowmold (Gray or Pink)	1 ½ tsp.	Make a single application in late fall.
	1 tsp.	or Apply twice in the fall prior to snow cover waiting 10-14 days between each application.
Pythium Blight Pythium Root Rot	½ - 1 tsp.	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.  Apply every 10-14 days.  For use on newly seeded as well as established lawns.
Zoysia Patch	½ - 1 tsp.	Apply 1 or 2 applications (14-28 days interval) approximately one month prior to zoysiagrass dormancy

## TREES, ORNAMENTS AND SHRUBS

### General Directions

Do not apply more than two applications before alternating with a fungicide that has a different mode of action.

Apply sufficient water to obtain coverage without excessive runoff.

Plant	Disease	How Much to Apply (per gal. of water)	Comments
Conifers Cedar Christmas Trees Cypress Fir Juniper Pine Spruce	Conifer Blights	1/8-1/2 tsp.	Apply every 7-28 days.
Abelia Alder Arborvitae Aspen Azalea Bougainvillea Butterfly bush ( <i>Buddleia</i> ) Boxwood Camellia Cedar Clethra Coleus Crapemyrtle Dogwood	Downy Mildew Leaf Spot Scab	1/8-1/2 tsp.	Apply every 7-28 days.

Plant	Disease	How Much to Apply (per gal. of water)	Comments
Dumbcane ( <i>Diffenbachia</i> ) Euonymus Fatsia Fig* Forsythia Grasses Hawthorn Heather Hibiscus Hydrangea Ivy Japanese andromeda Liriodendron Magnolia Maples Nandina Oak Oleander Palms Philodendron Photinia Plum (flowering or purple-leaf) Poinsettia Pothos Red Bud Rhododendron Rose of Sharon Rubber tree Sedum Viburnum Wiegela Zebra plant			* Figs: Upon treating, do not use for food and/or feed purposes.
Azalea Dogwood Rhododendron	Anthraco	1/2-1 tsp.	Apply every 7-28 days.
Ajuga Azalea Barberry Birch Bradford's pear Ceanothus Crapemyrtle Dogwood Gardenia Hawthorn Hibiscus Hydrangea Holly Lilac Laurel Mock-orange Oak Pampas & Ribbon grasses Photinia Rhododendron Sage Spirea	Powdery Mildew	1/8-1/2 tsp.	Apply every 7-28 days

Plant	Disease	How Much to Apply (per gal. of water)	Comments
Viburnum Willow			
Aucuba Azalea Boxwood Holiday cactus Caladium Cotoneaster Japanese andromeda Laurel Palm Rhododendron Rubber plant ( <i>Peperomia</i> ) Yew Yucca	Soil-Borne Diseases	1/8-1/2 tsp.	Use as a directed  Apply every 7-28 days
Azalea Rhododendron	Shoot/Stem Diseases	1/8-1/4 tsp.	Apply every 7-28 days

FRUIT AND NUT TREES, BERRIES AND VINES:

NOT FOR USE ON APPLES, CRAB-APPLES, OR FLOWERING (ORNAMENTAL) CHERRIES

General Directions

1. Do not apply more than two applications before alternating with a fungicide that has a different mode of action.
2. Do not make more than 4 applications of this product (.75 lbs. azoxystrobin per acre per year) per crop (unless specified in use directions).
3. Apply sufficient water to obtain coverage without excessive runoff.

Plant	Disease	How Much to apply (per gallon of water)	Days to Wait to Harvest	Comments
Almonds	Alternaria Leaf and Fruit Spot Anthracnose Brown Rot Blossom Blight Leaf Blight Leaf Rust Scab Shothole	1½ - 2 tsp.	28	Start applications during early bloom.  Apply every 7-14 days.  Do not make more than 4 applications of this product per crop.
Berries, Bushberry subgroup: Blueberry Currant Elderberry Gooseberry Huckleberry  Including all cultivars and/or hybrids of these Lingonberry Juneberry	Botryosphaeria Canker Powdery Mildew Septaria Blight Mummyberry Alternaria Fruit Rot Phomopsis Stem Canker Anthracnose Fruit Rot	1 – 2 tsp.	0	Apply every 7-14 days.  Do not make more than 3 applications of this product per crop.

Plant	Disease	How Much to apply (per gallon of water)	Days to Wait to Harvest	Comments
Salal				
Berries, Caneberry subgroup Blackberry Bingleberry Boysenberry Dewberry Lowberry Marionberry Olafieberry Youngberry Loganberry Red and black raspberry  Including all cultivars and/or hybrids of these	Botryosphaeria Canker Anthracnose Powdery Mildew Leaf Spot Colletotrichum Rot Spur Blight Rosette or Double Blossom of Blackberries	1 - 2 tsp.	0	Apply every 7-14 days.  Do not make more than 6 applications of this product per crop.
Citrus Fruit Calamondin Citron Citrus hybrids Grapefruit Kumquat Lemon Lime Mandarin Orange (sour and sweet) Pummelo Satsuma mandarin Tangerine  Including all cultivars and/or hybrids of these	Greasy Spot Melanose Scab Albinism Post Bloom Fruit Drop (PFD) Alternaria Leaf and Fruit Spot	1½ - 2 tsp	0	Apply every 7-21 days.  Do not make more than 4 applications of this product per crop.
Grapes Including Mascadines	Downy Mildew Phomopsis Cane and Leaf Spot Powdery Mildew Black Rot  Suppression Botrytis Bunch Rot	1½ - 2 tsp	14	Apply every 10-14 days.  Do not make more than 4 applications of this product per crop.
Pistachios	Alternaria Late Blight Botryosphaeria panicle and Shoot Blight Septoria Leaf Spot	1 ½ - 2 tsp	7	Apply every 7-21 days.  Do not make more than 3 applications of this product per crop.
Stone Fruits Apricot Cherry, Sweet Cherry, Tart Nectarine Peach Plum	Alternaria Spot and Fruit Rot Anthracnose Leaf Rust Powdery Mildew Scab Shot Hole	1½ - 2 tsp.	0	Apply every 7-14 days.  For Brown rot blossom blight begin spray at early bloom and continue through petal fall.  Do not make more than 4

Plant	Disease	How Much to apply (per gallon of water)	Days to Wait to Harvest	Comments
Plumcot Prune	Brown Rot Blossom Blight and Fruit Rot			applications of this product per crop.
Strawberry	Anthrachnose Powdery Mildew  Suppression of Botrytis on the Foliage	2 tsp	0	Apply every 7-10 days.  Do not make more than 4 applications of this product per crop.
Tropical Fruit Pawpaw Persimmon	Anthrachnose Cercospora Leaf Spot Powdery Mildew Rust	1- 2 tsp	0	Apply every 10-14 days.  Do not make more than 4 applications of this product per crop.

## VEGETABLES -

### General Directions

1. Do not apply more than two applications before alternating with a fungicide that has a different mode of action. This product is a group 11 fungicide. Do not alternate with another group 11 fungicide.
2. Do not make more than 6 applications of this product per crop (.75 lbs. of azoxystrobin per acre per year) unless specified in use directions.
3. Apply sufficient water to obtain coverage without excessive runoff.

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Asparagus	Stemphyllium purple spot	1½ - 2 tsp.	100	Apply on a 7-14 day schedule to asparagus foliage after harvest of spears is complete.
Brassica Head and Stem subgroup: Broccoli Chinese broccoli Brussel sprouts Cabbage Chinese cabbage Chinese mustard Cabbage Cauliflower Cavalo broccolo Kohlrabi  Including all cultivars and/or hybrids of these	Alternaria Leaf Spot Downy Mildew	1 – 1¾ tsp	0	Apply on a 7-14 day schedule.
Brassica Leafy Greens subgroup: Broccoli raab Cabbage Chinese Collards Kale Mizuna	White Rust Black Spot Cercospora Leaf Spot	1 – 2 tsp.	0	Apply on a 7-14 day schedule.
	Soil-Borne Diseases Seedling Root Rot, Basal Stem Rot	½ - 7/8 tsp. For 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Mustard greens Mustard spinach Rape greens  Including all cultivars and/or hybrids of these				emerged plants
Bulb Vegetables Garlic Leek Onion. Bulb Onion, green Welsh onion Shallot	Foliar Diseases Cladosporium Leaf Blotch Purple Blotch Rust White Rot	1 - 1½ tsp.	0	Apply on a 7-14 day schedule. Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.
	Botrytis Leaf Blight Downy Mildew	1¼ - 2 tsp.		
	Soil-Borne Diseases Rhizoctonia Damping Off	½ - 7/8 tsp. for 400 feet of row		
Carrots	Alternaria Leaf Spot Ascochyta Leaf Spot Cercospora Leaf Spot Early Blight Late Blight Powdery Mildew Rust White Mold White Rust	1¼ - 2 tsp.	0	Apply on a 7-14 day schedule.
	Soil-Borne Diseases Rhizoctonia Root	½ - 7/8 tsp. for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.
Celery	Alternaria Leaf Spot Anthracnose Cercospora Leaf Spot Downy Mildew Early Blight Late Blight Powdery Mildew Septoria Leaf Spot White Rust	1¼ - 2 tsp.	0	Apply on a 7-14 day schedule
	Soil-Borne Diseases Rhizoctonia Root Rot	½ - 7/8 tsp. for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Corn Pop Sweet	Rust	1 - 1¼ tsp.	7	Apply on a 7-14 day schedule.
	Anthracnose Leaf Blight Eye Spot Gray Leaf Spot Northern Corn Leaf Blight Northern Corn Leaf Spot Southern Corn Leaf Blight	1¼ -2 tsp.		Apply on a 7-14 day schedule.  For Gray leaf spot, the first application must be made at the onset of disease.
Cucurbits Cantaloupe Chayote Chinese-waxgourd Cucumber Gourds Honeydew Melons (bitter melon, balsam apple) Muskmelon Watermelon Pumpkin Squash Zucchini  Including cultivars and/or hybrids of these	Anthracnose Belly Rot Downy Mildew Gummy Stem Blight Leaf Spots Myrothecium Canker Powdery Mildew	1½ - 2 tsp.	1	Make a preventative application on a 5-7 day schedule.  For belly rot apply at the 1-3 leaf crop stage with second application made just prior to vine tip over or 10-14 days later, whichever occurs first.
	Soil-Borne Diseases Rhizoctonia root rot	½ - 7/8 tsp. for 200 feet of row		Apply after seeds planting by spraying over the row or directed to the base of transplants or emerged plants





Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
rice bean, southern pea, urd bean, yardlong bean),  Broad bean (Fava bean) Chickpea (garbanzo bean) Guar Jackbean Lablab bean (hyacinth bean).  Pea ( <i>Pisum</i> spp.) (includes dwarf pea, edible-pod pea, English pea, garden pea, green pea, field pea, snow pea, sugar snap pea). Pigeon pea Sword bean				
Mint (fresh)	Rust Powdery Mildew	1-2tsp.	0	Apply on a 7-10 day schedule.  Do not make more than 3 sprays/crop.
Pepper Bell Pepper Non-Bell Pepper Sweet Non-Bell Pepper	Powdery Mildew Anthracnose	1-2 tsp.	0	Apply on a 7-14 day schedule.
Eggplant Okra	Soil-Borne Diseases Rhizoctonia Seedling Rot	½ - 7/8 tsp. for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.
Potatoes	Early Blight Late Blight Black Dot Powdery mildew Late Blight	1-2 tsp.    2 tsp.	14	Begin application when plants are 6 inches high or at prior to first sign of disease.  Do not make more than 2 sprays before alternating to a different product with alternate chemistry.  Do not make more than 6 applications (1-2 tsp each application).
	Soil-Borne Diseases Black Scurf Silver Scurf	½ - 7/8 tsp. for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.
Tomatoes	Anthracnose Black Mold Buckeye Rot Early Blight Powdery Mildew Septaria Leaf Spot Target Spot  Late Blight	¾ tsp.	0	Do not apply until 21 days after transplanting or 35 days after seeding. Apply on a 7-21 day schedule.  For late blight apply on a 5-7 day schedule. Make only one

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
		1 tsp.		application before alternating to another product with a different chemistry.
Vegetables, leaves of root and tuber, group Beet, garden and sugar Burdock Carrot Cassava, bitter and sweet Celeriac (celery root) Chervil, turnip- rooted Chicory Dasheen (taro) Parsnip Radish Radish, oriental (daikon) Rutabaga Salsify, black Sweet potato Tanier Turnip Yam, true	Foliar Diseases Alternaria Leaf Spot Ascochyta Leaf Spot Cercospora Leaf Spot Powdery Mildew Rust White Rust	1-2 tsp.	0	Apply every 7-14 days. Use 1 ¼ - 2 tsp. rate for Cercospora leaf spot and Powdery Mildew. For Powdery Mildew apply on 5-7 day schedule.
	Soil-Borne Diseases Circular Spot, Southern Blight, Rhizoctonia Stem Canker, Crown Rot, Pythium Root Rot	½ - 7/8 for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.
Vegetables, root, subgroup Beet, garden and sugar Burdock Carrot Celeriac Chervil, turnip- rooted Chicory Ginseng Horseradish Parsley, turnip- rooted Parsnip Radish Radish, oriental Rutabaga Salsify Salsify, black Salsify, Spanish Skirret Turnip	Alternaria Leaf Spot Ascochyta Leaf Spot Rust White Rust  Cercospora Leaf Spot Powdery Mildew	1-2tsp.     1.25 -2tsp.	0	Apply every 7-14 days.     For Powdery Mildew apply on 5-7 day schedule.
	Soli-Borne Diseases Circular Spot, Southern Blight, Rhizoctonia Stem Canker, Crown Rot, Pythium Root Rot	½ - 7/8 for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.
Vegetables, tuberous and corm, subgroup Arracacha Arrowroot Artichoke, Chinese and Jerusalem Canna Cassava, edible, bitter and sweet Chayote (root) Chula Dasheen (Taro) Ginger	Foliar Diseases Alternaria Leaf Spot Ascochyta Leaf Spot Cercospora Leaf Spot Powdery Mildew Rust White Rust	1-2tsp.	14	Apply every 7-14 days. Use 1-2 tsp. rate for Cercospora leaf spot and Powdery Mildew. For Powdery Mildew apply on 5-7 day schedule.
	Soil-Borne Diseases Circular Spot, Southern Blight, Rhizoctonia Stem Canker, Crown Rot, Pythium Root Rot	½ - 7/8 tsp. for 200 feet of row		Apply after planting seeds by spraying over the row or directed to the base of transplants or emerged plants.

Plant	Disease	How Much to Apply (per gal. of water)	Days to Wait To Harvest	Comments
Leren Potato Sweet Potato Tanier Tumeric Yam, bean Yam, true				
Watercress	Cercospora Leaf Spot	1 - 2 tsp.	7	Apply on a 7-10 day schedule

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep in a cool place. Do not store at temperatures below 32°F. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### CONTAINER DISPOSAL:

Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

(non-refillable  $\leq 5$  gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(non-refillable  $>5$  gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container (250 gallon & bulk): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

#### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ALBAUGH, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold ALBAUGH, LLC and Seller harmless for any claims relating to such factors.

ALBAUGH, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or ALBAUGH, LLC, and, (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, ALBAUGH, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall ALBAUGH, LLC be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ALBAUGH, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ALBAUGH, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

ALBAUGH, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of ALBAUGH, LLC.