



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

September 2, 2025

Christina M. Swick
Agent for Albaugh, LLC
c/o Lewis & Harrison, LLC
2461 S. Clark Street, Suite 710
Arlington, VA 22202-3872

Subject: Label Amendment – Addition of Low Growing Berry Subgroup 13-07G, Updated Hotline Number and Booklet Statements, Addition of Optional Marketing Logos, Updated “Managing Mite Resistance” Section, Addition of Maximum Active Ingredient Rates and Various Other Label Changes
Product Name: HEXYTHIAZOX 1E
EPA Registration Number: 42750-311
Application Date: December 17, 2024
Action Case Number: 648798

Dear Christina M. Swick:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website

is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Carmen Rodia via e-mail at Rodia.Carmen@epa.gov.

Sincerely,

A handwritten signature in dark ink, appearing to read "Tamica Cain", is positioned above the typed name and title.

Tamica L. Cain, Product Manager 10
Invertebrate & Vertebrate Branch 2
Registration Division (7505T)

Enclosure: *Stamped "Accepted" Master Label, September 2, 2025*

{Note to reviewer: Text in braces { } is for informational purposes and will not appear on the end-use label.}

{Note to reviewer: Text in brackets [] is optional, and may or may not appear on the end-use label.}

{Text in parentheses () will appear on the end use label.}

HEXITHIAZOX	GROUP	10A	ACARICIDE
-------------	-------	-----	-----------

HEXYTHIAZOX 1E

ACTIVE INGREDIENT:

Hexythiazox


trans-5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide..... 11.80%

OTHER INGREDIENTS*: 88.20%

TOTAL 100.00%

*Contains petroleum distillate

HEXYTHIAZOX 1E contains 1.0 lb. active ingredient per gallon of formulated product



KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• DO NOT induce vomiting unless told to do so by the poison control center or doctor.• DO NOT give any liquid to the person.• DO NOT give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER: For non-emergency questions regarding exposure to this product, call 1-888-347-6732 (7 days/week, 24-hr/day). For medical emergencies, call the poison control center at 1-800-222-1222. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
NOTE TO PHYSICIAN: May pose an aspiration pneumonia hazard. Contains petroleum distillate.	
For 24-hour chemical spill, leak, fire, or accident response information, call CHEMTREC toll free at 1-800-424-9300.	

[See [inside] [side] [back panel] [booklet] [attached] [to] [pouch] [bag] [sealed pack] for [complete] [additional] [First Aid Statements,] [Precautionary Statements,] [Directions For Use,] [Storage and Disposal,] and [Conditions of Sale and Warranty.]]

EPA Reg. No. 42750-311

EPA Est. No. _____

NET CONTENTS: _____ Gallons

MANUFACTURED BY:

Albaugh, LLC
1525 NE 36th Street
Ankeny, IA 50021

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Contains petroleum distillate. Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes, or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Chemical-resistant gloves made of barrier laminate, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, or viton \geq 14 mils; and
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** contaminate water when cleaning equipment or when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT use, pour, spill or store near heat or open flame. **DO NOT** mix or allow contact with oxidizing agents. Hazardous reaction may occur.

PRODUCT INFORMATION

HEXYTHIAZOX 1E Miticide is recommended for use as a foliar application in labeled crops. HEXYTHIAZOX 1E Miticide provides effective control of Tetranychid spider mite species. It is an emulsifiable concentrate to be mixed with water and applied as a spray.

HEXYTHIAZOX 1E controls mites through activity on eggs and immature stages. Control is achieved from either direct contact with the spray or from contact with treated plant surfaces. Through its ovicidal activity, HEXYTHIAZOX 1E controls newly deposited mite eggs and eggs which are laid after application. HEXYTHIAZOX 1E is also highly effective in controlling immature motile stages of target mite species that are sprayed or move onto treated surfaces. Adult mites are not directly affected. However, eggs produced by females in contact with treated surfaces will be rendered nonviable. HEXYTHIAZOX 1E provides residual control of pest mite species. The degree and duration of control is dependent on the rate used, growth stage of the mite, species of mite, and climatic conditions under which the material is applied.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all directions on the EPA approved label and any supplemental labeling.

USE RESTRICTIONS ON ALL CROPS

- Apply this product only as specified the EPA-approved label and any supplemental labeling.
- **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- **DO NOT** graze or feed livestock on cover crops growing in treated areas.
- HEXYTHIAZOX 1E is an emulsifiable concentrate to be diluted with water for application in commercial plantings only.
- Apply HEXYTHIAZOX 1E prior to adult mite build up. HEXYTHIAZOX 1E will not control adult spider mites. Use higher rates on moderate-to-high mite infestations or for larger plants with a dense canopy. If adult mites are present in medium-to-high populations, better results may be obtained using HEXYTHIAZOX 1E in combination with a registered contact adulticide. The use of less than label rates is not recommended with HEXYTHIAZOX 1E.
- **DO NOT** make more than one application of HEXYTHIAZOX 1E or any other Hexythiazox product to the same crop per year.
- **DO NOT** plant rotational crops other than those on this label within 120 days of this application.
- **DO NOT** use in home plantings.
- For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals (REIs). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the REI of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over long-sleeved shirt and long pants;
- Chemical-resistant gloves made of barrier laminate, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, or viton \geq 14 mils; and
- Shoes plus socks

Managing Mite Resistance:

For resistance-management, this product contains a Group 10 acaricide. Any insect population may contain individuals naturally resistant to this product and other Group 10 acaricides. The resistant individuals may dominate the insect population if this group of acaricides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed. To reduce the potential for developing insect resistance, rotate to an acaricide with a different mode of action. Monitor treated pest populations for resistance development. Read product label before applying any acaricide and follow label directions.

To delay acaricide resistance, take the following steps:

- Rotate the use of this product or other Group 10 acaricides within a growing season, or among

growing seasons, with different groups that control the same pests. Avoid application of more than the maximum seasonal use rate or the total number of consecutive sprays of HEXYTHIAZOX 1E per season.

- Use tank mixtures with acaricides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual acaricides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance-management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance-management benefits of an acaricide mixture are greatest if the two components have similar periods of residual acaricidal activity. Mixtures of acaricides with unequal periods of residual acaricidal activity may offer an insect resistance management benefit only for the period where both acaricides are active.
- Adopt an integrated pest management program for acaricide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Albaugh, LLC representatives at 1-800-247-8013.

Mixing Directions: Turn on spray tank agitation. Pour recommended amount of this material on the surface of water in a nearly filled spray tank.

Add balance of water to fill the tank. Keep agitator running during filling and spraying operations. **DO NOT** allow mixture to stand.

Compatibility: HEXYTHIAZOX 1E is compatible with most agricultural chemicals. However, all possible combinations have not been evaluated. Before full-scale mixing of HEXYTHIAZOX 1E with other products, test mix small proportionate quantities of each to ensure compatibility.

Phytotoxicity Notice: As is common with most emulsifiable concentrate formulations, adverse effects such as spotting or discoloration of the treated surfaces can occur. Some conditions known to contribute to phytotoxicity include, but are not limited to, high temperatures, poor spray drying conditions, excessive spray deposit or run-off, certain spray mixtures, stage of crop development or tank mixes with other pesticides. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Application Information: Use sufficient spray volume to obtain thorough, uniform coverage of all plant surfaces. Apply using ground equipment, chemigation or air unless otherwise specified under the crop Use Directions.

Important: HEXYTHIAZOX 1E Miticide is a 1-pound active ingredient per gallon emulsifiable concentrate formulation to be diluted with water for application in commercial plantings only. **DO NOT** use in home plantings.

Use Recommendations: For best results, apply HEXYTHIAZOX 1E Miticide at first sign of mites before adult mite build up. HEXYTHIAZOX 1E Miticide will not control adult spider mites. Use higher labeled rates on moderate-to-high mite infestations, large trees with dense foliage, or for extended residual control. The lowest labeled rate of HEXYTHIAZOX 1E Miticide may be used in conjunction with other IPM strategies, or for shorter residual control later in the season. The use of less than label rates is not recommended since it will result in poor performance and contribute to resistance development. If adult mites are present in medium-to-high populations, better results may be obtained using HEXYTHIAZOX 1E Miticide in combination with a registered contact adulticide. HEXYTHIAZOX 1E Miticide may be tank mixed with horticultural grade oil. Refer to all precautions regarding phytotoxicity on the oil label. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

USE RESTRICTION: DO NOT make more than one application of HEXYTHIAZOX 1E Miticide per calendar year.

Chemigation Systems: HEXYTHIAZOX 1E may be applied through irrigation systems (chemigation) to any crop on the HEXYTHIAZOX 1E label unless otherwise specified under the crop Use Directions or on supplemental labeling supplied by Albaugh, LLC. **DO NOT** allow chemigation to run off field.

Types of Irrigation Systems: Apply HEXYTHIAZOX 1E only through sprinkler, including center pivot, lateral move, Low Energy Precision Applications (LEPA), end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. **DO NOT** apply HEXYTHIAZOX 1E through any other type of irrigation system.

USE DIRECTIONS FOR ALL RECOMMENDED TYPES OF IRRIGATION SYSTEMS

Uniform Water Distribution and System Calibration: The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. For best results, apply at 100% input/travel speed, for center pivots or 0.10 inch (2.716 gallons) up to 0.15 inch (4,073 gallons) of water/A, for other systems. Higher labeled rates of HEXYTHIAZOX 1E may be necessary for chemigation applications.

Chemigation Monitoring: A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift: DO NOT apply when wind speed favors drift beyond the area intended for treatment.

Required System Safety Devices: The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump; such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using Water from Public Water Systems: Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Cleaning the Chemical Injection System: In order to accurately apply pesticides, the chemical injection system must be kept clean; free of chemical or fertilizer residues and sediments. Refer to your owner's manual or ask your equipment supplier for the cleaning procedure for your injection system.

Flushing the Irrigation System: At the end of the application period, allow time for all lines to flush the pesticide through all nozzles or emitters before turning off irrigation water. To ensure the lines are flushed and free of pesticides, a dye indicator may be injected into the lines to mark the end of the application period.

Equipment Area Contamination Prevention: It is recommended that nozzles in the immediate area of control panels, chemical supply tanks, pumps and system safety devices be plugged to prevent chemical contamination of these areas.

Center-Pivot and Automatic-Move Linear Systems: Inject the specified dosage per acre continuously for one complete revolution or move of the system. **DO NOT** USE END GUNS. The system should be run at maximum speed.

Solid Set and Manually Controlled Linear Systems: Injection should be during the last 30 to 60 minutes of regular irrigation period or as a separate 30- to 60-minute application not associated with a regular irrigation. **DO NOT** USE END GUNS.

APPLICATION DIRECTIONS

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
ALFALFA (Western US Only*)	Two-spotted spider mite	10 - 24	<p>Ground Application: Apply the labeled rate of HEXYTHIAZOX 1E by ground equipment at 15 - 20 GPA.</p> <p>Air Application: Apply the labeled rate of HEXYTHIAZOX 1E by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary.</p> <p>Chemigation: See details under Chemigation Systems.</p>
			<p>USE RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply within 14 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. For use in the Western US only. <p>*May only be applied in the Western US - west of the line defined as follows: TX: W of Rt. 283 and NW of Rt. 377 OK: W of Rt. 281/183 KS, NE, SD, ND: W of Rt. 281</p>

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
TIMOTHY (West of the Rockies)	McDaniel spider mite, Two-spotted spider mite, Banks grass mite	10 - 24	<p>Ground Application: Apply the labeled rate of HEXYTHIAZOX 1E by ground equipment at 15 - 20 GPA.</p> <p>Air Application: Apply the labeled rate of HEXYTHIAZOX 1E by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary.</p> <p>Chemigation: See details under Chemigation Systems.</p>
			<p>USE RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply within 14 days of harvest for hay. • Animals may graze treated fields on the day of application. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year.

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
BEANS, DRY AND SUCCULENT (WESTERN US ONLY*)	Two-spotted spider mite, Pacific spider mite, Strawberry spider mite	10 - 24	<p>Ground Application: Apply a minimum of 20 gallons finished spray per acre.</p> <p>Air application: Apply a minimum of 10 gallons finished spray per acre.</p> <p>Chemigation: See details under Chemigation Systems.</p>
	<p>USE RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply within 14 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. • DO NOT harvest or graze bean vine forage or hay. <p>*May only be applied in the Western US - west of the line defined as follows: TX: W of Rt. 283 and NW of Rt. 377 OK: W of Rt. 281/183 KS, NE, SD, ND: W of Rt. 281</p>		

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
BERMUDAGRASS (West of the Rockies)	Banks grass mite, Two-spotted spider mite	12 - 24	<p>Apply at the first sign of mites before populations begin to build.</p> <p>Ground Application: Apply the labeled rate of HEXYTHIAZOX 1E by ground equipment at 15-20 GPA.</p> <p>Air Application: Apply the labeled rate by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary.</p> <p>Chemigation: See details under Chemigation Systems on Section 3 of the label.</p>
	<p>USE RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply within 7 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. 		

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
CITRUS FRUIT GROUP 10-10 (CA, AZ AND TX ONLY)	Tetranychid Mites (such as European Red Mite and Two- spotted Spider mite)	12 - 24	Application by ground equipment only. Use sufficient volume of water to obtain complete coverage but not less than 20 gallons per acre.
	USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 7 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. 		

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
COTTON	Two-spotted spider mite, Pacific spider mite, Strawberry spider mite, Carmine spider mite	12 - 20	Apply HEXYTHIAZOX 1E before boll opening and prior to adult mite buildup. Complete coverage of foliage is essential to optimal performance. Ground Application: Apply at labeled rates Air application: Use labeled per acre rate and apply a minimum of 20 gallons of finished spray per acre. Chemigation: See details under Chemigation Systems.
	USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 35 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 20 oz. (0.1562 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. 		

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
CORN, FIELD	Banks grass mite, Two-spotted spider mite, Pacific spider mite, Strawberry spider mite	6 - 24	Ground Application: Apply the labeled rate of HEXYTHIAZOX 1E with conventional spray boom equipment at 15 - 20 GPA. Thorough and uniform coverage of plants is required for optimal results. Air Application: Apply the labeled rate of HEXYTHIAZOX 1E by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: See details under Chemigation Systems.
	USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 30 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. • DO NOT harvest field corn forage, stover, and grain for 30 days after application. 		

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
CORN, SWEET (forage, kernel plus cob with husks removed) (Western US Only*)	Banks grass mite, Two-spotted spider mite, Pacific Spider Mite, Strawberry spider mite	10 - 24	Ground Application: Apply the labeled rate of HEXYTHIAZOX 1E with conventional spray boom equipment at a minimum of 20 gallons finished spray per acre. Thorough and uniform coverage of plants is required for optimal results. Air Application: Apply the labeled rate of HEXYTHIAZOX 1E by air in a minimum of 10 gallons finished spray per acre. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: See details under Chemigation Systems.
			USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. • DO NOT apply within 28 days of harvest of sweet corn or forage. *May only be applied in the Western US - west of the line defined as follows: TX: W of Rt. 283 and NW of Rt. 377 OK: W of Rt. 281/183 KS, NE, SD, ND: W of Rt. 281

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
CHRISTMAS TREE, CHRISTMAS TREE PLANTATIONS	Pacific Spider Mite, Spruce Spider Mite, Two-spotted Spider Mite	12 - 24	Ground Application: Apply at labeled rates Air application: Apply in 2-10 gallons finished spray per acre. For best coverage, apply in two opposing passes of 5 gallons each. Chemigation: See details under Chemigation Systems.
			USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year.

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
SMALL FRUIT VINE CLIMBING SUBGROUP 13- 07F, EXCEPT FUZZY KIWIFRUIT Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these	Tetranychid Mites (such as European Red Mite and Two- spotted Spider Mite), Pacific spider mite, Willamette mite	12 - 24	Ground Application: Apply at labeled rates. Use sufficient volume of water to obtain complete coverage but not less than 20 gallons per acre. Air Application: Apply a minimum of 10 gallons finished spray per acre. Chemigation: See details under Chemigation Systems.
			USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 7 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year.

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
GREENHOUSE TOMATOES	Tetranychid Mites (such as Two-spotted Spider Mite)	12 - 24	Use sufficient volume of water to obtain complete coverage but not less than 20 gallons per acre.
			USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 1 day of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year.

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
HOPS	Tetranychid Mites (such as Two-spotted Spider Mite)	12 - 24	Ground Application: Apply a minimum of 20 gallons of finished spray per acre. Air Application: Use sufficient volume of water to obtain complete coverage but not less than 10 gallons per acre.
			USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 30 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year.

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
LOW GROWING BERRY SUBGROUP 13-07G[*] Bearberry; bilberry; blueberry, lowbush; cloudberry; cranberry; lingonberry; muntries; partridgeberry; strawberry; cultivars, varieties, and/or hybrids of these	Two-spotted Spider Mites	12 - 24	Ground Application: Apply a minimum of 20 gallons of finished spray per acre. It is recommended to use a silicone surfactant to increase canopy coverage. Air Application: Apply a minimum of 20 gallons finished spray per acre. Chemigation: See details under Chemigation Systems
			USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 3 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than 32 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. • DO NOT use oil. [* Not Registered for Use by California]

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
NON-BEARING TREE FRUITS and NUTS, CITRUS, and VINES (FIELD GROWN AND NURSERY) Non-bearing crops are perennial crops that will not produce a harvestable raw agricultural commodity during the season of application	Citrus Red Mite, European Red Mite, Two-spotted Spider Mite, McDaniel Spider Mite, Pacific Spider Mite, Six-spotted Mite, Texas Citrus Mite	12 - 24	Ground Application: Apply at labeled rates Air Application: Apply a minimum of 5 gallons finished spray per acre. Chemigation: See details under Chemigation Systems.
			USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year.

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
PEPPER/EGGPLANT SUBGROUP 8-10B African eggplant; bell pepper; eggplant; martynia; nonbell pepper; okra; pea eggplant; pepino; roselle; scarlet eggplant; cultivars, varieties, and/or hybrids of these	Tetranychid Mites (such as Two-spotted spider mite), Pacific spider mite, Strawberry spider mite	12 - 24	Ground Application: Apply a minimum of 20 gallons of finished spray per acre. Air Application: Apply a minimum of 2 gallons finished spray per acre. Chemigation: See details under Chemigation Systems.
			USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 1 day of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year.

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
POME FRUIT GROUP 11-10 Apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these	European red mite, Two-spotted spider mite, McDaniel spider mite, Pacific spider mite, Willamette mite	12 - 24	Ground Application: Use sufficient volume of water to obtain complete coverage but not less than 20 gallons per acre. Air Application: Use sufficient volume of water to obtain complete coverage but not less than 10 gallons per acre. Chemigation: See details under Chemigation Systems.
			USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 28 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. • For D'Anjou or Bosc pear varieties, use higher labeled rates (16-24 oz. (0.125-0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre).

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
POTATOES (Western US Only*)	Two-spotted Spider Mites; Potato Psyllid	12 - 24	Ground Application: Apply at labeled rates Air Application: Apply a minimum of 5 gallons finished spray per acre. Chemigation: See details under Chemigation Systems.
			USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 21 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. *May only be applied in the Western US - west of the line defined as follows: TX: W of Rt. 283 and NW of Rt. 377 OK: W of Rt. 281/183 KS, NE, SD, ND: W of Rt. 281

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
SORGHUM (Western US only, including Oklahoma and Texas*)	Banks grass mite, Two-spotted spider mite	10 - 24	Ground Application: Apply the labeled rate of HEXYTHIAZOX 1E by ground equipment at 15 - 20 GPA. Air Application: Apply the labeled rate of HEXYTHIAZOX 1E by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: See details under Chemigation Systems
			RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 30 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. *May only be applied in the Western US - west of the line defined as follows: KS, NE, SD, ND: W of Rt. 281 Entire States of OK and TX included for use

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
STONE FRUITS GROUP 12 <i>(including Plums)</i>	European red mite, Two-spotted spider mite, McDaniel spider mite, Strawberry spider mite, Pacific spider mite, Pecan leaf scorch mite, Willamette mite	12 - 24	Ground Application: Apply a minimum of 20 gallons finished spray per acre. Air Application: Apply a minimum of 10 gallons finished spray per acre. Chemigation: See details under Chemigation Systems.
	USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 7 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. 		

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
SUGAR BEETS	Banks grass mite, Two-spotted spider mite	12 – 24	Apply at the first sign of mites before populations begin to build. Ground Application: Apply the labeled rate of HEXYTHIAZOX 1E by Ground equipment at 15-20 GPA. Air Application: Apply the labeled rate by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. Chemigation: See details under Chemigation Systems.
	USE RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply within 45 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. 		

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
TREE NUTS GROUP 14 and pistachio	European red mite, Two-spotted spider mite, McDaniel spider mite, Strawberry spider mite, Pacific spider mite, Pecan leaf scorch mite, Willamette mite, Citrus Red Mite	12 - 24	<p>Ground Application: Apply a minimum of 20 gallons finished spray per acre.</p> <p>Air Application: Apply a minimum of 5 gallons finished spray per acre.</p> <p>Chemigation: See details under Chemigation Systems.</p>
			<p>USE RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply within 7 days of harvest. • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year.

CROP	PEST	RATE FL OZS/ACRE	USE DIRECTIONS
WHEAT, including emmer wheat and tritcale (West of the Rockies)	Banks grass mite	10-24	<p>Apply at the first sign of mites before populations begin to build.</p> <p>Ground Application: Apply the labeled rate of HEXYTHIAZOX 1E by ground equipment at 15-20 GPA.</p> <p>Air Application: Apply the labeled rate by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary.</p> <p>Chemigation: See details under Chemigation Systems.</p>
			<p>USE RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT make more than one application of HEXYTHIAZOX 1E or any other hexythiazox-containing product per calendar year. • DO NOT apply more than a total of 24 oz. (0.1875 lb. hexythiazox) of HEXYTHIAZOX 1E per acre per calendar year. • DO NOT apply after the boot stage of growth. • DO NOT harvest hay or forage for at least 14 days after application.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: DO NOT store near heat or open flame. Keep this product in its tightly closed original container. Store in a cool, dry (preferably locked) area that is away from other pesticides, fertilizers, food, or feed, in a place that is inaccessible to children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer empty container for recycling if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

FOR BULK AND MINI-BULK CONTAINERS

CONTAINER HANDLING: Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Before refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices. REFILL ONLY WITH HEXYTHIAZOX 1E. The contents of this container cannot be completely removed by cleaning. Refilling with materials other than HEXYTHIAZOX 1E will result in contamination and may weaken container. After filling and before transporting, check for leaks. **DO NOT** refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

FOR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL **CHEMTREC®** at **1-800-424-9300**, 24 hours per day, 7 days per week.

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS


Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our recommendations for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Albaugh, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer and User.

Albaugh, LLC warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALBAUGH, LLC MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALBAUGH, LLC's EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT ALBAUGH, LLC's SOLE DISCRETION.

[OPTIONAL MARKETING GRAPHICS]



090225

[LABEL HISTORY

Not included in final printed label

File Name	Version Mark	Comment
042750-00311.20190617.DRAFT	061719	Label Amend add crops expand crop geographies
042750-00311.20210429.DRAFT	042921	Label Amend add crops expand crop geographies revisions
042750-00311.20210504.MASTER	AD050421	EPA Acceptance
042750-00311.20241202.DRAFT	120224	Add Low Growing Berry Subgroup 13-07
042750-00311.20250828.DRAFT	082825	(e) Label Revision
042750-00311.20250902.DRAFT	090225	(e) Label Revision

]