



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

42750-336

Date of Issuance:

11/15/18

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

ST Pre-Mix #8

Name and Address of Registrant (include ZIP Code):

Morris Gaskins
Albaugh, LLC
P.O. Box 2127
Valdosta, GA 31604

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.

Signature of Approving Official:

Hope Johnson, Product Manager 21
Fungicide Branch, Registration Division (7505P)

Date:

11/15/18

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 42750-336.”
 - Add Net Contents information
4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 03/05/2018

If you have any questions, please contact Marcel Howard by phone at (703)305-6784, or via email at howard.marcel@epa.gov.

Enclosure

METALAXYL	GROUP	4	FUNGICIDE
TOLCLOFOS-METHYL	GROUP	14	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

ST PRE-MIX # 8

Broad-spectrum seed treatment fungicide for protection against seed and seed-borne diseases in soybeans, specified legumes, cotton and cereals (barley, oats, triticale, wheat).

ACTIVE INGREDIENTS:

Fludioxonil:* 1.86%
Metalaxyl**..... 11.63%
Tolclofos-methyl*** 7.20%

OTHER INGREDIENTS: 79.31%

TOTAL: 100.00%

* CAS No. 131341-86-1

** CAS No. 57837-19-1

***CAS No. 57018-04-9

Contains 0.16 pound active ingredient of fludioxonil, 1.03 pound active ingredient of metalaxyl and 0.64 pound active ingredient tolclofos-methyl per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER - Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For chemical emergency: spill, leak, fire, exposure or accident, call CHEMTREC: 1-800-424-9300.	
NOTE TO PHYSICIAN: Treatment is supportive and symptomatic. This material contains an atypical weak cholinesterase inhibitor. If signs of cholinesterase inhibition appear, atropine sulfate is antidotal. 2-PAM (PROTOPAM) is also antidotal and may be used in conjunction with atropine but should not be used alone. Other non-cholinergic signs and symptoms should be treated symptomatically.	


See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 42750-

EPA Est. No. 42750-MO-002

NET CONTENTS: _____ Gallons

Manufactured By:
ALBAUGH, LLC
ANKENY, IA 50021



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (barrier laminate, butyl rubber ≥ 14 mils, natural rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton[®] ≥ 14 mils).
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet.
2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

PHYSICAL AND CHEMICAL HAZARDS

Do not mix or allow contact with oxidizing agents. Hazardous chemical reactions may occur.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater and estuarine/marine fish, oyster and shrimp and aquatic invertebrates. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory Statement: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Fludioxonil has properties and characteristics associated with chemicals detected in groundwater. Use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Use is permitted on-farm and in commercial seed treatment facilities. Do not use for at-plant applications (e.g. hopper box, planter box, etc.). This product is to be applied as a water-based slurry through standard slurry- or mist-type commercial seed treatment equipment.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

1. Coveralls
2. Chemical-resistant gloves (barrier laminate, butyl rubber ≥ 14 mils, natural rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils)
3. Shoes plus socks

Failure to follow the directions for use and precautions on this label may result in crop injury, poor disease control, or illegal residues.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination test on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed-quality conditions beyond the control of Albaugh, LLC, no claims are made to guarantee germination of carry-over seed.

PRODUCT INFORMATION

ST PRE-MIX # 8 is a seed treatment fungicide that protects against damping-off and seed rots due to *Pythium*, *Phytophthora*, *Fusarium*, *Rhizoctonia* spp., and early season *Phytophthora* root rot. ST PRE-MIX # 8 also suppresses seed-borne *Sclerotinia* and *Phomopsis* spp., particularly in soybeans. ST PRE-MIX # 8 contains the active ingredient metalaxyl that provides systemic protection against *Pythium* and early season *Phytophthora* root rot. *Helminthosporium*, and weakly pathogenic fungi as *Aspergillus* and *Penicillium*.

Tolcolfos-methyl provides protection against *Rhizoctonia solani* and seed borne and soil borne fungal pathogens that cause seed decay, damping off and seedling blight.

ST PRE-MIX # 8 also contains the active ingredient fludioxonil that provides protection against certain soil-borne and seed-borne diseases of crop plants. Fludioxonil is active against *Fusarium*, *Rhizoctonia*, *Helminthosporium*, and weakly pathogenic fungi such as *Aspergillus* and *Penicillium*.

RESISTANCE MANAGEMENT

METALAXYL	GROUP	4	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE
TOLCLOFOS-METHYL	GROUP	14	FUNGICIDE

For Resistance Management, ST PRE-MIX #8 contains metalaxyl, a Group 4 fungicide; fludioxonil a Group 12 fungicide and Tolclofos-methyl, a Group 14 fungicide.

Any fungal population may contain individuals naturally resistant to ST PRE-MIX #8 and other Group 4 or Group 12 or Group 14 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields.

Because resistance development cannot be predicted, the use of this product should follow appropriate resistance-management strategies. To delay fungicide resistance, use should be based on an IPM program that includes the following steps:

- Rotate the use of ST PRE-MIX #8 with other fungicide Groups within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Albaugh, LLC at **1-800-247-8013**. You can also contact your pesticide distributor or university extension specialist to report resistance.

All applications must be made according to the use directions that follow. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

RESTRICTIONS:

ST PRE-MIX #8 must be used only in accordance with use instructions on this label utilizing mechanical, slurry, or mist type seed treating equipment. Calibrate the equipment to accurately and uniformly apply the product to seed. This product is for both commercial and on-farm application.

- Do not apply this product in hopper-box or planter-box at planting time. Always mix product thoroughly before use.
- ST PRE-MIX #8 may only be used on true seeds and may not be used on propagative plant parts such as garlic cloves.
- The maximum number of applications per season is 2

- Do not plant treated seeds by broadcast.
- For all treated crops, do not exceed seeding rate of 200 lb/A.

MIXING INSTRUCTIONS FOR SOYBEANS

ST PRE-MIX # 8 is formulated for on-farm or commercial treatment to be used with liquid rhizobia products, using standard mechanical slurry or mist-type seed treatment equipment which accurately measures and mixes a flowable seed treatment. The equipment must provide uniform coverage of ST PRE-MIX # 8 on the seed. Uneven seed coverage may not give the desired level of disease control. ST PRE-MIX # 8 must be diluted with either water or liquid inoculants prior to application to soybeans. Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment.

Do not mix ST PRE-MIX # 8 fungicide with other pesticides unless sufficient compatibility, loading rates, and seed and crop safety factors are determined.

In a clean container large enough to contain the entire mixture add the desired amount of ST PRE-MIX # 8. Add half of the required water or liquid inoculant and mix thoroughly. Add the remaining amount of water or liquid inoculant and mix again. The slurry is ready for application to the soybeans. Mix as needed and do not store diluted mixture.

Additional or continuous agitation or mixing of the slurry mixture may be necessary to prevent the ST PRE-MIX # 8 from settling out of solution when using water or liquid inoculants.

ST PRE-MIX # 8 has been found to be compatible with some liquid inoculant products. ST PRE-MIX # 8 may be mixed or applied sequentially with approved liquid inoculants. Consult the maker of the liquid inoculants and a ALBAUGH, LLC representative for directions before applying ST PRE-MIX # 8 with liquid inoculants.

Precaution:

ST PRE-MIX # 8 is formulated to be compatible with some liquid inoculants. If water is used to further dilute ST PRE-MIX # 8 when rhizobia are present, use only non-chlorinated water. Chlorinated water will kill the live rhizobia contained in the inoculant.

MIXING INSTRUCTIONS FOR SPECIFIED LEGUMES

Apply ST PRE-MIX # 8 as a water based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease protection. Thoroughly mix the specified amount of ST PRE-MIX # 8 into the required amount of water for the slurry treater and dilution rate to be used.

Follow the manufacturer application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Allow seed to dry before bagging.

CROP USE DIRECTIONS

SEED CONTAINER LABEL REQUIREMENT

The Federal Seed Act requires that containers of treated seeds shall be labeled with the following statements:

- This seed has been treated with metalaxyl, fludioxonil, and tolclofos-methyl fungicides.
- Do not use for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on containers of seeds treated with ST PRE-MIX #8:

- Ground Water Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil has properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- Excess treated seed may be used for ethanol production only if:
 1. By-products are not used for livestock feed, and
 2. No measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.
- Do not allow children, pets, or livestock to have access to treated seed.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Dispose of all excess treated seed by burying away from bodies of water.
- Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Do not allow animals to graze the following crops grown from seeds treated with fludioxonil until 30 days after planting:
 - Forage, Fodder, and Straw of Cereal Grains - Crop Group 16
 - Grass Forage, Fodder, and Hay - Crop Group 17
 - Nongrass Animal Feeds (Forage, Fodder, Straw, and Hay) - Crop Group 18
- For all treated crops except for cotton and root vegetable, do not exceed seeding rate of 200 lb/A. The maximum allowed seeding rates for cotton and root vegetable are 40 lb/A.
- Do not apply more than a total of 0.02 lb Tolclofos-methyl (active ingredient) per acre per cropping cycle.

All seed must be adequately dyed with a suitable colorant to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, part 2.25. Any colorant added to treated seed must be cleared for use under 40 CFR, part 153.155.

Crop	Target Diseases	Use Rate (fl. oz./100 lb seed)
Soybeans	Seed rot, seedling blight, pre and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Rhizoctonia</i> spp. and <i>Pythium</i> spp. Early season seedling root rot caused by <i>Fusarium</i> spp. Early season <i>Phytophthora</i> spp. Seedling blight and seed rot caused by <i>Phomopsis</i> spp. Suppression of seed-borne <i>Sclerotinia</i>	2.0
Legumes Chickpea (garbanzo bean), Lentils, *Peas (<i>Pisum</i> spp.), **Beans (dry &	Seed rot, seedling blight, pre and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Rhizoctonia</i> spp. and <i>Pythium</i> spp.	2.0

Crop	Target Diseases	Use Rate (fl. oz./100 lb seed)
succulent)		
Cotton	Pre and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Rhizoctonia</i> spp. and <i>Pythium</i> spp.	2.0
Cereals wheat (winter & spring), barley, oats, & triticale	Seed rot, seedling blight, pre and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Rhizoctonia</i> spp. and <i>Pythium</i> spp	2.0
2.0 fluid ounces of ST PRE-MIX #8 contains the following active ingredient: Metalaxyl: 0.0161 lbs Fludioxonil: 0.0025 lbs Tolclofos-methyl: 0.010 lbs		

***Peas:** (dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, dry pea, snow pea, sugar snap pea) Pigeon pea Sword bean.

****Beans:** Bean (*Lupinus* spp.) grain lupin, sweet lupin, white lupin, white sweet lupin

Bean (*Phaseolus* spp.) field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean

Bean (*Vigna* spp.) adzuki bean, asparagus bean, black-eyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean)

Broad bean (fava bean)

Guar

Jackbean

If the target soybean fields have a history of high *Phytophthora* disease pressure then use a tank mix with additional metalaxyl seed treatment product. Alternatively, approved liquid rhizobia inoculants at rates recommended by manufacturer to achieve proper total slurry volumes can be used.

Where mefenoxam/metalaxyl products used allow the same maximum poundage of active ingredient per acre per calendar year:

If more than one product containing mefenoxam/metalaxyl active ingredient is used on an acre during the same calendar year and the mefenoxam/metalaxyl products used allow the same maximum poundage of active ingredient per acre per calendar year, then the total poundage of all such mefenoxam/metalaxyl products used must not exceed any of the specified individual mefenoxam/metalaxyl product maximum poundage of active ingredient allowed per acre per calendar year.

Where mefenoxam/metalaxyl products used allow different maximum poundage of active ingredient per acre per calendar year:

If more than one product containing mefenoxam/metalaxyl active ingredient is used on an acre during the same calendar year and the mefenoxam/metalaxyl products used allow different maximum poundage of active ingredient per acre per calendar year, then the total poundage of all such mefenoxam/metalaxyl products used must not exceed the lowest specified individual mefenoxam/metalaxyl product maximum poundage of active ingredient allowed per acre per calendar year.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original containers only in a cool, dry place. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ALBAUGH, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold ALBAUGH, LLC and Seller harmless for any claims relating to such factors.

ALBAUGH, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or ALBAUGH, LLC, and (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, ALBAUGH, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall ALBAUGH, LLC be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ALBAUGH, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ALBAUGH, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

ALBAUGH, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of ALBAUGH, LLC.