



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

42750-345

Date of Issuance:

4/2/19

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

ST Pre-Mix #13

Name and Address of Registrant (include ZIP Code):

Morris Gaskins
 Albaugh LLC
 P.O. Box 2127
 Valdosta, GA 31604

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Sincerely,

Venus Eagle, Product Manager 01
 Invertebrate & Vertebrate Branch 3
 Registration Division (7505P)
 Office of Pesticide Programs

Date:

4/2/19

2. You are required to comply with the data requirements described in the DCI identified below:
 - a. Imidacloprid GDCI: 129099-951
 - b. Metalaxyl GDCI: 113501-869
 - c. Tebuconazole GDCI: 128997-1598

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.
4. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 42750-345.”
5. Submit one copy of the final printed label for the record before you release the product for shipment.


Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 9/21/2018

If you have any questions, please contact Jasmin Jackson by phone at 703-347-0732, or via email at jackson.jasmin@epa.gov.

Enclosure



Metalaxyl	GROUP	4	FUNGICIDE
Tebuconazole	GROUP	3	FUNGICIDE
Imidacloprid	GROUP	5	INSECTICIDE

ST PRE-MIX #13
Seed Treatment

For early season protection of seedlings against insect injury and soilborne diseases for Barley, Wheat and Oats.

ACTIVE INGREDIENTS:

Imidacloprid: 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	11.38%
Metalaxyl: <i>N</i> -(2,6-dimethylphenyl)- <i>N</i> -(methoxyacetyl) alanine methyl ester	0.61%
Tebuconazole:	
Alpha-[2-(4-Chlorophenyl) ethyl]-alpha-(1, 1-Dimethylethyl)-1H-1,2,4-triazole-1-ethanol	0.45%
OTHER INGREDIENTS:	<u>87.56%</u>
TOTAL:	100.00%

Contains 1.00 pound Imidacloprid per gallon (120.0 grams per liter), 0.05 pound Metalaxyl per gallon (6.4 grams per liter), 0.04 pound Tebuconazole per gallon (4.8 grams per liter)

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continuing rinsing. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER: Have the product container or label with you when calling poison control center or doctor, or going for treatment. For 24- Hour Emergency Assistance spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300	

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 42750-GUL

EPA Est. No. 42750-MO-002

NET CONTENTS: ____ Gallons

Lot No. _____

Manufactured By:
ALBAUGH, LLC
ANKENY, IA 50021

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Avoid contact with eyes, skin or clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks
- Chemical-resistant gloves (Barrier Laminate, Butyl Rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils, Neoprene Rubber ≥ 14 mils, Polyvinyl Chloride (PVC) ≥ 14 mils, Viton ≥ 14 mils).

Persons involved in bagging treated seed, sewing or moving bags of treated seed, or cleaning up bagging areas or seed treatment equipment are pesticide handlers and must wear the PPE required on this label for pesticide handlers.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS
Users should: <ol style="list-style-type: none">1. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish and aquatic invertebrates. For terrestrial use, DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment wash water or rinsate. Cover or incorporate spilled treated seeds. Exposed treated seeds may be hazardous to birds.

Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PHYSICAL AND CHEMICAL HASZARDS

Do not mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

Coveralls, Chemical-resistant gloves (Barrier Laminate, Butyl Rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils, Neoprene Rubber ≥ 14 mils, Polyvinyl Chloride (PVC) ≥ 14 mils, Viton ≥ 14 mils, Shoes plus socks.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. The following plant-back intervals must be observed for listed crops.

ROTATIONAL PLANT-BACK INTERVALS*				
Immediate Plant-Back				30-Day Plant-Back
Artichoke Barley Borage Brassica (cole) leafy vegetable Bulb vegetables Canola Cilantro Corn, Field Corn, Sweet Cotton Cranberry	Crambe Cucurbits Eggplant Flax Groundcherry Leafy petiole vegetables Leafy vegetables Legume vegetables: Succulent or dried Including Soybean Millet	Mustard Seed Oats Okra Pepinos Pepper Popcorn Potato Rapeseed Rye Safflower Sorghum	Soybean Strawberry Sugarbeet Sunflower Tomatillo Tomato Triticale Root and Tuber vegetables Watercress Wheat	Cereals, including buckwheat rice
<p>*Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.</p> <p>*For all other crops not listed on an imidacloprid label, or for crops for which no tolerance for the active ingredient has been established, a 12-month plant-back interval must be observed.</p>				

Notification of the crop rotational restriction must be conveyed to the grower, by appropriate seed tag or bag printing on all seed units.

USE RESTRICTIONS

- **DO NOT** use as a planter (hopper) box treatment.
- **DO NOT** use treated seed for feed, food, or oil purposes.
- Store treated seed away from feeds and foodstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- The maximum application rate for imidacloprid (including seed treatments, foliar applications, soil applications) is 0.5 lbs. per acre per year. **DO NOT** apply more than 0.5 lbs. ai imidacloprid per acre per year.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Exposed treated seed may be hazardous to birds. Cover or incorporate spilled treated seed. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** graze or feed livestock on treated areas for 45 days after planting.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash water.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

SEED TREATMENT

This product is a combination of Imidacloprid insecticide and a mixture of the fungicides Metalaxyl, Tebuconazole, and Fludioxonil. When used on listed crops, the fungicide mixture in this product provides consistent control when disease pressure is high. This product aids in the control or suppression of the following insects, seed, seedling and soilborne diseases of Barley, Wheat, and Oats.

CROP	INSECT / DISEASE CONTROL	RATE
Barley	<p>Insect Control: Early season protection of seedlings against injury by Aphids (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid) and Wireworms.</p> <p>Disease Control: Early season disease control of <i>Pythium</i> damping-off and Covered smut, Loose smut, Barley stripe (suppression), Early season <i>Rhizoctonia</i> root rot, Early season common root rot, Early season <i>Fusarium</i> foot rot, Early season suppression of powdery mildew and Early season suppression of barley leaf rust.</p>	3.4 to 5.0 fluid ounces per 100 pounds of seed
Wheat and Oats	<p>Insect Control: Early season protection of seedlings against injury by Aphids (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid), Wireworms and Hessian fly.</p> <p>Disease Control: Early season disease control of <i>Pythium</i> damping-off, Stinking smut, Flag smut, Loose smut, Early season <i>Septoria</i> disease complex, Early season <i>Rhizoctonia</i> root rot, Early season common root rot, Early season <i>Fusarium</i> foot rot, Early season suppression of powdery mildew and early season suppression of wheat leaf rust.</p>	
<p>APPLICATION DIRECTIONS: Apply prior to planting as a slurry treatment. This product is to be used in liquid or slurry treaters only. Mix thoroughly before use or use entire container at one time. Ensure thorough coverage. All tank mixes should be pre-tested to determine physical compatibility between formulations.</p> <p>It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.</p> <p>RESTRICTIONS: DO NOT graze or feed livestock on treated areas for 45 days after planting.</p>		

SEED BAG LABELING REQUIREMENTS

Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act. The user of this product is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

THE FEDERAL SEED ACT REQUIRES THAT BAGS CONTAINING TREATED SEEDS BE LABELED WITH THE FOLLOWING STATEMENTS:

- This seed has been treated with ST PRE-MIX #13 (imidacloprid, metalaxyl, tebuconazole).
- **DO NOT** use for feed, food, or oil purpose.

THE U.S. ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON BAGS CONTAINING SEEDS TREATED WITH ST PRE-MIX #13 (imidacloprid, metalaxyl, tebuconazole):

- **Pollinator Precautions:** Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- **Groundwater Advisory:** Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- Store away from feeds and foodstuffs.

- Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Exposed treated seed may be hazardous to birds. Cover or incorporate spilled treated seeds. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** graze or feed livestock on treated areas for 45 days after planting.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash water.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- In the event of a crop failure or harvest of a crop grown from ST PRE-MIX #13 treated seed, the field may be replanted immediately to artichoke, barley, borage, Brassica (cole) leafy vegetables, bulb vegetables, canola, cilantro, corn (field or sweet), cotton, cranberry, crambe, cucurbits, eggplant, flax, groundcherry, leafy petiole vegetables, leafy vegetables, legume vegetables (succulent or dried including soybean), millet, mustard seed, oats, okra, pepinos, pepper, popcorn, potato, rapeseed, rye, safflower, sorghum, soybean, strawberry, sugarbeets, sunflower, tomatillo, tomato, triticale, root and tuber vegetables, watercress, and wheat. For cereals including buckwheat, and rice; the minimum plant-back interval is 30 days from the date ST PRE-MIX #13 treated seed was planted. Cover crops for soil building or erosion control may be planted at any time; but do not graze or harvest for food or feed. For all other crops not listed on an imidacloprid label, or for crops for which no imidacloprid, tolerance for the active ingredient has been established, a 12-month plant-back interval must be observed.
- The maximum application rate (including seed treatment, foliar application, and soil application) per acre per year for imidacloprid is 0.5 lbs.
- This seed has been treated with X lbs. imidacloprid, X lbs. metalaxyl, X lbs. tebuconazole, X lbs. of seed [or X mg per seed].
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

NOTE: If this product is undyed, the purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as a food for man or feed for animals. Refer to 21 CFR, Part 2.25. Any dye added to treated seed must be cleared for use under 40 CFR, Part 180.1001.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Store in a cool, dry place, **DO NOT** store in direct sunlight. Protect from freezing temperatures (<25° F) and hot temperatures (> 90° F). After prolonged storage in intense cold, allow product to return to approximately 75 deg. F before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[Nonrefillable Containers 5 Gallons or Less]

Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic

containers are also disposable by incineration.

[or]

[Nonrefillable containers larger than 5 gallons]

Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[or]

[Refillable containers larger than 5 gallons for return to Albaugh]

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Close all openings and replace all caps. Contact Albaugh's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

[or]

[Refillable containers for disposal]

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times. For final disposal, offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1)THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BYWAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BYTHE BUYER, USER, OR ITS CUSTOMERS. TOTHE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.