



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

January 5, 2023

Blake Cowen
Product Registration Manager
Albaugh, LLC
PO Box 2127
Valdosta, GA 31604

Subject: Registration Review Label Amendments Incorporating Mitigation Measures from the Interim Decisions for Metolachlor and Mesotrione and the National Marine Fisheries Services' (NMFS) Biological Opinion on the Effects of Metolachlor on Pacific Salmonids
Product Name: Metolachlor + Mesotrione SC
EPA Registration Number: 42750-346
Application Dates: 8/25/2021 and 04/19/2021
Decision Numbers: 573336; 583842 and 589296

Dear Blake Cowen:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Metolachlor and Mesotrione Interim Decisions. The Agency has concluded that your submission is acceptable.

This letter also addresses the label mitigation resulting from the NMFS' Biological Opinion on the effects of Metolachlor on Pacific salmonids. The Agency has concluded that your submission is also acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


If you have any questions about this letter, please contact Srijana Shrestha at shrestha.srijana@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington", with a stylized flourish at the end.

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure



Metolachlor	Group	15	Herbicide
Mesotrione	Group	27	Herbicide

METOLACHLOR + MESOTRIONE SC
 [Metolachlor + Meso]
 [Priority Meso]

Herbicide for pre-emergence and post-emergence control of grass and broadleaf weeds
 In Corn (field, seed, sweet and yellow popcorn), and Grain Sorghum

ACTIVE INGREDIENTS*:	By Wt.
Metolachlor	36.80%
Mesotrione	3.68%
OTHER INGREDIENTS:	59.52%
TOTAL:	100.00%

*Equivalent to 3.26 lbs. per gal. metolachlor and 0.33 lb. per gal. mesotrione

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. Emergency Phone Numbers: CHEMTREC 1-800-424-9300 (transportation and spills)	

Optional Language that may appear on the label:

[See additional Precautionary Statements and Directions for Use inside [the] booklet]

[See inside booklet for additional [complete] [First Aid,] Precautionary Statements and Directions For Use.]

EPA Reg. No. 42750-346

EPA Est. No. xxxxx-xx-xxx

NET CONTENTS: _____ Gallons

MANUFACTURED BY:
 Albaugh, LLC
 Ankeny, IA 50021

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All Mixers, Loaders, Applicators, and other handlers must wear:

1. Coveralls over short-sleeved shirt and short pants
2. Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride ≥ 14 mils, or Viton ≥ 14 mils.
3. Chemical-resistant footwear plus socks
4. Chemical-resistant headgear for overhead exposure
5. Chemical-resistant apron when cleaning equipment, mixing or loading.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

REPORTING ECOLOGICAL INCIDENTS

To report ecological incidents, including mortality, injury, or harm to plants and animals, call 1 (800)247-8013.

GROUND WATER ADVISORY

Metolachlor is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

SURFACE WATER ADVISORY

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks or months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of metolachlor from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

The active ingredients in this product have the potential to contaminate surface water through ground spray drift. Under some conditions, the active ingredients may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These include poorly drained or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

Sound erosion control practices will reduce this product's contribution to surface water contamination.

MIXING/LOADING INSTRUCTIONS

This product must be used in a manner that will prevent back siphoning into wells and prevent spills.

Dispose of excess pesticide, spray mixtures or rinsates properly.

Mixing equipment must have check valves or anti-siphoning devices in use.

Do not mix or load this product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This restriction does not apply to plugged abandoned well or wells that are properly capped and does not apply to impervious pads or mixing/loading areas that are properly diked.

Mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well is strictly prohibited unless on an impervious pad constructed to withstand the weight of the heaviest load that could be on or moved across the pad.

The pad must be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rainwater that may fall on the pad. Surface water must not be allowed to flow over or from the pad. To facilitate material removal, the pad must be sloped. A pad that is not under cover must have capacity to hold a minimum of 110% of the capacity of the largest pesticide product container or application equipment that will be on the pad. Covered pads that are completely protected from precipitation must have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment that will be on the pad.

The containment capacities must be specified and maintained at all times. Minimum specific containment capacities do not apply to vehicles that deliver pesticides to the mixing/loading site. There may be additional state requirements regarding containment and well setback restrictions. Consult local authorities for additional information.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not mix or allow contact with oxidizing agents, as a

hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Failure to follow the DIRECTIONS FOR USE, RESTRICTIONS and PRECAUTIONS on this label may result in reduced weed control, adverse crop response, or illegal crop residues.

NOTE: Not for sale, distribution or use in Nassau or Suffolk Counties in New York.

ENDANGERED SPECIES

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/espp/>, call 1-844-447-3813, or email ESPP@epa.gov. You must use the Bulletin valid for the month in which you will apply the product.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

The following PPE is required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water:

1. Coveralls over short-sleeved shirt and short pants
2. Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride \geq 14 mils, or Viton \geq 14 mils.
3. Chemical-resistant footwear plus socks
4. Chemical-resistant headgear for overhead exposure

PRODUCT INFORMATION

METOLACHLOR + MESOTRIONE SC is for use in field corn and seed corn for pre-emergence and early post-emergence control of many annual grass and broadleaf weeds.

METOLACHLOR + MESOTRIONE SC may also be applied to sweet corn, yellow popcorn and grain sorghum as pre-emergence control of many annual grass and broadleaf weeds.

Refer to the Weeds Controlled tables for lists of weeds. This product must be used before weeds emerge to effectively control most grass weeds.

If applications are made according to labeled directions for use and under normal growing conditions, METOLACHLOR + MESOTRIONE SC will not cause crop injury to the treated crop. During germination and early stages of growth, environmental conditions or other factors that contribute to stress of the crop may cause poor or slow growth and may weaken crop seedlings. Using METOLACHLOR + MESOTRIONE SC under these conditions can result in crop injury.

USE RESTRICTIONS

- Not for sale, use, and distribution in Nassau and Suffolk Counties in the State of New York is prohibited.
- Do not make applications of this product through any type of irrigation system.
- **DO NOT** use flood irrigation to make applications with this product or to incorporate this product.
- **DO NOT** apply this product by air.
- **DO NOT** contaminate water used for domestic purposes or irrigation water used for crops that are not on this label.
- **DO NOT** make applications under conditions that favor runoff or wind erosion to soil that has been treated with this product or drift to non-target areas.
- To prevent movement to off-site areas due to runoff or wind erosion:
- When conditions are favorable for wind erosion, avoid treating powdery dry or light sand soils. Allow the soil surface to settle by rainfall or irrigation first under these types of conditions.
- **DO NOT** make applications to impervious substrates, such as paved or highly compacted surfaces or snow covered/frozen soils.

RESISTANCE MANAGEMENT

Metolachlor	Group	15	Herbicide
Mesotrione	Group	27	Herbicide

METOLACHLOR + MESOTRIONE SC is a combination of two active herbicide ingredients - mesotrione and metolachlor (Group 15 and 27 Herbicides). Two modes of action can be an effective component of a weed resistance management program.

There is potential risk of resistance development in some weeds against the herbicides that have been used repeatedly. While the development of resistance is well understood, it is not easily predicted. Therefore, herbicides must be used in conjunction with resistance management strategies in the area. Consult the local or State agricultural advisors for details. If weed resistance develops in the area, this product used alone may not continue to provide sufficient levels of weed control. If the reduced levels of control cannot be attributed to improper application timing, unfavorable weather conditions or abnormally high weed pressure, a resistant strain may have developed.

To reduce the potential for weed resistance, use this product in a rotation program with other classes of chemistry and modes of action. Always apply this product at the specified labelled rates and in accordance with the use directions. Do not use less than specified label rates alone or in tank mixtures. Do not use reduced rates of the tank mix partner. For optimum performance, scout fields carefully and begin applications when weeds are smaller rather than larger. If resistance is suspected, contact the local or State agricultural advisors.

There are naturally occurring biotypes of broadleaf weeds with known resistance to triazines, ALS, PPO, Glycine (glyphosate) and HPPD herbicides. If weed biotypes that are resistant to triazines, ALS, PPO and glycine inhibitors are in the field, this herbicide should control them if they are listed in the WEEDS CONTROLLED tables in this label.

To reduce the potential of weeds developing resistance to HPPD inhibitors, implement a spray program using multiple modes of action that includes both pre-emergence and post-emergence herbicides that provide effective control of all weeds. Consider weed resistance management strategies that includes multiple modes of action where a minimum of two modes of action are labeled for good control of the target weed when either are applied alone.

Integrated Pest (Weed) Management

Integrate METOLACHLOR + MESOTRIONE SC into an overall weed pest management strategy whenever the use of an herbicide is required. Practices known to reduce weed development (tillage, crop competition) and herbicide use (weed scouting, proper application timing, banding) should be followed wherever possible. Consult local agricultural and weed authorities for additional Integrated Pest (Weed) Management strategies established for your area.

APPLICATION INFORMATION

Ground Application

Space spray nozzles uniformly using the same size and type nozzle to provide accurate and uniform application. To avoid drift and produce good coverage, use nozzles that will produce medium to coarse size droplets. Only use 50-mesh or coarser screens in all inline strainer and nozzle screens. Using agitation, maintain proper product dispersion in the tank, and use a pump that can maintain pressure of at least 35 to 40 PSI at the nozzles. If using extended range or drift reduction nozzles, reduced pressure may be used provided that adequate coverage is maintained. Ensure proper and consistent agitation during spraying through duration until spraying is complete – even when there are brief periods of time where spraying has stopped. Stop and run a full agitation before resuming spray if the spray tank is allowed to sit for more than 5 minutes to re-suspend the solution.

Pre-Emergence Applications

Make pre-emergence applications of METOLACHLOR + MESOTRIONE SC in a spray volume of 10 to 80 gals./A.

Post-Emergence Applications

For optimum weed control, good weed coverage is essential. Make applications in a spray volume of 10 to 30 gals./A.

If weed pressure is high and foliage is dense, use a minimum spray volume of 20 gals/A. For post-emergence applications, use flat fan nozzles for best coverage. Do not use flood jet or venture type nozzles or controlled droplet application. Use only clean water as a carrier.

Aerial Application

Do not apply this product by air.

MANDATORY SPRAY DRIFT MANAGEMENT
GROUND BOOM APPLICATIONS: <ul style="list-style-type: none">• Do not release spray at a height greater than 3 feet above the ground or crop canopy.• Applicators are required to select the nozzles and pressure that deliver medium or coarser droplets (ASABE S572).• DO NOT apply when wind speeds exceed 15 miles per hour at the application site.• DO NOT apply during temperature inversions.
BOOMLESS GROUND APPLICATIONS:

- Applicators are required to select the nozzle and pressure that deliver medium or coarser droplet size (ASABE S572.3) for all applications.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boomless Ground Applications:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Sensitive Areas

Only apply METOLACHLOR + MESOTRIONE SC when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

ADDITIVES/ADJUVANTS

For applications where an adjuvant will be used, it is recommended to select one that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification.

Post Emergence Applications - After Corn Has Emerged

Add either a non-ionic surfactant at 0.25% v/v (1 qt./100 gals.) or crop oil concentrate at a rate of 1% v/v (1 gal./100 gals.) after field corn has emerged. Using a COC will provide better control than using an NIS, but temporary crop injury may occur.

Do not use methylated seed oil (MSO) with this product when applied alone to emerged field corn, or when applied as a post-emergence tank mixture with other products.

Pre-Emergence Applications - Before Corn Emergence

To increase burndown activity on weeds that have emerged, any adjuvant may be used at a pre-emergence or pre-plant timing.

MIXING PROCEDURES

Use either clean water or liquid fertilizers (excluding suspension fertilizers) as carriers for pre-emergence applications.

If using fluid fertilizers, a compatibility test must be conducted. See **COMPATIBILITY TEST** section for additional information. Even if METOLACHLOR + MESOTRIONE SC is determined to be physically compatible with a fluid fertilizer, constant agitation will be necessary to maintain a uniform solution during application. Use only clean water as a carrier.

The spray tank must be thoroughly rinsed, decontaminated and clean before adding either METOLACHLOR + MESOTRIONE SC alone or with tank mix partners. Use only clean water, if water is used as the carrier.

Refer to specific tank mix recommendation sections in this label. Always refer to the tank mix partner label(s) for mixing directions and precautions. Do not exceed maximum label use rates, or combined total maximum seasonal use rates for mesotrione or metolachlor. Do not mix this product with any product bearing a label prohibition against such mixing. If a tank mixture is used, a compatibility test must be conducted. See COMPATIBILITY TEST section below for information on conducting a compatibility test.

COMPATIBILITY TEST

To ensure compatibility of a tank mix partner with METOLACHLOR + MESOTRIONE SC, a compatibility test should be conducted.

Complete liquid fertilizers or nitrogen solutions (excluding suspension fertilizers) may replace all or part of the water in the spray, as recommended in directions for use. Always conduct compatibility test and make actual applications according to label directions and use recommended carrier. Always check compatibility of liquid fertilizers with pesticide(s) before use because, even within the same analysis, liquid fertilizers vary. Tank mixture incompatibility is more common with mixtures of fertilizers and pesticides.

COMPATIBILITY TEST PROCEDURE

(Assuming a 25 gals./A spray volume)

1. Add 1.0 pt. of water or fertilizer carrier to each of two - 1 quart jars with tight lids. It is important to use the same source of water that will be used in the tank mix and to conduct the test at the same temperature the tank
2. mix will be applied as water and temperature can affect compatibility.
3. Add ¼ tsp. or 1.2 mL of a compatibility agent approved for the intended use to one of the jars (¼ tsp equals 2.0 pts./100 gals. of spray). Mix by shaking or gently stirring (if shaking place lid on jar).
4. Add the appropriate amount of pesticide(s) based on described label rates to both jars. If more than one pesticide product will be used, add them separately in the order as described in the Tank Mix Instructions section of this label. Shake or stir gently after each addition to thoroughly mix (if shaking place lid on jar).
5. After all ingredients have been added, place lids on tightly, and invert each jar ten times. Allow the mixtures to stand 15 to 30 minutes. Look for separation, precipitates, gels, heavy oily film on the jar, large flakes, or other signs of incompatibility. Compare the two jars to determine if the compatibility agent is needed. If mixtures separate, but can be easily and readily remixed, the mixture can be sprayed but good agitation must be used. If it is determined the mixtures are incompatible, use the following methods to test for improving compatibility:
 - a) Make a slurry of the dry pesticide(s) in water before addition, or
 - b) Add ½ of the compatibility agent to the carrier (fertilizer or water) and the other ½ to the emulsifiable concentrate (EC) or flowable pesticide before adding to the mixture. If mixture is still not compatible, do not use the mixture.
6. Dispose of any pesticide wastes in accordance with the Storage and Disposal section in this label.

TANK MIXTURES

Tank Mix Instructions

Use sprayers and equipment that are in good, clean condition and maintain adequate agitation. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

If the tank mix partner is determined to be compatible, fill the tank half full of the carrier. Begin agitation and maintain throughout mixing and application. Make sure all return lines to the spray tank discharge below the liquid level. Prepare the tank mixture components and add to the tank in the following order:

1. If using ammonium sulfate (AMS) – add and continue until it is completely dispersed.
2. If using a wettable powder or dry flowable formulation, make a slurry with water first and then add it slowly through the screen into the tank. Maintain agitation during this step.
3. If using a flowable formulation, add slowly through screen into the tank. Diluting the flowable with water before adding to the tank may improve mixing and compatibility with dry flowable formulations.
4. Add METOLACHLOR + MESOTRIONE SC.
5. Add any other tank mix products, adding emulsifiable concentrates last.
6. If an adjuvant will be used, add as the final step. Maintain agitation.
7. Complete filling the spray tank with the carrier and maintain agitation. Make application as soon as possible after spray mixture is prepared. Do not leave mixture in spray tank overnight unattended or without agitation.

If METOLACHLOR + MESOTRIONE SC is added to the spray tank via induction, compatibility of the spray mixture may be compromised. If using an induction tank (or comparable equipment), add each tank mixture product separately and allow each to fully disperse into the spray tank before adding the next product. For optimum compatibility, rinse the induction tank with clean water before adding each component.

The addition of METOLACHLOR + MESOTRIONE SC to the spray tank via in-line injection is not

recommended.

Cleaning Equipment Post Application

Careful attention must be use when cleaning equipment before spraying a crop other than field corn following applications with this product. Mix the volume of spray solution based on the area of application and mix only as much spray solution as needed.

Tank and Sprayer Clean Out

1. Use clean water to flush the tank, hoses, boom, and nozzles.
2. Add 1 gal. of household ammonia per 25 gals. of water. Or alternatively, use a commercially available spray tank cleaner.
3. Using pressure washer, clean the inside of the spray tank with this solution. Wash all parts of the tank, including the inside and top surface. If there is not a pressure washer available, fill the sprayer completely with the cleaning solution to provide contact with all internal surfaces of the tank and plumbing. Begin agitation in the sprayer and thoroughly recirculate the solution in the tank for at least 15 minutes. Remove all visible deposits from the spray equipment.
4. Use the cleaning solution to flush the hoses, spray lines, and nozzles for at least 1 minute.
5. Flush dead space areas with water by removing boom end caps, and then replace caps.
6. Dispose of rinsate from the clean-out according to all local State and federal regulations.
7. Repeat the steps 2 to 5 above.
8. After completing the above procedures, remove and clean the nozzles, screens, and strainers separately in the cleaning solution.
9. Completely rinse the spray tank and equipment with clean water.

WEEDS CONTROLLED

Make applications of METOLACHLOR + MESOTRIONE SC as directed in this label to control or partially control the weeds listed in the tables below. Tank mixtures may control additional weeds. See the Tank Mixtures sections for specific and additional information. Always refer to the tank mix partner label(s) for specific use rates, directions and restrictions.

Weed control may be reduced, if a sufficient rainfall is not received within 7 days after application. Apply 0.5 to 1 inch of water, if irrigation is available. Conduct a uniform, shallow cultivation as soon as weeds emerge, if irrigation is not available. Post-emergence control may be reduced or delayed when weeds are stressed or not actively growing due to drought, heat, lack of fertility, flooding, or prolonged cool temperatures.

Pre-Emergence Applications: Weeds Controlled or Suppressed

Common Name	Scientific Name	C = Control S=Suppression
Amaranth, Palmer	<i>Amaranthus palmeri</i>	C
Amaranth, Powell	<i>Amaranthus powellii</i>	C
Barnyardgrass	<i>Echinochloa crus-galli</i>	C
Buffalobur	<i>Solanum rostratum</i>	C
Carpetweed	<i>Mollugo verticillata</i>	C
Cocklebur, common	<i>Xanthium strumarium</i>	S
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	C
Cupgrass, prairie	<i>Eriochloa contracta</i>	C
Cupgrass, Southwestern	<i>Eriochloa acuminata</i>	C
Cupgrass, woolly	<i>Eriochloa villosa</i>	S
Foxtail, giant	<i>Setaria faberi</i>	C
Foxtail, green	<i>Setaria viridis</i>	C
Foxtail, robust (purple, white)	<i>Setaria</i> spp.	C
Foxtail, yellow	<i>Setaria pumila</i>	C
Galinsoga	<i>Galinsoga parviflora</i>	C

Common Name	Scientific Name	C = Control S=Suppression
Goosegrass	<i>Eleusine indica</i>	C
Jimsonweed	<i>Datura stramonium</i>	C
Johnsongrass, seedling	<i>Sorghum halepense</i>	S
Kochia	<i>Kochia scoparia</i>	S
Lambsquarters, common	<i>Chenopodium album</i>	C
Millet, foxtail	<i>Setaria italica</i>	C
Millet, wild proso	<i>Panicum miliaceum</i>	S
Morningglory, entireleaf	<i>Ipomoea hederacea</i>	S
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>	S
Nightshade, black	<i>Solanum nigrum</i>	C
Nightshade, Eastern black	<i>Solanum ptycanthum</i>	C
Nightshade, hairy	<i>Solanum sarrachoides</i>	C
Nutsedge, yellow	<i>Cyperus esculentus</i>	C
Panicum, browntop	<i>Panicum fasciculatum</i>	C
Panicum, fall	<i>Panicum dichotomiflorum</i>	C
Panicum, Texas	<i>Panicum texanum</i>	S
Pigweed, redroot	<i>Amaranthus retroflexus</i>	C
Pigweed, smooth	<i>Amaranthus hybridus</i>	C
Purslane, common	<i>Portulaca oleracea</i>	C
Pusley, Florida	<i>Richardia scabra</i>	C
Ragweed, common	<i>Ambrosia artemisiifolia</i>	S
Ragweed, giant	<i>Ambrosia trifida</i>	S
Rice, red	<i>Oryza sativa</i>	C
Sandbur, field	<i>Cenchrus incertus</i>	S
Shattercane	<i>Sorghum bicolor</i>	S
Sida, prickly	<i>Sida spinosa</i>	S
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>	S
Smartweed, ladythumb	<i>Polygonum persicaria</i>	C
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>	C
Sprangletop, red	<i>Leptochloa filiformis</i>	C
Velvetleaf	<i>Abutilon theophrasti</i>	C
Waterhemp, common	<i>Amaranthus rudis</i>	C
Waterhemp, tall	<i>Amaranthus tuberculatus</i>	C
Witchgrass	<i>Panicum capillare</i>	C

Early Post-Emergence Applications: Weeds Controlled or Suppressed

Applied early post-emergence, METOLACHLOR + MESOTRIONE SC will provide control or suppression of small emerged broadleaf weeds that are less than 3 inches tall but will not provide good control of weeds resistant to post-emergence HPPD inhibitors.

Common Name	Scientific Name	C = Control S=Suppression
Amaranth, Palmer	<i>Amaranthus palmeri</i>	C
Amaranth, Powell	<i>Amaranthus powellii</i>	C
Buffalobur	<i>Solanum rostratum</i>	C
Carpetweed	<i>Mollugo verticillata</i>	C
Cocklebur, common	<i>Xanthium strumarium</i>	C
Dandelion	<i>Taraxacum officinale</i>	S
Galinsoga	<i>Galinsoga parviflora</i>	C
Hemp	<i>Cannabis sativa</i>	C
Horsenettle	<i>Solanum carolinense</i>	C
Horseweed (marestail)	<i>Conyza canadensis</i>	C
Jimsonweed	<i>Datura stramonium</i>	C

Common Name	Scientific Name	C = Control S=Suppression
Kochia	<i>Kochia scoparia</i>	S
Lambsquarters, common	<i>Chenopodium album</i>	C
Morningglory, entireleaf	<i>Ipomoea hederacea</i>	S
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>	S
Mustard, wild	<i>Brassica kaber</i>	C
Nightshade, black	<i>Solanum nigrum</i>	C
Nightshade, Eastern black	<i>Solanum ptycanthum</i>	C
Nightshade, hairy	<i>Solanum sarrachoides</i>	C
Nutsedge, yellow	<i>Cyperus esculentus</i>	S
Pigweed, redroot	<i>Amaranthus retroflexus</i>	C
Pigweed, smooth	<i>Amaranthus hybridus</i>	C
Pokeweed	<i>Phytolacca americana</i>	C
Potatoes, volunteer	<i>Solanum spp.</i>	C
Purslane, common	<i>Portulaca oleracea</i>	S
Pusley, Florida	<i>Richardia scabra</i>	C
Ragweed, common	<i>Ambrosia artemisiifolia</i>	C
Ragweed, giant	<i>Ambrosia trifida</i>	C
Sida, prickly	<i>Sida spinosa</i>	S
Smartweed, ladysthumb	<i>Polygonum persicaria</i>	C
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>	C
Thistle, Canada	<i>Cirsium arvense</i>	S
Velvetleaf	<i>Abutilon theophrasti</i>	C
Waterhemp, common	<i>Amaranthus rudis</i>	C
Waterhemp, tall	<i>Amaranthus tuberculatus</i>	C

ROTATIONAL CROPS

The crop rotational intervals listed below should be observed following application of METOLACHLOR + MESOTRIONE SC. For tank mixtures of other products with METOLACHLOR + MESOTRIONE SC, follow the most restrictive product's crop rotation interval listed on the tank mix partner label.

Crop Rotational Intervals

Crop	Crop Rotational Interval*
Corn (all types) and grain sorghum**	Anytime
Cereals (barley, oats, rye, wheat)	4.5 Months
Cotton, peanuts, potatoes, and soybeans	Spring following the application
Beans (dry and snap), cucurbits, peas, red clover, sugar beets, tomatoes.	18 Months

*Period between METOLACHLOR + MESOTRIONE SC application and planting of the rotational crop.

**Seed for grain sorghum must be treated with a product safener to provide tolerance to metolachlor.

CORN - Directions for Use

Make a pre-emergence application of METOLACHLOR + MESOTRIONE SC for control of annual grass and broadleaf weeds in field corn, seed corn, sweet corn, and yellow popcorn. Make an early post-emergence application of METOLACHLOR + MESOTRIONE SC for the control of broadleaf weeds in field corn and seed corn. See the **WEEDS CONTROLLED** section of this label for a list of weeds controlled or suppressed. Do not make applications of METOLACHLOR + MESOTRIONE SC to yellow popcorn or sweet corn after the crop has emerged, or crop injury may result.

APPLICATION TIMINGS

Reduced Tillage - Burndown Applications

In reduced or no-till corn and prior to crop emergence, METOLACHLOR + MESOTRIONE SC may be applied alone or in tank mixtures with paraquat or glyphosate for the burndown of weeds that have emerged.

See the **WEEDS CONTROLLED** section of this label for a list of weeds controlled or suppressed.

Refer to the paraquat or glyphosate product label for additional information on weeds controlled, directions for use, restrictions and precautions.

See the **ADDITIVES** and **TANK MIX** sections on this product label for additional information.

Early Pre-Plant and Pre-Emergence Applications

Make an early pre-plant application of METOLACHLOR + MESOTRIONE SC up to 14 days before planting or pre-emergence application in field corn, seed corn, sweet corn and yellow popcorn.

Post-Emergence Applications

Make a post-emergence application of METOLACHLOR + MESOTRIONE SC to field or seed corn after emergence up to the time when the plants reach 30 inches in height or up to the 8-leaf stage of corn growth. Use only clean water as the carrier. Do not make post-emergence applications in liquid fertilizer or severe crop injury will result.

See the ADDITIVES section on this label for recommendations on adjuvants for burndown applications.

RESTRICTIONS:

Do not make applications of METOLACHLOR + MESOTRIONE SC to emerged yellow popcorn or sweet corn, or severe crop injury may result.

METOLACHLOR + MESOTRIONE SC Use Rates

Make application of METOLACHLOR + MESOTRIONE SC at 2.0 - 2.4 qts./A (1.68 lbs. a.i./A metolachlor and 0.165 lb. a.i./A mesotrione to 2.016 lbs. a.i./A metolachlor and 0.198 lb. a.i./A mesotrione) for control or suppression of the weeds listed in the WEEDS CONTROLLED section of this label. Before making applications of this product, determine the soil organic matter content of the field.

For soils with <3% organic matter content – use 2.0 quarts of METOLACHLOR + MESOTRIONE SC per acre. For soils with \geq 3% organic matter content – use 2.4 quarts of METOLACHLOR + MESOTRIONE SC per acre.

Use of this product on soils with >10% soil organic matter is not recommended and may result in poor weed control.

TANK MIXTURES

Pre-Emergence Applications (before crop has emerged)

Tank mix partners listed in the table below may be used in conventional, reduced, or no-till operations and by the same application methods and the same timings as METOLACHLOR + MESOTRIONE SC unless otherwise directed in the tank mix partner product label.

Follow all tank mix product labels for use rates, precautions and restrictions.

Pre-Emergence Tank Mix Applications METOLACHLOR + MESOTRIONE SC in Field Corn

Tank Mixture Recommendations* for use with products containing:	Target Use
atrazine	Broadleaf and grass weed control improved
glyphosate	Burndown of emerged existing weeds
metribuzine	Broadleaf weed control improved
paraquat	Burndown of emerged existing weeds
simazine	Broadleaf and grass weed control improved
2,4-D	Burndown of emerged existing weeds
lambda-cyhalothrin	Insect control (see product label)
* Reference the tank mix partner(s) product label for directions for use, precautions, and restrictions. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.	

Early Post-Emergence Applications (after crop has emerged)

Tank mix products listed in the table below may be used in conventional, reduced, or no-till systems and applied by the same methods and at the same timings as METOLACHLOR + MESOTRIONE SC unless otherwise specified in the tank mix product label. Follow all tank mix product labels for use rates and restrictions.

Post-Emergence Tank Mixture Applications METOLACHLOR + MESOTRIONE SC in Field Corn

Tank Mixture Recommendations^{1,2} for use with products containing:	Target Use Target Use
atrazine	Broadleaf and grass weed control improved
nicosulfuron	Emerged grass weed control
rimsulfuron + thifensulfuron-methyl	Emerged grass weed control
glufosinate-ammonium	Refer to instructions under the Metolachlor 36.8% + Mesotrione 3.68% Spray Programs in LibertyLink® Corn section of this label.
primisulfuron-methyl + dicamba	Broadleaf and grass weed control improved
prosulfuron	Broadleaf and grass weed control improved
glyphosate	Refer to the instructions under the Metolachlor + Mesotrione Spray Programs in glyphosate-tolerant Corn section of this label.
primisulfuron-methyl + prosulfuron	Broadleaf and grass weed control improved
dicamba + diflufenzopyr	Emerged grass weed control
nicosulfuron + rimsulfuron	Emerged grass weed control
lambda-cyhalothrin	Insect control (see product label)

¹ Reference the tank mix partner(s)' product label for directions for use, precautions, and restrictions.

² Refer to the Additives section of this label for recommendations when making applications of METOLACHLOR + MESOTRIONE SC alone or in tank mixture to field corn that has emerged.

METOLACHLOR + MESOTRIONE SC Spray Programs in Glyphosate-Tolerant Corn

Make early post-emergence tank mixture applications of METOLACHLOR + MESOTRIONE SC with a solo glyphosate product (example: Touchdown® or Roundup® brands) that is registered for use over-the-top in glyphosate tolerant field corn (example: Roundup® Ready or Agrisure® GT Corn) at rates as low as 1.6 qts./A.

To reduce weed competition with the crop, application of this mixture should be targeted to weeds that are 1 to 2 inches. If the glyphosate product has an adjuvant included in the formulation (the product label does not call for an adjuvant being added), only spray-grade ammonium sulfate (AMS) at 8.5

lbs./100 gals. should be added to the tank mixture. If the glyphosate product label recommends an adjuvant in addition to AMS, add a non-ionic surfactant (NIS) at 0.25% v/v and AMS to this spray tank mixture. Do not use urea ammonium nitrate (UAN), crop oil concentrate (COC), or methylated seed oil (MSO) type adjuvants in these tank mixtures, or crop injury may result. Read and follow all directions for use, precautions and restrictions on the tank mix partner glyphosate label.

As an alternative, a pre-emergence application of METOLACHLOR + MESOTRIONE SC may be made at rates as low as 1.6 qts./A as part of a two-pass weed control program when followed by a post-emergence application of a glyphosate-containing product in glyphosate-tolerant corn (example: Roundup® Ready or Agrisure® GT Corn). When this type of application is made, METOLACHLOR + MESOTRIONE SC will provide reduced competition of the weeds listed in the Pre-Emergence Applications: Weeds Controlled or Suppressed table for a period of 30+ days, improving the flexibility in application timing and effectiveness of the glyphosate-based product application. Follow all directions for use, precautions and restrictions on the glyphosate product label.

A pre-emergence application of METOLACHLOR + MESOTRIONE SC may be made at 1.0 to 1.2 qts./A as part of a two-pass weed control program when followed by a tank mix of METOLACHLOR + MESOTRIONE SC and glyphosate in glyphosate-tolerant corn (example: Roundup® Ready or Agrisure GT Corn). Make application of METOLACHLOR + MESOTRIONE SC at 1.0 qt./A for soils with less than 3% organic matter, and 1.2 qts./A for soils with greater than 3% organic matter. Follow all directions for use, precautions and restrictions on each product label.

METOLACHLOR + MESOTRIONE SC Spray Programs in Glufosinate-Tolerant Corn

A post-emergence application of METOLACHLOR + MESOTRIONE SC may be made at 1.6 qts./A in tank mixture with Ignite® and applied over-the-top in field corn designated as LibertyLink®. To reduce weed competition with the crop, application of this mixture should be targeted to weeds that are 1 to 2 inches. Ammonium sulfate (AMS) may be added as an adjuvant as directed on the Ignite® label. However, AMS should be the only adjuvant used in this tank mixture. Do not make tank mixture applications with urea ammonium nitrate (UAN), crop oil concentrate (COC), non-ionic surfactants (NIS), or methylated seed oil (MSO) type adjuvants in these type of spray programs, or crop injury may result. Follow all directions for use, precautions and restrictions on the Ignite® product label.

As an alternative, a pre-emergence application of METOLACHLOR + MESOTRIONE SC may be made at 1.6 qts./A as part of a two-pass weed control program when followed by a post-emergence application of Ignite® in field corn designated as LibertyLink®. When this type of application is made, METOLACHLOR + MESOTRIONE SC will provide reduced competition of the weeds listed in the **Pre-Emergence Applications: Weeds Controlled or Suppressed** table for a period of 30+ days, improving the flexibility in application timing and effectiveness of the Ignite® product application. Follow all directions for use, precautions and restrictions on the Ignite® product label.

RESTRICTIONS FOR ALL CORN USES

- **DO NOT** make applications of more than 2.4 qts. (2.016 lbs. a.i./A metolachlor and 0.198 lb. a.i./A mesotrione) of METOLACHLOR + MESOTRIONE SC per year.
- **DO NOT** make applications of METOLACHLOR + MESOTRIONE SC to corn that is taller than 30 inches in height or corn that is larger than the 8-leaf stage of growth.
- **DO NOT** graze or feed forage from treated areas for 45 days following last application.
- **DO NOT** harvest corn for grain, forage, or stover within 45 days after a post-emergence application of METOLACHLOR + MESOTRIONE SC.
- **DO NOT** apply METOLACHLOR + MESOTRIONE SC as a post-emergence application in a tank mix with any organophosphate or carbamate insecticide, or severe corn injury may result.

PRECAUTIONS FOR ALL CORN USES

- Severe adverse crop response and corn injury can result if applying METOLACHLOR + MESOTRIONE SC post-emergence to corn that has emerged and that has received an at-plant application of Counter® insecticide. Environmental conditions that promote poor growth will increase the likelihood and risk of severe crop injury.

- Severe corn injury can occur when an organophosphate or carbamate insecticide post-emergence application is made to corn within 7 days before or 7 days after a METOLACHLOR + MESOTRIONE SC application. Environmental conditions that promote poor growth will increase the likelihood and risk of severe crop injury.

METOLACHLOR + MESOTRIONE SC Applications – Grain Sorghum

Make a non-incorporated, pre-plant application of METOLACHLOR + MESOTRIONE SC in sorghum that has been seed-treated with Concep® III (or equivalent safener that provides tolerance to metolachlor) up to 21 days before planting and up through pre-emergence for weed control. See the **Pre-Emergence Applications: Weeds Controlled or Suppressed** table for a listing of weeds.

Make a broadcast, non-incorporated spray application at 2.0 qts./A starting at 21 days pre-plant and up through planting, but before sorghum has emerged. Making application less than 7 days before the sorghum planting can increase the risk of crop injury, particularly if there is rainfall or irrigation after the application. Symptoms of crop injury include temporary bleaching of young sorghum leaves, or in severe conditions, stunting or partial stand loss. Making the application of METOLACHLOR + MESOTRIONE SC at greater than 7 days (and no more than 21 days) before the sorghum planting will reduce the risk of adverse crop response.

When METOLACHLOR + MESOTRIONE SC application is made before planting, do not incorporate and minimize soil disturbance of the treatment area during planting to minimize the potential for reduced weed control.

Split applications of METOLACHLOR + MESOTRIONE SC may be made to sorghum as an early pre-plant (7 to 21 day prior to planting), non-incorporated application at 1.0 to 1.25 qts./A of METOLACHLOR + MESOTRIONE SC following with a second application of METOLACHLOR + MESOTRIONE SC made at 0.75 to 1.0 qt./A before the sorghum has emerged. Do not exceed 2.0 qts./A of product for the split applications.

It is recommended to use a nonionic surfactant (NIS) type adjuvant at 0.25% v/v or a crop oil concentrate (COC) at 1% v/v in the spray solution if weeds are present at the time of application. A spray grade UAN at 2.5% v/v or AMS at 8.5 lbs./100 gallons of spray may also be added in addition to the COC or NIS to the mixture to improve control of weeds that have already emerged. The addition of additives is not recommended, if weeds have not emerged at the time of application.

SORGHUM USE RESTRICTIONS:

- **DO NOT** make applications of more than 2.0 quarts (1.68 lbs. a.i./A metolachlor and 0.165 lb. a.i./A mesotrione) of METOLACHLOR + MESOTRIONE SC per year.
- **DO NOT** make applications of METOLACHLOR + MESOTRIONE SC to sorghum that is grown on sandy soils (sand, sandy loam, or loamy sand).
- **DO NOT** make applications of METOLACHLOR + MESOTRIONE SC to grain sorghum that has emerged or severe crop injury will result.
- **DO NOT** make applications of METOLACHLOR + MESOTRIONE SC to sorghum grown for forage, sweet sorghum (sorgo), sudangrass, sorghum-sudangrass hybrids, or dual-purpose sorghum.
- Seeds must be treated with Concep® III herbicide or an alternate seed safener that provides tolerance to metolachlor before planting, or severe adverse crop response and injury may result.
- **DO NOT** apply METOLACHLOR + MESOTRIONE SC to sorghum that is grown south of Interstate 20 (I-20) or east of Highway 277 in the state of Texas.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep container tightly closed when not in use. Do not store near seeds,

fertilizers, or foodstuffs. Can be stored at temperatures as low as -10°F. Keep away from heat and flame.

PESTICIDE DISPOSAL: Open dumping is prohibited. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Rinse spray equipment. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of as described above, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING [5 gallons or less]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER HANDLING [greater than 5 gallon]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

DO NOT USE CONTAINERS FOR THE STORAGE OF FOOD, FEED, OR DRINKING WATER!

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of this product, which are beyond the control of ALBAUGH, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ALBAUGH, LLC and Seller harmless for any claims relating to such factors.

ALBAUGH, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ALBAUGH, LLC and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALBAUGH, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, in no event shall ALBAUGH, LLC or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ALBAUGH, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ALBAUGH, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

ALBAUGH, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of ALBAUGH, LLC.

All trademarks are the property of their respective owners.

LABEL HISTORY

(not to be included in the final printed label)

FILE NAME	VERSION MARK	COMMENT
042750-00346.20200206.DRAFT Clean	020620	Changes requested by US EPA
042750-00345.20200207.MASTER	020720	Section 3 Approval
042750-00345.20200211.ABN Clean	021120	Notification
042750-00345.20200324.ABN MASTER	042920	Notification
042750-00346.20210825.DRAFT	082521	EPA Reg Review plus NMFS BioP
042750-00346.20221212.DRAFT	12122022	EPA Reg Review