



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Pesticide Programs  
 Registration Division (7505P)  
 1200 Pennsylvania Ave., N.W.  
 Washington, D.C. 20460

**EPA Reg. Number:**  
 42750-351

**Date of Issuance:**  
 9/9/19

**NOTICE OF PESTICIDE:**

Registration  
 Reregistration  
 (under FIFRA, as amended)

**Term of Issuance:**  
 Conditional

**Name of Pesticide Product:**  
 ST Pre-Mix #17

**Name and Address of Registrant (include ZIP Code):**

Morris Gaskins, Director Product Registrations  
 Albaugh, LLC  
 P.O. Box 2127  
 Valdosta, GA 31604-2127

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

**Signature of Approving Official:**

Venus Eagle, Product Manager 01  
 Invertebrate and Vertebrate Branch 3, Registration Division (7505P)  
 Registration Division (7505P)

**Date:**

9/9/19

2. You are required to comply with the data requirements described in the DCI identified below:
  - a. Difenaconazole GDCI-128847-1602
  - b. Thiabendazole GDCI-060101-1453

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.
4. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, “EPA Reg. No. 42750-351.”
5. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 12/19/18

If you have any questions, please contact Marianne Lewis by phone at (703) 308-8043 or via email at [lewis.marianne@epa.gov](mailto:lewis.marianne@epa.gov).

Enclosure

IMIDACLOPRID	GROUP	4A	INSECTICIDE
METALAXYL	GROUP	4	FUNGICIDE
DIFENACONAZOLE	GROUP	3	FUNGICIDE
TEBUCONAZOLE	GROUP	3	FUNGICIDE
THIABENDAZOLE	GROUP	1	FUNGICIDE

## ST PRE-MIX #17

FOR USE ONLY IN COMMERCIAL SEED TREATMENT FACILITIES ON BARLEY, OATS, TRITICALE, AND WHEAT FOR PROTECTION OF SEEDS AGAINST LISTED PESTS and SOILBORNE DISEASES


### ACTIVE INGREDIENTS:

Imidacloprid .....	19.38%
Difenoconazole .....	4.22%
Metalaxyl .....	3.17%
Tebuconazole .....	0.70%
Thiabendazole .....	2.25%

<b>OTHER INGREDIENTS:</b> .....	<u>70.28%</u>
<b>TOTAL:</b> .....	100.00%

### Contains:

- 1.76 pound Imidacloprid per gallon (211.1 grams per liter)
- 0.38 pound Difenoconazole per gallon (45.6 grams per liter)
- 0.29 pound Metalaxyl per gallon (34.8 grams per liter)
- 0.064 pound Tebuconazole per gallon (7.7 grams per liter)
- 0.20 pound Thiabendazole per gallon (24.0 grams per liter)



**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For transportation or medical emergencies call CHEMTREC toll free at 1-800-424-9300.	

EPA Reg. No. 42750-GLR/351

EPA Est. No. XXX-XX-XXXX

NET CONTENTS: \_\_\_\_\_ Gallons

Batch No. \_\_\_\_\_

### MANUFACTURED FOR:

Albaugh, LLC  
Ankeny, IA 50021

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION.** Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist or vapor.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks,
- Chemical-resistant gloves made out of barrier laminate, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, or viton  $\geq 14$  mils.
- Protective eyewear

**Important!** Persons involved in bagging treated seed, sewing or moving bags of treated seed, or cleaning up bagging areas or seed treatment equipment are pesticide handlers and must wear the PPE required on this label for pesticide handlers.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

Users should:

Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### PHYSICAL AND CHEMICAL HAZARDS

DO NOT place product near or allow product to come into contact with strong oxidizing substances (such as potassium permanganate).

### ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to mammals, birds, fish, and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate. Cover or incorporate spilled treated seeds.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

**Pollinator Protection Statement:** Imidacloprid is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

**Groundwater Advisory Statement:** Metalaxyl and Tebuconazole are known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Surface Water Advisory:** This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements, specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## **USE RESTRICTIONS**

- Only for use in commercial seed treatment facilities. Not for use on agricultural establishments in hopper box, planter box, slurry box or other seed treatment applications at or immediately before planting.
- Not for use on Oats in California.
- Do not use treated seed for feed, food, or oil purposes.
- Store treated seed away from feeds and foodstuffs.
- Do not allow children, pets, or livestock to have access to treated seed.
- Regardless of the method of application, do not apply more than 0.5 lbs active ingredient imidacloprid per acre per year.
- As per 40 CFR 153.155, all seed treated with ST PRE-MIX #17 must be colored with an EPA approved dye or colorant to visually distinguish seed that is treated with a pesticide. It is mandatory that such dye or colorant be added into the slurry mixture containing ST PRE-MIX #17, prior to application. This regulation is necessary to prevent treated seed from accidental use as a food for man or feed for animals.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Exposed treated seed is highly toxic to birds. Cover or incorporate spilled treated seed. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.
- Do not graze or feed livestock on green barley forage until 31 days after planting.
- Do not graze or feed livestock on all other areas until 55 days after planting.
- Do not plant any crop other than those on this label within 30 days to fields in which treated seeds were planted.
- Do not store treated seed in burlap bags or impervious bags/container or in areas that are poorly ventilated
- Do not store leftover treated seed where there is a chance of it becoming mixed with untreated seed.
- Store treated seed in cool, dry conditions.
- Do not use empty seed bags for any other purpose.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that

- are used in agronomic practice.
- Dispose of seed packaging in accordance with local requirements

### **RESISTANCE MANAGEMENT RECOMMENDATIONS**

For resistance management ST Pre-Mix #17 contains imidacloprid, a Group 4a insecticide. Any insect population may contain individuals naturally resistant to ST Pre-Mix #17 and other Group 4A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of ST Pre-Mix #17 or other Group 4A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
  - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
  - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
  - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Albaugh, LLC at 1-800-247-8013.

For resistance management ST Pre-Mix #17 contains difenoconazole and tebuconazole, Group 3 fungicides; thiabendazole, a Group 1 fungicide, and; metalaxyl, a Group 4 fungicide. Difenoconazole and tebuconazole belong to the DMI class of chemistry. Thiabendazole belongs to the methyl benzimidazole carbamates class of chemistry. Metalaxyl belongs to the phenylamide class of chemistry.

Fungal populations may contain individuals naturally resistant to active ingredients in ST Pre-Mix #17 and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies such as:

- Alternation with fungicides of a different mode of action and/or tank mixes established for the crop and use area.
- Use based on an IPM program that includes

- field sanitation,
- scouting,
- historical information related to pesticide use, and crop rotation.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- The IPM program should also consider cultural, biological, and other chemical control practices.

Albaugh encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label. For additional information on Fungicide Resistance Management contact an Albaugh, LLC representatives at 1-800-247-8013, your local extension specialist or certified crop advisor. Visit the Fungicide Resistance Action Committee (FRAC) at <http://www.frac.info> for additional information on fungicide resistance management

### PRODUCT INFORMATION

This product is a mixture of the fungicides Difenoconazole, Tebuconazole, Metalaxyl, Thiabendazole and the insecticide Imidacloprid. When used on listed crops, this product aids in the control or suppression of the following seed, soilborne diseases and insects on seed and young seedlings of Barley, Oats, Triticale, and Wheat.

Apply as a seed treatment using standard slurry or mist-type commercial seed treatment equipment. Apply a uniform application to seed to ensure seed safety and best disease and insect protection. Seed must be sound and well cured prior to treatment. Product must be diluted with sufficient water to ensure complete seed coverage.

**Tank Mixing:** It is the pesticide user’s responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Compatibility:** All tank mixes should be pre-tested to determine physical compatibility between formulations. Before mixing ST PRE-MIX #17 with other seed treatment products conduct a compatibility test before use:

- Use a clear glass container, thoroughly mix all intended seed treatment products with appropriate amount of water;
- Allow the mixture to sit for one hour;
- Remix and observe for incompatibility

### SEED TREATMENT

CROP	DISEASE CONTROL	RATE
Barley	Control of seed rots such as <i>Penicillium</i> and <i>Aspergillus</i> and seedborne <i>Fusarium</i> Scab and <i>Septoria</i> .	Fl. Oz. / 100 LB Seed
	Control of Dwarf bunt, Stinking smut (Common bunt), Flag smut, Loose smut, Covered smut of barley.	4.0* <sup>3</sup>
	Control of early season Barley stripe, <i>Pythium</i> damping-off, <i>Septoria</i> disease complex, <i>Rhizoctonia</i> root rot, Common root rot ( <i>Cochliobolus</i> spp.), <i>Fusarium</i> foot rot.	

CROP	DISEASE CONTROL	RATE
Oats* <sup>1</sup>	Early season suppression of powdery mildew, and wheat leaf rust.	
Triticale	<b>INSECT CONTROL</b>	
Wheat	Control of light to moderate pressure of wireworms. Controls aphids (including Bird cherry-oat, English grain, Greenbug and Russian wheat), and Hessian fly	
<p><b>APPLICATION DIRECTIONS:</b> Apply this product using standard slurry or mist-type commercial seed treatment equipment in a manner that insures uniform coverage of the seed. Depending upon type of equipment used, uniform coverage of the seed can be attained using this product alone or combined with a diluent such as water to a total slurry rate of 15 ounces / 100 lbs of seed. Mix the slurry thoroughly with the seed to insure uniform coverage.</p> <p><b>RESTRICTIONS:</b> Regardless of application method, do not apply more than 0.5 lbs / A / calendar year imidacloprid active ingredient.</p> <p>*1 - Not approved for use in California *2 - 4.0 fl. oz. of ST PRE-MIX #17 contains 0.055 lbs (25.0 grams) of imidacloprid.</p>		

### SEED BAG LABELING REQUIREMENTS

Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act. The user of this product is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

#### THE FEDERAL SEED ACT REQUIRES THAT BAGS CONTAINING TREATED SEEDS BE LABELED WITH THE FOLLOWING STATEMENTS:

- This seed has been treated with ST PRE-MIX #17 (difenoconazole, metalaxyl, tebuconazole, imidacloprid, and thiabendazole).
- **DO NOT** use for feed, food, or oil purpose.

#### THE US ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON BAGS CONTAINING SEEDS TREATED WITH ST PRE-MIX #17 (difenoconazole, metalaxyl, tebuconazole, thiabendazole and imidacloprid):

- **Pollinator Precautions:** Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- **Groundwater Advisory:** Metalaxyl and Tebuconazole are known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- **Surface Water Advisory:** This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.



- Store treated seeds away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Exposed treated seed is highly toxic to birds. Cover or incorporate spilled treated seeds.
- Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** graze or feed livestock on green barley forage until 31 days after planting.
- **DO NOT** graze or feed livestock on green wheat, oats, or triticale for 55 days after planting.
- **DO NOT** contaminate bodies of water when disposing of planting equipment wash water.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- **DO NOT** plant any crop other than those on this label within 30 days to fields in which treated seeds were planted.
- Regardless of the type of application (seed, foliar, or soil application) do not apply more than 0.5 lbs imidacloprid per acre per year.
- This seed has been treated with \_X\_ lbs imidacloprid, \_X\_ lbs difenoconazole, \_X\_ lbs metalaxyl, \_X\_ lbs tebuconazole, \_X\_ lbs thiabendazole.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.
- Do not use empty seed bags for any other purpose.
- Do not store leftover treated seed where there is a chance of it becoming mixed with untreated seed.
- Do not store treated seed in burlap bags or impervious bags/container or in areas that are poorly ventilated
- Store treated seed in cool, dry conditions.
- Not for use on Oats in California.

## STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool place, DO NOT store in direct sunlight. Protect from freezing temperatures.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING:

[Nonrefillable Containers 5 Gallons or Less]

Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration.

[Nonrefillable containers larger than 5 gallons]

Nonrefillable container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Refillable containers larger than 5 gallons]

Refillable container. Refill this container with pesticide only. DO NOT reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. For final disposal, offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

### **WARRANTY DISCLAIMER**

The Directions For Use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

### **LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.