

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: MAY BE HARMFUL IF SWALLOWED Do not take internally. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Wash contaminated clothing before reuse. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine administered by injection only is an antidote.

ENVIRONMENTAL HAZARD

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Use this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Keep away from food, feed-stuffs, and domestic water supplies. Keep container closed and out of reach of children and pets.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

TURF AND LAWNS

Apply Red Panther Dursban Granules insecticide uniformly to home lawns and other ornamental and recreational turf grass areas using any hand or power operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear using Red Panther Dursban Granules at the rate of 200 pounds per acre. To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply the balance at a right angle. Immediately after application sprinkle the turf with 1300 gallons of water per acre so as to wash the granules into the zone of insect infestation. Consult the following table to determine amounts of Red Panther Dursban Granules and water needed for smaller sized areas. Re-treat as necessary.


RATES TO USE

Area to be Treated	Pounds of Dursban Granules	Gallons of water to Sprinkle on Turf after Treatment
1000 sq. ft.	5	30
5000 sq. ft.	25	150
10700 sq. ft.	50	325

To control lawn white grubs (larvae of European chafer and Japanese beetle) use two to four times the amounts of Red Panther Dursban Granules shown in the table, i.e. 10 to 20 pounds per 1,000 sq. ft. Apply the granules uniformly over the area treated followed immediately by thorough watering to wash the insecticide into the turf. For best results treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area. To control Hyperodes (turfgrass) weevils use twice the rate of Red Panther Dursban Granules shown in the table. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

NOTICE:

Red Panther Chemical Company warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to Red Panther Chemical Company, and buyer assumes the risk of any such use. Red Panther Chemical Company shall not be responsible for incidental or co-incidental damages, if any, resulting from a breach of warranty.



RED PANTHER

DURSBAN GRANULA

TURF AND LAWN INSECTICIDE

CONTROLS BROWN DOG TICKS — CHINCH BUGS. Also Sod Webworms (Lawn Moths), Ants, Crickets, Cutworms, Earwigs, Sowbugs (Pillbugs), Certain White Grubs and Hyperodes (Turfgrass) Weevils infesting Home Lawns and Other Ornamental Turf Areas.

ACTIVE INGREDIENT:

Chlorpyrifos O,O-diethyl O-(3, 5, 6-trichloro-2-pyridyl) phosphorothioate 0.5%

INERT INGREDIENTS: 99.5%
100.0%

*DURSBAN®—Trademark of the Dow Chemical Company
U. S. Patent No. 3,244,586

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

If Inhaled: Remove victim to fresh air and apply respiration if indicated.

If On Skin: Wash promptly with soap and water. Rinse thoroughly.

If In Eyes: Rinse eyes for at least 15 minutes with water and call a physician immediately.

NOTE TO PHYSICIAN

Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine administered by injection only is an antidote.

EPA Reg. No. 42761-3
EPA Est. No. 42761-MS 1

Manufactured By
RED PANTHER CHEMICAL COMPANY
Clarksdale, Mississippi 38614

Net Contents

PRE-AUTOMATIC STATEMENTS
WARNING TO HUMANS AND DOMESTIC ANIMALS
 Do not take internally. Do not get on eyes or on skin. Do not breathe while using. Wash thoroughly after handling. Wash contaminated clothing before reuse. Flush contaminated skin with plenty of water and get medical attention. Avoid breathing spray mist or vapors. Do not spray vegetable gardens or other plants used for food purposes. Keep away from food, feedstuffs, and domestic water supplies. Keep container closed.

ENVIRONMENTAL HAZARDS
 This product is highly toxic to some species of birds, mammals, and aquatic animals. It is also highly toxic to fish, birds, and other wildlife. Keep out of lakes, streams, ponds, rivers, marshes, and wetlands. Spraying and creels may be killed if application rates recommended on this label. Do not apply where birds are reported nesting. Do not spray where runoff is likely to occur. Do not apply where water contamination may occur. Do not spray where birds are nesting or where they are likely to be exposed. Do not apply where birds are nesting or where they are likely to be exposed. Do not apply where birds are nesting or where they are likely to be exposed.

PHYSICAL AND CHEMICAL HAZARDS
CAUTION
 DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. Contains 100% Chlorpyrifos in a Water-based Emulsion.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL
 STORAGE: Keep away from food, feedstuffs, and domestic water supplies. Keep container closed and out of reach of children and pets. DISPOSAL: Do not pour into waterways. Wrap container and put in trash container.

USE PRECAUTIONS
 Use DURSBAN I to control the pests listed in the following table by application of the recommended dosage and in accordance with the directions given below. Check DURSBAN I for complete and in accordance with the directions given below. Check DURSBAN I for complete and in accordance with the directions given below. Check DURSBAN I for complete and in accordance with the directions given below.

Pest	Amount of DURSBAN I per gallon of water	Remarks
Ants	1.0 to 2.0 fl. oz.	Spray when ants are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Beetles	1.0 to 2.0 fl. oz.	Spray when beetles are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Crickets	1.0 to 2.0 fl. oz.	Spray when crickets are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Culicids	1.0 to 2.0 fl. oz.	Spray when culicids are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Flies	1.0 to 2.0 fl. oz.	Spray when flies are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Grasshoppers	1.0 to 2.0 fl. oz.	Spray when grasshoppers are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Houseflies	1.0 to 2.0 fl. oz.	Spray when houseflies are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Leafhoppers	1.0 to 2.0 fl. oz.	Spray when leafhoppers are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Mealybugs	1.0 to 2.0 fl. oz.	Spray when mealybugs are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Midges	1.0 to 2.0 fl. oz.	Spray when midges are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Planthoppers	1.0 to 2.0 fl. oz.	Spray when planthoppers are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Scavenger Beetles	1.0 to 2.0 fl. oz.	Spray when scavenger beetles are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Stinkbugs	1.0 to 2.0 fl. oz.	Spray when stinkbugs are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Termites	1.0 to 2.0 fl. oz.	Spray when termites are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Wasp	1.0 to 2.0 fl. oz.	Spray when wasps are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Whiteflies	1.0 to 2.0 fl. oz.	Spray when whiteflies are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.
Yellowjackets	1.0 to 2.0 fl. oz.	Spray when yellowjackets are active, control when needed. For and outdoors only. Do not spray where birds are nesting or where they are likely to be exposed.



RED PANTHER DURSBAN I

LAWN AND ORNAMENTAL PLANT INSECTICIDE

CONTROLLED PARTS OF YOUR YARD: GRASS, SOFT WOOLYWEED, ALIVE, BROWN DOG TURTLE, CONCRETE, CUTTING, LANTANA, ORNAMENTAL, CANTON WHITE GRASS, HYPERICUM, WHEAT, AND ORNAMENTAL PLANTS (LILIES, BALSAM, CUTTING, ORNAMENTAL, LEAFHOPPERS, MEALYBUGS, MIDGES, GOUTLETTED, GRASSHOPPER, WINDMILL).

Active Ingredients:
 Chlorpyrifos [O,O-Diethyl O-(4,4,4-trichloro-3-hydroxybutyl) phosphorothioate] 12.5%
 Xylene Range Aromatic Solvent 88.3%
 Inert Ingredients 0.2%

WARNING

REPORT OF BLEACHING

MAY BE FATAL IF SWALLOWED • MAY BE ABSORBED THROUGH SKIN • MAY BE IRRITATED TO EYES AND SKIN • CHLORPYRIFOS IS A NEUROTOXICANT

If swallowed: Drink one or two glasses of water and induce vomiting by drinking back of throat with finger. Do not induce vomiting if you are unconscious or if you are pregnant. Get medical attention immediately.

If inhaled: Remove victim to fresh air and apply respiration if indicated. If the victim breathes normally with easy and water, then thoroughly wash the face and hair with water and soap. Do not use a physician or dentist.

Chlorpyrifos is a cholinesterase. Treat symptomatically. Atropine is indicated only in severe cases.

EPA Reg. No. 1274-1
 EPA Est. No. 4264-1

Parts of Ornamentals: Use DURSBAN I to treat flowers, shrubs, trees, shade and flowering trees and evergreens. Do not apply to the parts listed on the label and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage. For best results apply a wetting agent to both upper and lower leaf surfaces and adjacent limbs and trunks. Avoid to contact with foliage but avoid over-spraying to the point of stream runoff. Treat when plants appear and report 11 to 15 days after application. Do not use on plants, complete, permanent, and repeat or re-sprayed by means of possible injury to these plants.

Pest Controlled	Amount of DURSBAN I in Water in Gallons		
	1 gallon	2 gallons	3 gallons
Ants	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Beetles	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Crickets	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Culicids	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Flies	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Grasshoppers	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Houseflies	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Leafhoppers	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Mealybugs	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Midges	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Planthoppers	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Scavenger Beetles	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Stinkbugs	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Termites	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Wasp	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Whiteflies	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.
Yellowjackets	1.0 fl. oz.	2.0 fl. oz.	3.0 fl. oz.

*Equivalent to 1 ounce of chlorpyrifos per 100 gallons
 **Equivalent to 2 ounces of chlorpyrifos per 100 gallons
 ***Equivalent to 3 ounces of chlorpyrifos per 100 gallons

NOTE: Maximum dosage to use on areas treated by control of soil and surrounding plant parts will be listed for various periods of time after treatment. Exceeding an exposure of treated areas to untreated areas.

Additional Use Information:
 Outside Surface of Buildings: Use DURSBAN I at the rate of 1/4 to 1/2 fluid ounce per gallon of water (1.5% chlorpyrifos emulsion) to control ants, silverfish, spiders, and ticks by application to exterior walls, floors, ceilings, and other surfaces of buildings including porches, steps, window frames, doors, utility porches, eaves, and other areas where these pests congregate or have access. Do not use on areas where the spray is likely to contact plants, flowers, plants, shrubs, trees, evergreens, etc. (Exposure of possible injury to these plants or animals is not advised.)

Notice:
 Red Panther Chemical Company warrants that the product conforms to its chemical description and is suitable for the purpose stated on the label when used in accordance with directions under normal conditions of use, but without the warranty of any other warranty of performance or fitness for a particular purpose, right or standard, except to the extent of the label. Red Panther Chemical Company and its agents assume the risk of any such use. Red Panther Chemical Company shall not be responsible for incidental or consequential damage if any resulting from a breach of warranty.

Manufactured by
RED PANTHER CHEMICAL COMPANY
 Charlotte, North Carolina 28211

15