



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS**

CAUTION Harmful if swallowed, inhaled or absorbed through the skin. Do not breathe dust. Do not take internally. Avoid contact with skin. Wear gloves when applying this product. Avoid getting it on eyes. Wash hands before eating. Wash all contaminated clothing with soap and water before reuse. Avoid contamination of foodstuffs, feedstuffs and water receptacles.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct contact with treated plants. Protective information may be obtained from your county cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

GENERAL INFORMATION

Application rates are for pounds per acre or 50 feet of row unless otherwise indicated. Where a dosage range is indicated use the lower rate on small or young plants and the higher rate on large or mature plants. Always use sufficient dust to obtain good coverage of the crop plants being treated. Apply when insects or their damage appear and repeat applications as necessary, or at 7 to 10 day intervals, unless shorter intervals are specified in "DIRECTIONS FOR USE".

This product is a combination of Bacillus Thuringiensis, a biological insecticide, specifically for control of chewing caterpillars and Carbaryl, a broad spectrum insecticide, and will control the following insect pests on vegetables in the home garden:


VEGETABLES

Asparagus — Use 20 to 40 pounds per acre or 1 to 2 oz. per 50 ft. of row for Asparagus Beetle on seedlings or spears. Do not apply later than 1 day before harvesting asparagus. Use 40 to 50 pounds per acre or 2 to 4 oz. to 50 ft. of row for Asparagus Beetle and Apache Cucumber ferns or brush growth in the post-harvest period. Do not apply more often than once every three days.

Beans (green beans, lima beans, snap beans, cowpeas and black-eyed peas) — Use 10 pounds per acre or 1/2 oz. to 50 ft. of row for Mexican Bean Beetle, For Bean Leaf Beetle, Cucumber Beetle, Flea Beetle, Japanese Beetle, Leafhoppers, Velvetbean Caterpillar. Use 40 pounds per acre or 2 oz. to 50 ft. of row. Use 40 lbs. per acre or 2 oz. to 50 ft. of row for Armyworms, Corn Earworm, Stink Bugs and Tarnished Plant Bug.

Corn — Use 40 pounds per acre or 2 oz. to 50 ft. of row for Corn Earworm, Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetle, Sap Beetles and Leafhoppers.

For larvae in whorl. Build up the whorl and damage from top to bottom to entire plant. Repeat as necessary.



**RED PANTHER
GARDEN SPECIAL**

Contains SEVIN and
DIPEL Insecticides

ACCEPTED
SEP 22 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act, this product is amended for the pesticide registered under EPA Reg. No. 4280-100

ACTIVE INGREDIENTS	
Bacillus Thuringiensis, Berliner Potency of 240 International Units per mg. (at least 0.34 billion viable spores per g.)*	0046
Carbaryl (1-naphthyl N-methylcarbamate)	5000
INERT INGREDIENTS	9490
TOTAL	100000

*SEVIN is a registered trademark of Union Carbide Corp. for carbaryl insecticide.
DIPEL is a registered trademark of Abbott Laboratories.
*Equivalent potency: 1,021 million International Units per pound of the product.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SHIELD PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink one or two glasses of water and induce vomiting by sticking back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

If Inhaled: Remove victim to fresh air and apply respiration if indicated.
If On Skin: Wash promptly with soap and water. Rinse thoroughly.

If In Eyes: Rinse eyes for at least 15 minutes with water, and call a physician immediately.

NOTE TO PHYSICIAN: Carbaryl is a non-reversible cholinesterase inhibitor. Atropine is antidotal.

NOTICE:

Red Panther Chemical Company warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions. This warranty does not extend to any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under conditions or in circumstances not reasonably foreseeable to Red Panther Chemical Company, and Buyer assumes the risk of any such use. Red Panther Chemical Company shall not be responsible for any damages or claims of liability resulting from a breach of warranty.

EPA Reg. No. 4280-100
EPA F. N. 4280-100-MS-1
NET WEIGHT

VEGETABLES (Continued)

Brussels Sprouts — Use 20 pounds per acre or 1 oz. to 50 ft. of row for Flea Beetles, Leafhoppers and Squash Bug. Avoid excessive application. Repeat treatments as necessary when tender foliage is wet or in the presence of prolonged high humidity.

Cucumbers — Use 20 pounds per acre or 1 oz. to 50 ft. of row for Flea Beetles, Leafhoppers and Squash Bug. Avoid excessive application. Repeat treatments as necessary when tender foliage is wet or in the presence of prolonged high humidity.

Cucumbers, Melons, Pumpkins and Squashes — Use 20 pounds per acre or 1 oz. to 50 ft. of row for Flea Beetles, Leafhoppers and Squash Bug. Avoid excessive application. Repeat treatments as necessary when tender foliage is wet or in the presence of prolonged high humidity.

Garden Peas — Use 20 pounds per acre or 1 oz. to 50 ft. of row for Leafhoppers, Flea Beetles, Potato Beetles, European Potato Beetle, Colorado Potato Beetle, and Colorado Potato Beetle. Control insect application to avoid contamination in processed peas. Use 40 pounds per acre or 2 oz. to 50 ft. of row for Armyworms.

Tomatoes, Eggplant and Peppers — Use 20 pounds per acre or 1 oz. to 50 ft. of row for Colorado Potato Beetle, Flea Beetles, and Leafhoppers. Use 40 pounds per acre or 2 oz. to 50 ft. of row for European Corn Borer, Fall Armyworm, Lace Bugs, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bugs and Stink Bugs.

Okra — Use 20 to 40 pounds per acre or 1 to 2 oz. to 50 ft. of row for Corn Earworm and Stink Bugs. Repeat at 7 to 10 day intervals or as necessary.

Carrots, Chinese Cabbage, Collards, Dandelion, Endive (Escarole), Garden Beets, Horseradish, Kale, Lettuce, Mustard Greens, Parsnips, Radishes, Rutabagas, Salsify, Spinach, Swiss Chard and Turnips — Use 20 pounds per acre or 1 oz. to 50 ft. of row for Flea Beetles, Harlequin Bug and Leafhoppers. To control Six-Spotted Leafhopper (aster yellow vector) apply 20 to 40 pounds per acre or 1 to 2 oz. to 50 ft. row in a 5 to 7 day schedule. Use 40 pounds per acre or 2 oz. to 50 ft. row for Armyworms, Imported Cabbageworms, Diamondback Moth Larvae, Cabbage Looper, Cross-Striped Cabbageworm, Corn Earworm, Tarnished Plant Bug and Stink Bugs. For optimum control of Corn Earworm after lettuce heads begin to form, use 40 pounds per acre or 2 oz. per 50 ft. row in a 5 to 7 day schedule.

Do not apply to Chinese cabbage, collards, dandelion, endive (escarole), kale, mustard greens, parsley, spinach, or Swiss chard, later than 14 days before harvest for horseradish, parsnips, radishes and rutabagas later than 3 days before harvest.

Do not use on garden beets, turnips, and salsify within 14 days of harvest for top, and 3 days of harvest for roots.

Do not use on head lettuce within 3 days of harvest and leaf lettuce within 14 days of harvest.

Cabbage, Broccoli, Brussels Sprouts, Cauliflower and Kohlrabi — Use 20 pounds per acre or 1 oz. to 50 ft. of row for Flea Beetles and Harlequin Bug.

Use 40 pounds per acre or 2 oz. to 50 ft. of row for Armyworms, Imported Cabbageworm, Diamondback Moth Larvae, Cabbage Looper and Cross-Striped Cabbageworm, and Corn Earworm. Do not apply to these crops later than 3 days before harvest.

Manufactured By

RED PANTHER CHEMICAL COMPANY

Clarkdale, Mississippi 38614