

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

Caution: May be harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Do not take internally. Avoid contact with eyes, skin or clothing. Wear gloves when applying this product. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash thoroughly after handling and before eating or smoking. Wash all contaminated clothing before reuse. Avoid contamination of feed and food-stuffs.

ENVIRONMENTAL HAZARDS

"This pesticide is extremely toxic to aquatic and estuarine invertebrates." Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes."

BEE CAUTION

"This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow to drift to blooming crops or weeds if bees are visiting the treatment area."

STORAGE AND DISPOSAL

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

Application rates are for pounds per acre or 50 feet of row unless otherwise indicated. Where a dosage range is indicated use the lower rate on small or young plants and the higher rate on large or mature plants. Always use sufficient dust to obtain good coverage of the crop plants being treated. Apply when insects or their damage appear and repeat applications as necessary, or at 7 to 10 day intervals, unless shorter intervals are specified in "DIRECTIONS FOR USE."

This product is a combination of Bacillus Thuringiensis, a biological insecticide, specifically for control of chewing caterpillars and Carbaryl, a broad spectrum insecticide, and will control the following insect pests on vegetables in the home garden.

"Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels."

VEGETABLES

Asparagus — Use 20 to 40 pounds per acre or 1 to 2 oz. per 50 ft. of row for Asparagus Beetle on seedlings or spears. Do not apply later than 1 day before harvesting asparagus. Use 40 to 80 pounds per acre or 2 to 4 oz. to 50 ft. row for Asparagus Beetle and Apache Cicada on ferns or brush growth in the pre-harvest period. Do not apply more often than once every three days.

Beans (green beans, lima beans, snap beans, cowpeas and black-eyed peas) — Use 10 pounds per acre or 1/2 oz. to 50 ft. of row for Mexican Bean Beetle, For Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Japanese Beetle, Leafhoppers, Vegetable Caterpillar use 40 pounds per acre or 2 oz. to 50 ft. of row. Use 40 lbs. per acre or 2 oz. to 50 ft. of row for Armyworms, Corn Earworm, Stink Bugs, and Tarnished Plant Bug. Do not apply to cowpeas later than 3 days before harvest.

Corn — Use 40 pounds per acre or 2 oz. to 50 ft. of row for Corn Earworm, Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetle, Sap Beetles and Leafhoppers.

For larvae in whorl (Budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary.

BEST AVAILABLE COPY



RED PANTHER GARDEN SPECIAL

Contains SEVIN® and DIPEL® Insecticides

ACTIVE INGREDIENTS:

Bacillus thuringiensis, Berliner Potency of 240 International Units per mg. (at least 0.38 billion viable spores per g.)*	0.048%
Carbaryl (1-naphthyl N-methyl carbamate)	5.000%
INERT INGREDIENTS:	94.952%

TOTAL 100.000%

Ⓢ SEVIN is a registered Trademark of Union Carbide Corp. for carbaryl insecticide.

Ⓢ DIPEL is a registered trademark of Abbott Laboratories.

*Equivalent to potency of 0.11 billion International Units per pound of this product.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENTS

If Swallowed: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

If Inhaled: Remove victim to fresh air and apply respiration if indicated.

If On Skin: Wash promptly with soap and water. Rinse thoroughly.

If In Eyes: Rinse eyes for at least 15 minutes with water and call a physician immediately.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

NOTICE:

Red Panther Chemical Company warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Red Panther Chemical Company, and buyer assumes the risk of any such use. Red Panther Chemical Company shall not be responsible for incidental or co-incidental damages, if any, resulting from a breach of warranty.

Reg. No. 42761-4

Est. No. 42761-MS-1

NET WT

VEGETABLES (Continued)

For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control.

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

Cucumber, Melons, Pumpkins and Squash — Use 20 pounds per acre or 1 oz. to 50 ft. of row for Pickleworm and Melonworm, Cucumber Beetles, Flea Beetles, Leafhoppers and Squash Bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

Garden Peas — Use 20 pounds per acre or 1 oz. to 50 ft. of row for Leafhoppers and Colorado Potato Beetle. For Colorado Potato Beetle control direct application to weeds in fields of garden peas. This application is intended to help prevent insect contamination in processed peas. Use 40 pounds per acre or 2 oz. to 50 ft. of row for Armyworms.

Potato, Tomato, Eggplant and Peppers — Use 20 pounds per acre or 1 oz. to 50 ft. of row for Colorado Potato Beetle, Flea Beetles and Leafhoppers. Use 40 pounds per acre or 2 oz. to 50 ft. of row for European Corn Borer, Fall Armyworm, Lace Bugs, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bugs and Stink Bugs.

Okra — Use 20 to 40 pounds per acre or 1 to 2 oz. to 50 ft. of row for Corn Earworm and Stink Bugs. Repeat at 5 to 7 day intervals or as necessary.

Carrots, Chinese Cabbage, Collards, Dandelion, Endive (Escarole), Garden Beets, Horseradish, Kale, Lettuce, Mustard Greens, Parsnips, Radishes, Rutabagas, Salsify, Spinach, Swiss Chard and Turnips — Use 20 pounds per acre or 1 oz. to 50 ft. of row for Flea Beetles, Harlequin Bug and Leafhoppers. To control Six-Spotted Leafhopper (aster yellows vector) apply 20 to 40 pounds per acre or 1 to 2 oz. to 50 ft. row in a 5 to 7 day schedule. Use 40 pounds per acre or 2 oz. to 50 ft. row for Armyworms, Imported Cabbageworms, Diamondback Moth Larvae, Cabbage Looper, Cross Striped Cabbageworm, Corn Earworm, Tarnished Plant Bug and Stink Bugs. For optimum control of Corn Earworm after lettuce heads begin to form, use 40 pounds per acre or 2 oz. per 50 ft. row in a 5 to 7 day schedule.

Do not apply to Chinese cabbage, collards, dandelion, endive (escarole), kale, mustard greens, parsley, spinach, or Swiss chard, later than 14 days before harvest or to horseradish, parsnips, radishes and rutabagas later than 3 days before harvest.

Do not use on garden beets, turnips, and salsify within 14 days of harvest for tops and 3 days of harvest for roots.

Do not use on head lettuce within 3 days of harvest and leaf lettuce within 14 days of harvest.

Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi — Use 20 pounds per acre or 1 oz. to 50 ft. of row for Flea Beetles and Harlequin Bug.

Use 40 pounds per acre or 2 oz. to 50 ft. of row for Armyworms, Imported Cabbageworm, Diamondback Moth Larvae, Cabbage Looper and Cross Striped Cabbageworm, and Corn Earworm. Do not apply to these crops later than 3 days before harvest.

Manufactured By

RED PANTHER CHEMICAL COMPANY

Clarksdale, Mississippi 38614