

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING

LITTLE JOHN TOBACCO SPRAY

--- DANGER ---



KEEP OUT OF REACH OF CHILDREN

WARNING---BE SURE TO READ AND UNDERSTAND THE PRECAUTIONARY STATEMENTS BEFORE USING

DIRECTIONS FOR USE

TOBACCO PLANT BEDS: Green Peach Aphids, Cabbage Looper, Tobacco Budworm, Tobacco Hornworm, Tobacco Flea Beetle, Use 2 quarts per 100 gallons of water. Apply 6 gallons of finished spray per 100 sq. yards seed beds.

TOBACCO FIELD: Suckfly--Use 1½ quarts per acre. Green Peach Aphids, Cabbage Looper, Tobacco Budworms, Flea Beetles, Tobacco Hornworms--Use 2 to 4 quarts per acre. Stink Bugs -- Use 4 quarts per acre.

Use the above field dosages in sufficient water to cover one acre. Application of LITTLE JOHN TOBACCO SPRAY should begin when insects first appear and repeat at 3 to 5 days intervals depending upon the severity of infestation and local conditions. Full coverage of plants during application is essential for good control. Do not apply when weather conditions favor drift from areas treated.

To mix, fill the spray tank ¼ to 1/3 full of water and with agitator running, add the required amount of this material, then finish filling the spray tank to the desired level with water.

When applying this material by aircraft, mix the recommended amount with sufficient water to provide a minimum of 2 gallons of finished spray per acre. When applying this material by conventional ground equipment, mix the required amount with sufficient water to provide a minimum of 10 gallons of finished spray per acre.

Make no application within 5 days of priming or 15 days of cutting. Avoid plant juices coming in contact with the skin or other parts of the body of those engaged in priming and cutting the crop. Workers entering treated areas within 24 hours of application should wear protective clothing.

Not for use or storage in or around the home. This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Apply this product only as specified on this label.

Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural extension Service.

Do not use in undiluted form. Do not store below 20° F.

Do not use or store near heat or open flame. Rinse equipment and containers and dispose of wastes by burying at least 18" deep in non-crop lands away from water supplies.

Do not re-use the container. Sewer and bury with wastes.

EPA REG. NO. 4711-13

EPA EST. NO. 240-NC-1

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

34

MANUFACTURED BY

RIVERSIDE CHEMICAL CO.

A SUBSIDIARY OF COOK INDUSTRIES INC. MEMPHIS TN 38117