

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid inhalation of spray mist. Avoid contamination of feed and foodstuffs.

When using this product as a space spray in food processing plants, foods must be removed or covered during treatment. Apply only when the facility is not in operation. All food processing surfaces must be thoroughly cleaned with an appropriate cleaning agent and rinsed with potable water before reuse.

Do not remain in treated areas and ventilate the area after treatment is completed.

Remove pets, birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS: Extremely Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.


STORAGE: Store in a cool, dry, secure area.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

Remove protective cap. Aim spray opening away from person. Push button to spray.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal use, thoroughly spray entire animal as it is being released to pasture. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

POULTRY OPERATIONS: For poultry houses, close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. Treat daily or as necessary.



**CT-25
Dairy Fly Bomb
PRESSURIZED**

For use in Bottling Plants, Breweries, Homes, Bakeries, Food Plants, Stores, Restaurants, Schools, Hotels, Motels, Office Buildings, Hospitals, Warehouses, Food Handling Plants, Dairies, Milk House, Cattle, Horse, Poultry, and Swine Quarters. For use on cattle, horses and swine.

**with DUAL ACTIVATORS
KILLS & REPELS**

House Flies, Horn Flies, Stable Flies, Horse Flies, Face Flies, Deer Flies, Gnats, Mosquitos, Small Flying Moths (millers), Roaches, Waterbugs, Sowbugs, Spiders, Centipedes, Ants, Crickets, Wasps and Hornets.

ACTIVE INGREDIENTS:

Pyrethrins 0.25%

Piperonyl Buloxide, Tech* 1.0%

N-Octyl Bicycloheptene Dicarboximide 1.0%

INERT INGREDIENTS 97.75%

*Equivalent to 0.8% (Butylcarbityl)

(6-Propylpiperonyl) ether and 0.2% of Related Compounds

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Contents Under Pressure

See Side Panels for Additional Precautions

NET WEIGHT: 1.5 lbs.

Mfg. By:
CHEM-TECH, LTD.
4515 Fleur Dr., #303
Des Moines, IA 50321



NOTICE TO USER
LABEL NO. 47000-84
PER PR NOTICE 650-0

DATE: JAN 4 1997

DIRECTIONS FOR USE Con'l.

HOME AND FOOD PROCESSING AREA USE: For flies, mosquitoes, small flying moths (millers), and gnats: Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist then leave the treated areas. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed. **Roaches, Waterbugs, Sowbugs, Spiders and Centipedes:** spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible. Repeat as necessary. **ANTS:** Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

STORED PRODUCT INSECTS (accessible stages): Flying Mediterranean flour moths, Indian meal (chocolate) moths, angoumois grain moths: Close doors and windows. Direct aerosol mist to all parts of the room, especially into dark enclosures and around stored product containers where adult moths may be resting. Use a minimum treatment of six to eight seconds per 1000 cubic feet of space. Keep area closed for 30 minutes after each treatment. Repeat treatment weekly or as flying moths appear. Infested stored food products should be fumigated or treated with an approved product.

DAIRY FARM USE: Milking parlor and milk room. Close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. If rooms cannot be closed, double the dosage. Stand in middle of room. Direct spray in all directions. Keep room closed for 15 minutes after treatment. Do not remain in treated area. Ventilate the area after treatment is completed. Cover milking utensils and milk to prevent contamination from spray and dead or falling insects.

DAIRY ANIMAL USE: From approximately 2 foot distance thoroughly spray entire animal as it is being released to pasture. Do not spray directly toward animal's eyes. Spot treat withers, shoulders and back where saliva accumulates from head tossing.

STANCHION BARN USE: Walk behind animals and direct spray over backs allowing 1 second per animal. For most effective results, apply each morning.

HOG OPERATIONS: In hog houses, close all windows and doors. Apply at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. For animal application direct spray over backs and spray for 1 to 2 seconds per hog. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

HORSE BARN AND STABLES: In horse and pony barns close all windows and doors. Spray at the rate of 1 to 2 seconds per 1000 cubic feet. Keep area closed for 15 minutes following application. If area cannot be closed, double the dosage. In animal application, lightly mist over backs of horses and ponies from a distance of at least 2 feet. Do not spray directly toward horses face or eyes. Repeat application daily or as necessary.

NOTICE TO USER: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the products on these conditions.

EPA Reg. No. 47000-84

EPA Est. No. 47000-IA-01

5/90

BEST AVAILABLE COPY

FLEA BUSTER™ PET POWDER

This is an effective material for use against FLEAS, TICKS, AND LICE on DOGS, CATS and PETS.

ACTIVE INGREDIENTS

Pyrethrins	0.2%
Technical Piperonyl Butoxide*	1.0%
Diatomaceous Earth	88.0%
INERT INGREDIENTS	10.8%
TOTAL	100.0%

*Equivalent to .8% of (butylcarbityl) (6-propylpiperonyl) ether and 2% of related compounds.

NET WT

BATCH #

Manufactured for
DIALECT INTERNATIONAL INC
P.O. BOX 342
SMITH CENTER, KANSAS 66867
EPA Registration No. 42850-3
EPA Establishment No. 58960-KS-1


PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN CAUTION

Avoid contact with eyes and skin. In case of contact immediately flush eyes and skin with water. Get medical attention if irritation persists. Use in adequate ventilation and avoid breathing of dust. Use dust mask for prolonged periods of use. Wash hands thoroughly after using and before smoking or eating.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, ponds or public waters. Do not apply outdoors when weather conditions favor drift from the treatment area. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.



STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting unless directed by a physician. Contact a physician or Poison Control Center immediately.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

STORAGE AND DISPOSAL

STORAGE: Store in original container in cool, dry place out of sunlight. DISPOSAL: PROHIBITION Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Lightly work the material into the hair of the animal, down to the skin. Pay strict attention to places where the animal cannot reach, such as the scalp. Dust the ears of the animal with care. Dust the animal's bedding and areas where the animal sleeps, as well as cracks and crevices adjacent to them. Repeat once a month or as necessary. Individual ticks observed on the animal should be dusted thoroughly.

WARRANTY STATEMENT

Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, seller makes no claims other than those stated on this label.

DATE _____
FEB 17 1990