12850-4 6-16-18 Form Approved, OMB No. 2070-0060, Approval expires 05-31 OPP Identifier Number United States Registration **Environmental Protection Agency Amendment** 253373 Washington, DC 20460 Other Application for Pesticide - Section I 1. Company/Product Number 2. EPA Product Manager 3. Proposed Classification EPA Reg. No. 42850-4 George T. LaRocca Restricted 4. Company/Product (Name) PM# DIATECT MULTIPURPOSE INSECTICIDE II 5. Name and Address of Applicant (Include ZIP Code) 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) DIATECT INTERNATIONAL, INC. (b)(i), my product is similar or identical in composition and labeling c/o RegWest Company P.O. Box 2220; Greeley, CO 80632-2220 EPA Reg. No. Check if this is a new address Product Name Section - II Amendment - Explain below. Final printed labels in response to 05/13/98 Agency letter dated NOTIFICATION Resubmission in response to Agency letter dated_ "Me Too" Application. JUN 1 6 1998 Notification - Explain below. Other - Explain below. Explanation: Use additional page(s) if necessary. (For section I and Section II.) Submit label revision notification to add a Warranty Statement as per PR Notice 95-2 and submit final printed labeling. This notification is consistent with the provisions of PR Notice 95-2 and EPA regulations at 40 CFR § 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. [5] further understand that if this notification is not consistent with the terms of PR Notice 95-2 and 40 CFR § 152.46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. Section - III 1. Material This Product Will Be Packaged in: Child-Resistant Packaging Unit Packaging Water Soluble Packaging 2. Type of Container Metal Yes Yes Yes **Plastic** No Glass Paper If "Yes" No. per "Yes" No. per Certification must Unit Packaging wgt. Package wgt container Other (Specify) be submitted 5. Location of Label Directions 3. Location of Net Contents Information 4. Size(s) Retail Container On Label Container Label On Labeling accompanying product 6. Manner in Which Label is Affixed to Product Lithograph Other Paper glued Stenciled Section - IV 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Telephone No. (Include Area Code) Name Title Kim Davis, CC, APC Consultant/Agent (970) 35360611 6. Date Application Certification Received I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or (Stamped both under applicable law. 2. Signature 3. Title Consultant/Agent

5. Date

June 3, 1998

Kim Davis, CC, APC

4. Typed Name

DIATECT MULTIPURPOSE INSECTICIDE II

ACTIVE INGREDIENTS:

and 0.2% related compounds.

Pyrethrins	0.2%
Piperonyl Butoxide, Technical*	1.0%
Silicon Dioxide	
INERT INGREDIENTS	15.9%
Total	00.0%
*Equivalent to 0.8% (butylcarbityl)(6-propylpiperony	l) ether

KEEP OUT OF REACH OF CHILDREN

CAUTION

See other panels for additional precautionary statements

EPA Reg. No. 42850-4 Net Wt: One (1) lb. EPA Est. 42850-KS-1 Batch No.

Diatect International, Inc. P.O. Box 367 Smith Center, Kansas 66967

WARRANTY STATEMENT

Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment will vary, Seller makes no claims other than those stated on this label.

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

CAUTION: Avoid contact with eyes and skin. Use in adequate ventilation and avoid breathing dust.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothes.

Personal Protective Equipment Applicators and other handlers must wear coveralls or long-sleeved shirt and pants, waterproof gloves, shoes and socks, and a dust/mist filtering respirator (MSHA/NIOSH) approval number prefix TC-21C. Follow manufacturer's instructions for cleaning and maintaining Personal Protective Equipment (PPE). If no instructions for washable are given, use detergent and hot water. Keep and wash PPE separately from other laundry.

Statement of Practical Treatment (First Aid)

If Swallowed: Call a physician or poison control center. Drink one or two glasses of water and induce vormiting by touching back of throat with finger. Do not induce vormiting or give anything by mouth to an unconscious person. If Inhaled: Remove person to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention. If on Skin: Wash affected areas with plenty of soap and water. Get medical attention if irritation persists. If in Eyes: Flush eyes with plenty of water. Contact a obvisician if irritation persists.

Environmental Hazards This product is toxic to fish. Do not apply directly to any body of water. Do not apply when weather conditions favor drift from the treatment area. Do not contaminate water by disposal of equipment wash rinses. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard, and that involves contact with anything that has been treated, such as plants, soil, or water is: coveral(s, water-proof gloves and shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep unprotected persons, children and pets out of the treated area until sprays have dried.

STORAGE AND DISPOSAL

Storage: Store in original container in a cool, dry place out of sunlight. Protect from moisture. Disposal: Do not reuse empty container; wrap and put in trash.

GENERAL INFORMATION: Carefully follow the directions for each type of use. This material may be used on edible crops growing outdoors or in greenhouses, up to and including the day of harvest. PPE required for early entry to treated areas prior to the time the sprays have dried, that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: long-sleeved shirt and long pants, waterproof gloves, shoes and socks. Use rates will vary with insect populations and adequate control should be obtained using one to six pounds per acre of this product.

Use Rates for the Fruits, Nuts, Vegetables, Field Crops, Ornamentals and Miscellaneous Crops: Use 1 to 6 pounds of product per acre depending upon the insect population. The material can be used either: As a dust: The best results are obtained when the plants are wet or dew still remains on the leaf surface. It is suggested that the plants be misted prior to application to obtain the maximum benefits and use the least amount of product for control. Reapply only as necessary; As a wettable powder, Apply in sufficient water to obtain maximum coverage. Dilution of the material should be at the rate of one half pound in one half gailon of water or at a maximum of one pound to 25 gallons of water. It is best to keep the material agitated during application. Apply the material uniformly to both the upper and lower leaf surfaces. Reapply only as necessary.



GRASSES, LAWNS, & TURF (INCLUDING GOLF COURSES)

To control Ants, Armyworms, Caterpitlars, Chinch Bugs, Cutworms, Fire Ants, Fleas and Ticks. Use Rate: Use 16 oz. (1 lb.) per 1000 square feet of area. Apply either as a Dust or as a Wettable Powder.

Fire Ants: Mound Drench or Water Method: Use 0.6 ounces (4 tablespoons) to one gallon of water. For best results, saturate the perimeter of the mound first, working toward the center of the mound in a circular motion (you may want the mixture to puddle). An ant mound 12-14 inches in diameter requires approximately one gallon of mix. Hose End Sprayer: Use 16 oz (1 lb.) per 1000 square feet of area Use a hose end sprayer for mounds in lawns. Add a small amount of water to the jar and add the amount needed for the promeasured area, stir to make a sturry, then add the remaining water to the top of the par Empty entire contents of the jar on the pre-measured area. As a Dust: For best results, dust the perimeter of the mound first. With a stick, disturb the mound, then dust mound in a circular motion working toward the center until the mound is thoroughly dusted.

Ants, Cinch Bugs, Caterpillars, Cut Worms, Armyworms, Fleas and Ticks: Hose End Sprayer: Use 16 oz (1 lb.) per 1000 square feet of area. Use of a hose end sprayer for lawns. Add a small amount of water to the jar and add the amount needed for the pre-measured area, stir to make a sturry, then add the remaining water to the top of the jar. Empty entire contents of the jar on the pre-incasured area. As a <u>Dust</u>: The best results are obtained when applied with a hand or power duster or other suitable means of application to the intested area. Apply lightly and uniformly to the treatment area. Hand Sprayer: Use 0.6. oz (4 tablespoons) to one gallon of water. It is best to keep the material agitated during application. Apply the contents to pre-measured area.

ANIMAL QUARTERS

including but not limited to Barns, Dairies, Milkrooms, Pet Kennets, Poultry Houses, fruns and Stables For control of Gnats, Fleas, Flies, Lice, Mites, Darkling Beetles, Mosquitoes, and Ticks. Use Rate: Use at the rate of 2 oz. per 100 sq. ft. of surface area. Apply with a hand or power duster or sprayer or other suitable means to the hiding places where the posts are found. Repeat treatment as necessary to maintain control. Do not use this product directly on pets.

ON HARVESTED TOMATOES AND FRUIT (including Grapes)

To Control Fruit Flies and Vinegar Flies. Apply with a hand or power sprayer or other suitable means on tomatoes and fruit in baskets, on trucks or on plants, and row stock stacked in the yard. Apply liberally at the rate of 8 oz. per 100 sq. ft. of area.

AS A GRAIN PROTECTANT

To protect against grain storage insects, dilute one pound in 5 gallons of water and apply directly to the grain at a rate of 4 to 5 gallons per 1000 bushels as the grain is carried along a belt or as it enters the auger or elevator. Monthly inspections should be made. If the top two or three inches are found to be reinfested, retreat at the rate of 1 to 2 gallons of dilution per 1000 bushels of stored product.

IN MUSHROOM PRODUCTION AND PROCESSING

To control Mushroom Flies and Fungus Gnats Apply dilution at a rate of 5 ounces per 1000 cubic feet of space. Direct spray 8 to 10 feet away from bods and slightly upwards over beds. Keep circulating fairs running but turn off refrigerators during application. In processing plants, use at the rate of 5 ounces per 2,000 cubic feet of space.

MOSQUITO CONTROL

May be used on croplands as an adulticide to protect agricultural workers during harvesting. Use 2 to 12 ounces per acre when applied by fixed wing aircraft, helicopters or truck mounted equipment. Do not apply in a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

VEGETABLE CROPS

For the control of insects such as: Aphids, 12-Spotted Cucumber Beetle, Armyworms, Beet Webworms, Blister Beetle, Cabbage Looper, Cabbageworms, Caterpillars, Celery Leaftiers, Colorado Potato Beetle, Corn Farworm, Queumber Beetles, Diamondback Moth Larva, European Corn Borer, Flea Beetles, Garden Fleahoppers, Harlequin Bugs, Imported Cabbage Worms, Japanese Beetles, Leafhoppers, Leaf Miners, Leaftiers, Loopers, Lygus Bugs, Mexican Bean Beetle, Mites, Oblique-Banded Leafrollers, Plant Bugs, Stink Bugs, Squash Vine Borers, Thrips, Vegetable Weevils, Webworms, and White Flies.

Growing Crops (Outdoors and Greenhouses): Root and Tuber Vegetables, including but not limited to. Arracacha, Arrowroot, Purple Arrowroot, Japanese Artichoke, Jerusalem Artichoke, Beets, Sugar Beets, Edible Burdock, Carrots, Cassava (Bitter & Sweet), Cerarlac (Celery Root), Chervil (Turnip Root), Chicory, Chotia, Dasheen, Ginger, Ginseng, Horseradish, Laren, Parsley (turnip rooted), Parsnip, Potato, Radish, Japanese Radish, Rutabaga, Salsity, Black Salsity, Sweet Potato, Tamer, Tarrow Root, Turmeric, Turmp, Yam, Yam Bean. Leaves of Root and Tuber Vegetables, including but not limited to, Beet, Sugai Beet, Edible Burdock, Carrot, Cassava (Bitter & Sweet), Celery, Chervil, Chicory, Dasheen, Parsnip, Radish. Japanese Radish, Rutabaga, Black Salsily, Sweet Polato, Tanier, Turnip, and Yam (True) Bulb Vegetables, including but not limited to, Garlic, Leek, Onion (Bulb & Green), Shallot, Leaty Vegetables, including but not limited to, Amaranth, Lealy Amaranth, Chinese Spinach, Tompala, Arugula, Celery, Celluce, Chervil, Cilantro, Corn Salad, Chrysanthemum (edible leaves), Chrysanthemum garland, Cress (garden), Upland Cress (yellow rocket, winter cress), Dandelion, Dock, Endive, Fennel, Lettuce (Head & Leaty), Orach, Parstey, Purstane (garden & winter), Rhubarb, Spinach, Fine Spinach (Metabar, Ceylon), Spinach (New Zealand), Swiss Chard Brassica (Cote) Leafy Vegetables, including but not limited to, Broccoli, Chinese Broccoli,

Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage (Bok Choy, Napa), Chinese Mustard Cabbage (Gai Choy), Caulillower, Collards, Kale, Kohlrabi, Mustard Greens, Rape Greens. Legume Vegetables (Succulent or Dried), including but not limited to, Adzuki Beans, Field Beans, French Beans, Kidney Beans, Lima Beans, Moth Beans, Mung Beans, Navy Beans, Pinto Beans, Runner Beans, Snap Beans, Tepary Beans, Urd Beans, Wax Beans, Asparagus Beans, Black-eyed Peas, Caljang, Chinese Longbeans, Cowpeas, Chowder Peas, Southern Peas, Yard Longbeans, Broad Beans (Fava Beans), Chick Peas (Garbanzo Beans), Guar, Jack Beans (Sword Beans), Lablab Beans (Hycacinth Beans), Lentils, Peas (garden, field, sugar), Pigeon Peas, Soybeans Foliage of Legume Vegetables, including but not limited to, plant parts of any legionie vegetable included in the Legionie Vegetables group that will be used as animal feed including any variety of Beans, Field Peas, Soybeans. Fruiting Vegetables, including but not limited to, Eggplant, Ground Cherry, Okra, Pepinos, Peppers (Bell, Chili, Cooking and Sweet Peppers and Pimentos), Tomatillo, and Torristoes Cucurbit Vegetables, including but not limited to, Balsam Pear (Bitter Melon), Chinese Waxgourd, Cition Melon, Cucumber, Cherkin, Edible Gourds, Melons (including hybrids, Cantaloupe, Casaltia, Crenshaw, Honeydew, Honey Balls, Mango, Muskmelon and Persian Melons), Pumpkin, Squash (summer & winter), Watermelon, (including hybrids)

ORNAMENTALS

For the control of insects such as Aphids, Armyworms, Caterpillars, Chinch Bugs, Flea Beetles, Fleahoppers, Files, Fruit Flies, Japanese Beetles, Leafhoppers, Leafminers, Leatrotters, Loopers, Lygus Bugs, Mealy Bugs, Mites, Plant Bugs, Thrips and White Flies.

Qritamentats, such as Mrican Violets, Aster, Azalea, Begonia, Calceolaria, Calendula, Calia, Camelia, Carnation, Cinerada, Ulrysanthennum, Cypress, Dathold, Dathia, Dogwood, Flin, Eucalyphis, Esti, Ecus, Geranium, Gladiolus, Gypsophila, Holly, Juniper, Lily, Marigold, Oak, Palm, Peony, Petunia, Philodendron, Pine, Roses, Snapdragons, Sweetpeas, Tulips, Viburnum, Wandering Jew, Yew, Zinnia.

HERBS, SPICES AND SPECIALTY CROPS

For the control of such insects as Aphilds, Loopers, Mites, Plant Bugs, Thrips and White Flies.

Herby and Spices, including but not limited to, Anise, Balin, Basil, Burnel, Borage, Chamomile, Caraway, Cathip, Chives, Clary, Corrander, Costmary, Curnin, Curry Leaf, Dill, Fennet (Italian & Sweet), Fenugreek, Horehound, Hyssop, Marigold, Marjoram (Sweet & Wild), Mint, Nasturtium, Oregano, Pennyroyal, Rosemary, Rue, Sage, Savory (Winter & Summer), Sweet Bay (Bay Leal), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood. Specially Crops, such as Artichoke, Chayote, Aspargus, Coffee, Cotton, Hops, Jojoba, Ornamental Turf Grass, Sesame, Sunflower (leaves & seed) and Tea.

FIELD GRAIN CROPS

For the control of insects such as Aphids, Armyworms, Chinch Bugs, Boll Weevil, Bollworm, Budworm, Caterpillars, Corn Earworms, Fleabeetles, Fleahoppers, Flies, Horn Worms, Loopers, Lygus Bugs, Midges, Mites, Pear Cucurlia, Pink Bollworms, Thrips and White Flies.

Coreal Grains, including but not limited to, Barley, Buckwheat, Corn, Millet (proso & pearl), Oats, Popcorn, Rice, Rye, Sorghum (mile). Leosinte, Triticale, Wheat and Wild Rice. Grass Forage, Fodder and Hay, including but not limited to, Bermuda Grass, Blue Grass, Bromegrass, Fescue and any type grass, Gramineae Family (green or cured), Barley, Buckwheat, Corn, Millet (proso & pearl), Oats, Popcorn, Rice, Rye, Sorghum (Milo), Teosinte, Triticate, Wheat, Wild Rice, that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or sitage. Non Grass Animal Feeds, including but not limited to, Alalfa, Velvet Bean, Clover, Kudzu, Lespedeza, Lupine, Saintoin, Trefoil, Vetch, Crown Vetch and Milk Vetch.

FRUIT, NUT, VINE CROPS & ORIENTAL VEGETABLES

For the control of such insects as Aphids, Armyworms, Blueberry Maggots, Cabbage Looners, Caterpillars, Cherry Fruit Flies, Cutworms, Fireworms, Fruit Flies, Fuller Rose Beetle, Gooseberry Fruit Worms, Imported Currant Worms, Japanese Beetles, Leathoppers, Mites, Peach Borers, Pecan Weevil, Red-Necked Borers, Raspberry Fruit Worms, Rose Chafers, Stink bugs, Strawberry Leaf Rollers, Thrips, Weevils, and White Flies.

Citrus Fruits, including but not limited to, Calamondin, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquats, Lemons, Limes, Mandarm (Tangerine), Orange (sweet & sour), Pummelo, and Satsuma Mandarin. Pome Fruits, including but not limited to, Apple, Crabapple, Loquat, Pear, Oriental Pear and Quince Stone Fruits, including but not limited to, Apricot, Cherry (sweet & sour), Nectarines, Peaches, Plum, Prunes, Chicasaw Plums, Damson Plum, and Japanese Plum. Small Fruits and Berries, including but not limited to, Blackberry, Blueberry, Boysenberry, Cranberry, Currants, Dewberry, Elderberry, Gooseberry, Grape Huckleberry, Logariberry, Olahe Berry, Raspberry, (Black & Red), Strawberry, and Youngberry, Subtropical Fruits, including but not limited to, Avocado, Banana, Carob, Barbados Cherry, Cherimova, Dates, Feijoa Figs (Adriatic, Calimyrna, Kadota, Black Mission, California Brown Turkey, Brunswick), Guava, Kiwifruit, Lychee, Marigo, Papaya, Passion Fruit, Persimmon, Pineapple, Pomegranate. Tree Nuts, including but not limited to, Amond, Beech Nut, Brazil Nut, Butter Nut, Cashews, Chestnut, Chinquapin, Filbert (Hazelnut). Tlickory Nut, Japanese Horsechestnut, Macadamia Nut (Bushnut), Pecan, Pistachio, Walnut (Black & 🛬 Hickory Nut, Japanese Horsecheshitu, macadania hut (Change) Parker Broccoti (Gai Lon), Unglish) Oriental Vegetables, including but not limited, to Japanese Artichioke, Clinese Broccoti (Gai Lon), Change Ch Chinese Cabbage (Bok Choy, Napa), Chinese Mustard Cabbage (Gai Choy), Cilantro, Dasheen, Ginger, Ginseng, Chinese Longbeans, Mung Beans, Citron Melon, Balsam Bear (Bitter Melon), Japanese Radish (Daikon), Chinese Spinach, and Chinese Waxgourd.