

ACCEPTED

42871-2
SEP 27 1979

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO.

MICROBIOCIDE 330A

ACTIVE INGREDIENT

Polyoxyethylene (dimethyliminio) ethylene-
(dimethyliminio) ethylene dichloride 15.0%

INERT INGREDIENT 85.0%

This product contains 1.29 lb. of active ingredient per gallon and weighs
8.58 lbs. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact
with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg
whites, gelatin solution, or, if these are not available, drink large quantities
of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of
lakes, streams, or ponds. Permits may be required for discharges contain-
ing this pesticide into lakes, streams, ponds or public water. For guidance,
contact the regional office of the Environmental Protection Agency.

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling.

MICROBIOCIDE 330A is used to control algae, bacteria and fungi in
recirculating commercial and industrial water cooling towers. Prior to its
use, systems must be cleaned to remove algal growth, microbiological
slime, and other deposits. An initial slug addition of 4.0 to 10.0 fluid ounces
of MICROBIOCIDE 330A per 1000 gallons of water to provide a concentra-
tion of 32 to 80 parts per million of MICROBIOCIDE 330A, based on the
total weight of water in the system, is recommended. Repeat initial dosage
until control is evident.

Subsequent slug additions of 1.0 to 10.0 fluid ounces of MICROBIOCIDE
330A per 1000 gallons of water (8 to 80 parts per million of MICROBIO-
CIDE 330A) should be employed every 2 to 5 days, or as needed. The fre-
quency of addition depends upon the relative amount of bleedoff and the
severity of the microbiological problem. Slug additions should be made
in the sump water cooling towers.

MICROBIOCIDE 330A is used to control bacteria in industrial air-wash-
ing systems that maintain effective mist eliminating components. Prior to
its use, systems should be cleaned to remove bacterial slime and other
deposits. An initial slug dose of 14.8 to 24.7 fluid ounces of MICROBIO-
CIDE 330A per 1000 gallons of water is recommended. Repeat initial dose-
age until control is evident. Subsequent slug additions of 10.1 to 24.7 fluid
ounces of MICROBIOCIDE 330A per 1000 gallons of water should be em-
ployed each 1 to 5 days, or as needed. The frequency of addition depends
upon the relative amount of bleedoff and severity of the bacterial problem.
Slug additions may be made to the sump or to the water collection trays
of the airwash system.

STORAGE & DISPOSAL: Keep container closed when not in use. Do not
contaminate water, food, or feed by storage, disposal, or cleaning of
equipment. Do not reuse empty container. Triple rinse (or equivalent) and
dispose of container in an incinerator or in an approved landfill, or bury in
a safe place. Rinse that cannot be used or reprocessed should be dis-
posed of in a landfill approved for pesticides or buried in a safe place
away from water supplies. Open dumping is prohibited.

Manufactured by:

Alde Chemicals, Inc.

5258 North Maroa - Fresno, California 93704

EPA REG. NO. 42871-2

EPA EST. NO. _____