


HAZ 42364-1/ 10/1

DISINFECTANT VIRUCIDE

DISINFECTANT VIRUCIDE



BEST COPY AVAILABLE

ACCEPTED
SEP 21 1995
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 42964-17

AseptiCare® Aerosol Disinfectant Virucide

Air Sanitizer/Odor Counteractant with Residual Bacteriostatic Odor Control

ACTIVE INGREDIENTS:
n-Alkyl (60% C₁₂, 30% C₁₆, 5% C₁₂, 5% C₁₈) dimethyl benzyl ammonium chlorides 0.15%
n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl ethylbenzyl ammonium chlorides 0.15%
Ethanol 62.74%
INERT INGREDIENTS 36.96%
(Includes Essential Oil Odor Control Agents)

**KEEP OUT OF REACH OF CHILDREN
WARNING**

Do not spray in face.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS
WARNING: Avoid getting into eyes. Avoid contamination of food.
First Aid: In case of contact, flush with water. If irritation persists, get medical attention.
Physical and Chemical Hazards: Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

FOR EMERGENCY MEDICAL INFORMATION, CALL TOLL FREE 1-800-328-0026



Net Weight:
14 oz. (397 G)

EPA Reg. No. 42964-17
EPA Est. 334-IL-2

KILLS MOST GERMS—ON ENVIRONMENTAL SURFACES: Staphylococcus, Streptococcus, Pseudomonas, Mycobacterium tuberculosis, Candida albicans, *viruses like: Influenza A2 (Acht), Influenza B, Parainfluenza 1 (Sendai), Vaccinia, Canine distemper, Feline pneumonitis, Herpes simplex (Types 1 and 2), Adenovirus (Type 2), as well as Athlete's Foot fungi and most disease germs. It provides residual bacteriostatic action against odor-causing bacteria if not washed or rubbed off.

* **KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care setting or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS).

Special Instructions for cleaning and decontamination against HIV-1 on surfaces/objects soiled with blood/body fluids:

PERSONAL PROTECTION: When handling items soiled with blood or body fluids, wear personal protection such as disposable latex gloves, gowns, masks, or eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of disinfectant.

CONTACT TIME: 10 minutes.

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

CONTROLS MOLD & MILDEW—AseptiCare controls mold and mildew on hard nonporous surfaces if not washed or rubbed off. **ELIMINATES ODORS AND SANITIZES THE AIR.** AseptiCare temporarily reduces the number of airborne bacteria such as Staphylococcus aureus.

Use AseptiCare on door handles, clean-up carts, light switches, dressing or linen carts, hampers, diaper pails, toilet seats, bedpans, plastic mattress covers, and lockers. AseptiCare is designed for use in hospitals, offices, pet care areas, motels, and other areas where odor and bacterial control is desired.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To Disinfect: Clean surface prior to application. Hold container 6-8 inches from the surface. Spray 3-5 seconds or until surface is wet. Allow to air dry without wiping, or allow to remain on treated surface for 10 minutes before wiping. Avoid using spray on delicate fabrics, lacquered, shellacked, or other finished wood surfaces.

To Control Mold & Mildew: Clean surface prior to application. Spray as above. Repeat application in 7 days or as necessary to maintain control.

To Sanitize Air: Close doors and windows and spray 20-30 seconds in most average size rooms, moving the spray in a circular arc.

To Eliminate Odors: Spray 1-2 seconds in room as required.

AseptiCare is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. AseptiCare may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

STORAGE AND DISPOSAL

Storage: Store in cool, dry place away from flames, sparks or other sources of heat or ignition. Containers heated above 130°F (54°C) may burst. **Disposal:** This container may be recycled in aerosol recycling centers. At present, there are only a few such centers in the country. Before offering for recycling, empty the can by using the product according to the label (DO NOT PUNCTURE!). If recycling is not available, wrap the container and discard in the trash.

ADVERTENCIA: SI NO PUEDE LEER EN INGLES, PREGUNTE A SU SUPERVISOR SOBRE LAS INSTRUCCIONES DE USO APROPIADAS ANTES DE TRABAJAR CON ESTE PRODUCTO

Manufactured For



Airkem Professional Products, Division of ECOLAB INC.
Ecolab Center, St. Paul, MN 55102
9126 Torrance Road, Minneapolis, MN 55425

701319505-0485
Made in U.S.A. © 1995