

2 ACCEPTED

43140-1
APR 3 1980

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

DANGER: Corrosive, may cause severe skin irritation or chemical burns to broken skin. Causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until odors have dissipated.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Treated effluent may not be discharged into lakes, streams, ponds or public waters without a valid discharge permit. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS:
STRONG OXIDIZING AGENT: Mix only with water according to label directions. Mixing this product with gross filth such as feces, urine, etc., or with ammonia, acids, detergents or other chemicals may release hazardous gases irritating to eyes, lungs and mucous membranes.

manufactured by
LIMA POP STOP
7272 West Main Street
Lima, NY 14485

EPA REG. NO. 43140-1 EPA EST. NO.
NET CONTENTS:

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON RESISTANCE
ED UNDER NO.

LIMA

POOL SANITIZER

**"FOR SWIMMING POOL CHLORINATION
AND SANITIZING."**

ACTIVE INGREDIENT: SODIUM HYPOCHLORITE 12.5%
INERT INGREDIENTS 87.5%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

FIRST AID: If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk, or gelatin solution or, if these are not available, drink large quantities of water. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel

**DIRECTIONS
GENERAL CLASS**

**IT IS A VIOLATION OF FEDERAL LAW
TO USE THIS PRODUCT IN A MANNER INCONSISTENT
WITH THE LABEL DIRECTIONS.**

DIRECTIONS FOR SWIMMING POOL CHLORINATION
For a new pool or for spring start-up, superchlorinate with sodium hypochlorite solution for each 3000 gallons of water to 5 ppm available chlorine by weight.
For Pool Maintenance
A) Adjust pool water pH to 7.6 - 7.2 range.
B) Add manually or by a feeder device this product to maintain an available chlorine residual of 1.0 ppm (1.0 ozs.) of this sodium hypochlorite solution per 10,000 gallons of water. A 1.0 ppm available chlorine will depend on pool size. Use test kit to make certain the pH and chlorine levels are correct.

DIRECTIONS FOR DISINFECTING HOME WELLS
Dilute this sodium hypochlorite solution in the solution to 11 parts softened water. Mix thoroughly and begin feeding of solution with a minimum of 1.0 ppm available chlorine residual of at least throughout the distribution system, as determined by bacteriological sampling must be conducted at a frequency of 100 ppm available chlorine will depend on well size. Use a DPD chlorine test kit.
*Contact your local Health Department for further information.
DIRECTIONS FOR SANITIZING OR DAIRY EQUIPMENT

Clean equipment in the normal manner. Just before use, immerse in this sodium hypochlorite solution containing contact with disinfectant for a minimum of 15 minutes after treatment with sodium hypochlorite solution of one fluid oz. (2 tbsps.) of this sodium hypochlorite solution per 100 gallons of water. Rinse thoroughly. NOTE: This product degrades with age. Use as necessary, to obtain the required level of disinfection.

STORAGE AND DISPOSAL
Store in original container in a cool, dry place away from direct sunlight. In case of spill, flood an empty container thoroughly with water and empty by placing in trash collection or burying in an approved landfill. Do not be diluted with water. Do not contaminate food, or feed by storage or disposal.

34

2 ACCEPTED

43140-1
APR 9 1980

3 T

**CAUTIONARY STATEMENTS
PRECAUTIONS TO HUMANS
AND DOMESTIC ANIMALS**

Corrosive, may cause severe skin irritation on broken skin. Causes eye damage. Do not use on clothing. Wear goggles or face shield when handling this product. Wash thoroughly after use. Vacate poorly ventilated areas. Do not return until odors have dissipated.

ENVIRONMENTAL HAZARDS: This pesticide should not be applied to lakes, streams or ponds. Treated effluent should not be discharged into lakes, streams, ponds or public water supply without a discharge permit. For guidance, contact the Environmental Protection Agency.

ADDITIONAL CHEMICAL HAZARDS:
IRRITATING AGENT: Mix only with water. Follow label directions. Mixing this product with acids, urine, etc., or with ammonia, acids, or other chemicals may release hazardous gases and mucous membranes.

Manufactured by
MA POP STOP
West Main Street
Lima, NY 14485

140-1 EPA EST. NO.
CONTENTS:

UNDER THE FEDERAL INSECTICIDE ACT, AS AMENDED
FUNGICIDE AND RODENTICIDE
FOR ECONOMIC PEST CONTROL
REGISTERED UNDER NO.

LIMA

POOL SANITIZER

"FOR SWIMMING POOL CHLORINATION AND SANITIZING."

ACTIVE INGREDIENT: SODIUM HYPOCHLORITE 12.5%
INERT INGREDIENTS 87.5%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

FIRST AID: If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk or gelatin solution or, if these are not available, drink large quantities of water. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel.

**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

DIRECTIONS FOR SWIMMING POOL CHLORINATION

For a new pool or for spring start-up, superchlorinate with one (1) pint (16 fl. oz.) of sodium hypochlorite solution for each 3000 gallons of water. This dosage is equivalent to 5 ppm available chlorine by weight.

For Pool Maintenance:
A) Adjust pool water pH to 7.6 - 7.2 range and maintain.
B) Add manually or by a feeder device this sodium hypochlorite solution at a rate to maintain an available chlorine residual of 0.5 to 1.0 ppm. One-fourth pint (4 fl. oz.) of this sodium hypochlorite solution for each 4000 gallons of water will provide 1.0 ppm available chlorine by weight. Frequency of additions to maintain 0.5 to 1.0 ppm available chlorine will depend on temperature and number of swimmers. Use test kit to make certain the pH and chlorine residual are in the proper range.

DIRECTIONS FOR DISINFECTION OF POTABLE WATER FOR HOME WELL WATER SYSTEMS

Dilute this sodium hypochlorite solution in the ratio of one part sodium hypochlorite solution to 11 parts softened water. Mix sodium hypochlorite solution and water thoroughly and begin feeding of solution with a hypochlorinator (metering pump). Maintain a free available chlorine residual of at least 0.2 ppm and no more than 0.8 ppm throughout the distribution system, as determined by a DPD chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Interim Primary Drinking Water Regulations. Check water frequently with a DPD chlorine test kit.
*Contact your local Health Department for further details.

DIRECTIONS FOR SANITIZING FOOD PROCESSING OR DAIRY EQUIPMENT

Clean equipment in the normal manner. Just before using, rinse all surfaces thoroughly with this sodium hypochlorite solution containing 200 ppm available chlorine. Maintain contact with disinfectant for a minimum of two minutes. Do not rinse with water after treatment with sodium hypochlorite solution. Do not soak overnight. Addition of one fluid oz. (2 tblsp) of this sodium hypochlorite solution per 5 gallons of water will provide approximately 200 ppm available chlorine by weight.
NOTE: This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

STORAGE AND DISPOSAL: Store in a cool, dry area away from direct sunlight. In case of spill, flood area with large quantities of water. Rinse empty container thoroughly with water and either return to manufacturer or discard by placing in trash collection or burying in an approved landfill. Product or residue that cannot be used should be diluted with water and disposed of in a sanitary sewer. Do not contaminate food or feed by storage, disposal or cleaning of equipment.

34

34-A