

4

1. Turn on cold water and open valve on filtered water faucet allowing water to run into sink. Slowly turn the "T" bar on the self-tapping valve (attached to cold water line under counter-see Fig. "A") until water has stopped flowing from filtered water faucet into sink. Double-check to be sure water will not flow through filtered faucet and has been completely shut off at self-tapping valve.

2. Release the spent cartridge from the cartridge holder bracket.

3. Holding the spent cartridge in one hand, turn the nut on the plastic compression fitting located on top of the unit (see Fig. "B") counterclockwise until red vinyl tubing may be easily removed. NOTE: place pan under sink to catch excess water when tubing is disconnected.

4. Repeat same procedure with compression nut on bottom of spent cartridge, releasing blue vinyl tube (see Fig. "C"). Wrap spent cartridge in newspaper and discard with trash.

5. Loosen compression nut on top of new Replacement Cartridge. Insert red vinyl tube into compression fitting and tighten by hand only.

6. Repeat same procedure with compression nut on bottom of new Replacement Cartridge. Insert blue vinyl tube, and tighten by hand only.

7. Place new Replacement Cartridge into cartridge holder bracket and secure.

8. With valve on filtered faucet still open and cold water turned on, slowly open self-tapping valve until unit fills and water runs out filtered faucet into sink. Slowly close self-tapping valve, measuring flow with a container, until flow-rate of 1/2 gallon-per-minute is reached. Important for maximum quality filtration.

9. Let filter run for approximately 10 minutes, watching under counter for leaks. (Do not drink this water). Slightly tighten any fitting that leaks.




10. Complete warranty card and mail immediately.

This filter cartridge is designed for use with the Aqua Guard Bacteriostatic Undercounter Water Filter Model U-602. It should only be used in conjunction with municipally-treated tap water in the home.

AQUA GUARD, INC. • Rt. 15, Box 588 • Tyler, Texas 75707 • (214) 561-4545

JUL 23 1984

Under the Federal Insecticide, Fungicide, and Plant Pesticide Act as amended, for the pesticide registered under EPA Reg. No. 5042-5



46FY
BEST DOCUMENT AVAILABLE

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

DANGER: Corrosive, may cause severe skin irritation or chemical burns to broken skin. Causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until odors have dissipated.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Treated effluent may not be discharged into lakes, streams, ponds or public waters without a valid discharge permit. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS:
STRONG OXIDIZING AGENT: Mix only with water according to label directions. Mixing this product with gross filth such as feces, urine, etc., or with ammonia, acids, detergents or other chemicals may release hazardous gases irritating to eyes, lungs and mucous membranes.

manufactured by

POOL WORLD, INC.
15326 North Nebraska
Tampa, FL 33612

EPA REG. NO. 43141-1 EPA EST. NO.
NET CONTENTS

ACCEPTED

43141-1
NOV 30 1979

UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED

SUPR-CHLOR

**"FOR SWIMMING POOL CHLORINATION
AND SANITIZING."**

ACTIVE INGREDIENT
SODIUM HYPOCHLORITE 9.2%
INERT INGREDIENTS 90.8%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

FIRST AID: If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk, or gelatin solution or, if these are not available, drink large quantities of water. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel.

**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT
IN A MANNER INCONSISTENT WITH ITS LABELING.

DIRECTIONS FOR SWIMMING POOL CHLORINATION

For a new pool or for spring start-up, superchlorinate with one (1) pint (16 fl. oz.) of this sodium hypochlorite solution for each 2500 gallons of water. This dosage is equivalent to 5 ppm available chlorine by weight.

For Pool Maintenance

- A) Adjust pool water pH to 7.6 - 7.2 range and maintain.
B) Add manually or by a feeder device this sodium hypochlorite solution at a rate to maintain an available chlorine residual of 0.6 to 1.0 ppm. One-half pint (8 fl. oz.) of this sodium hypochlorite solution for each 6000 gallons of water will provide 1.0 ppm available chlorine by weight. Frequency of additions to maintain 0.6 to 1.0 ppm available chlorine will depend upon temperature and number of swimmers. Use test kit to make certain the pH and chlorine residual are in the proper range.

**DIRECTIONS FOR DISINFECTION OF POTABLE WATER
FOR HOME WELL WATER SYSTEMS**

Dilute this sodium hypochlorite solution in the ratio of one part sodium hypochlorite solution to 9 parts softened water. Mix sodium hypochlorite solution and water thoroughly and begin feeding of solution with a hypochlorinator (metering pump). Maintain a free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm throughout the distribution system, as determined by a DPD chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed in the National Interim Primary Drinking Water Regulations. Check water frequently with a DPD chlorine test kit.

Contact your local Health Department for further details.

DIRECTIONS FOR SANITIZING FOOD PROCESSING OR DAIRY EQUIPMENT

Clean equipment in the normal manner. Just before using, rinse all surfaces thoroughly with this sodium hypochlorite solution containing 200 ppm available chlorine. Maintain contact with disinfectant for a minimum of two minutes. Do not rinse with water after treatment with sodium hypochlorite solution. Do not soak overnight. Add one fluid oz. (2 fl. oz.) of this sodium hypochlorite solution per 4 gallons of water to provide approximately 200 ppm available chlorine by weight.

NOTE: This product degrades with time. Use a chlorine test kit and increase dosage as necessary to obtain the required level of available chlorine.

STORAGE AND DISPOSAL: Store in a cool, dry area away from direct sunlight. In case of spill, flood area with large quantities of water. Empty container thoroughly with water and either return to manufacturer or dispose by placing in trash collection or burying in an approved landfill. Product or rinseate cannot be used, should be diluted with water and disposed of in a sanitary sewer not for use in food or feed by storage, disposal or cleaning of equipment.

BEST AVAILABLE COPY