

PH 32 43-96-1

BEST AVAILABLE COPY



NOV 8 1982

SUPER SANI 80

Disinfectant - Sanitizer

Fungicide - Virucide - Deodorizer with
Organic Soil Tolerance for Hospital, Institutional,
Industrial, School, Dairy and Other Farm Use

Active Ingredients:	
Octyl decyl dimethyl ammonium chloride	2.250%
Dioctyl dimethyl ammonium chloride	1.125%
Dioctyl dimethyl ammonium chloride	1.125%
Alkyl (C14, 50%, C12, 40%, C16 10%) dimethyl benzyl ammonium chloride	3.000%
Inert Ingredients	92.500%
	100.00%

DANGER: Keep out of reach of children. See side panel for additional cautions.

For sale for use and storage by maintenance personnel only
EPA Registration No. 43198-1 EPA Establishment No. 43198-NM-01

REK-CHEM MANUFACTURING CORP.
121 Virginia N.E. Albuquerque, New Mexico 87108
Phoenix, Arizona-San Diego California 505-883-7695

Sanitizing of Food Processing Equipment and Other Hard Surfaces in Food Contact Locations: For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard non-porous surfaces. **No Potable water rinse is required.**

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. Super Sani 80 in 4 gallons of water (150 ppm active). Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when the solution becomes visibly dirty. For mechanical application, use solution may not be reused for sanitizing applications.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush or mechanical spray device. **No Potable water rinse is required.**

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz./4 gallon solution of Super Sani 80. **No Potable water rinse is required.**

At 1 oz./4 gallons Super Sani 80 fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinance 1978 Recommendations of the U.S. Public Health Service in waters up to 800 ppm of hardness calculated as CaCO3 when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*.

The udders, flanks and teats of dairy cows can be sanitized by washing with a solution of 1 oz. Super Sani 80 in 4 gallons of warm water. **No Potable water rinse is required.** Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

DIRECTIONS FOR USE - GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply Super Sani 80 with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions: For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard non-porous surfaces.

Add 3.5 oz. Super Sani 80 to 4 gallons of water. Apply to previously cleaned hard surfaces. At this use level, Super Sani 80 is effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella choleraesuis* in the presence of 5% blood serum when evaluated by the AOAC Use-Dilution Test.

Virucidal Performance: At a 3.5 oz./4.5 gallon use level Super Sani 80 was evaluated in the presence of 5% serum and found to be effective against the following viruses: Herpes simplex, Vaccinia, and Influenza A2 (Hong Kong) on inanimate environmental surfaces.

Disinfection in Institutions, Industry and Schools: Add 2 oz. of Super Sani 80 to 5.5 gallons of water.

Fungicidal Performance: At 0.5 oz. to 2.25 gallons of water use level, Super Sani 80 is an effective fungicide against *Trichophyton mentagrophytes* (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities, utilizing the AOAC Fungicidal Test.

Disinfection of Barber Tools: Pre-cleaned barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersion in a 0.5 oz./gallon solution of Super Sani 80.

Disinfection of Poultry Equipment, Animal Quarters and Kennels: Poultry brooders, watering founts, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 2 oz. Super Sani 80 to 4.5 gallons of water. Small utensils should be immersed in this solution. Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Precautionary Statements - Hazards to Humans and domestic animals

DANGER: Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Do not breathe spray mist. Avoid contamination of food.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to federal or approved state procedures under subtitle C of the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

GENERAL: Consult Federal, State or Local disposal authorities for approved alternative procedures.

STATEMENT OF PRACTICAL TREATMENT: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

NET CONTENTS - GALLON