




FOSBAN™ 2E INSECTICIDE CONTROLS

Numerous Pests of Households, Turf, and Ornamental Plants. Also
Recommended for Mosquito Control and Area Control of Ticks and Chiggers.

ACTIVE INGREDIENTS: Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 22.4 Aromatic
petroleum derivative solvent 56.41 INERT INGREDIENTS 21.19 Contains 2 pounds of chlorpyrifos per gallon



CONTENTS
1 U.S. GALLON
(3.785 LITER)

KEEP OUT OF REACH OF CHILDREN
WARNING



5651 Dawson Street
Hollywood, Florida 33023

EPA Reg. #43227-16
EPA Est. #43227 FL-01

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

ENVIRONMENTAL HAZARDS: This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service. FOSBAN 2E is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where fish and other aquatic life are important resources or where water is used for irrigation, recreational or domestic purposes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

STORAGE AND DISPOSAL Prohibitions: Do not contaminate water, food, or feed by storage or disposal. **Pesticide Disposal:** Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by federal, state or local disposal authorities. **Container Disposal:** Triple rinse for equivalent and dispose of in an approved landfill or bury in a safe place. **General:** Consult federal, state or local disposal authorities for approved alternative procedures.

**To Be Applied Only By or Under the Supervision of Pest Control Operators,
Public Health Organizations, Mosquito Abatement Districts, and Other
Trained Personnel Responsible for Insect Control Programs.**

KEEP OUT OF REACH OF CHILDREN

WARNING

PRECAUTIONARY STATEMENTS

MAY BE FATAL IF SWALLOWED • MAY BE ABSORBED THROUGH SKIN

• MAY BE INJURIOUS TO EYES AND SKIN

Do Not Take Internally • Do Not Get in Eyes or on Skin • Wash Thoroughly After Handling • Wash Contaminated Clothing Before Reuse • Flush Contaminated Eyes with Plenty of Water and Get Prompt Medical Attention • Avoid Breathing Vapors and Spray Mist • Avoid Spraying Food Crops • Keep Away from Food, Foodstuffs, and Domestic Water Supplies • Keep Container Closed • Handle Concentrate in a Ventilated Area

Statement of Practical Treatment

If Swallowed: Do not induce vomiting. Contains aromatic petroleum solvent. Call a physician immediately. **If On Skin:** In case of contact, remove contaminated clothing and immediately wash skin with soap and water. **If In Eyes:** Flush eyes with plenty of water for 15 minutes. Call a physician. **Note To Physician:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

Physical or Combustible Hazards

COMBUSTIBLE

Do Not Use or Store Near Heat or Open Flame
Do Not Cut or Weld Container

AGRICULTURE CHEMICAL

DO NOT SHIP OR STORE WITH FOODS, FEEDS DRUGS OR CLOTHING

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
**READ AND FOLLOW COMPLETE DIRECTIONS FOR USE AND OTHER IMPORTANT
INFORMATION GIVEN IN BOOKLET ATTACHED TO PRODUCT CONTAINER**



SANEX

Numerous pests of households, turf and ornamental plants. Also recommended for mosquito control and area control of ticks and chiggers.



ENVIRONMENTAL HAZARDS

The Department of State has approved the proposed
 and is hereby authorized to execute the same
 in accordance with the provisions of the
 Act of August 1, 1914, Chapter 136.

[illegible]

USE DIRECTIONS

11. The transfer of the subject matter of the contract to a third party is not permitted.

PESTS Indoors

General Information: Use FOSBAN 2E insecticide to control the pests indicated in the areas listed below by application as a 0.25- or 0.5% chlorpyrifos spray. Use the 0.25% spray to control light infestations and the 0.5% spray to quickly reduce heavy infestations or as directed. To prepare the spray dilute FOSBAN 2E with water or with a suitable base oil as shown in the following **SPRAY DILUTION CHART**. Repeat the treatment as needed.

... ..

NOTE: When you check out, you will be asked to sign a document that says you agree to pay for the books. If you do not have a credit card, you will need to pay for the books in cash. If you do not have cash, you will need to pay for the books in installments. If you do not have a credit card, you will need to pay for the books in cash. If you do not have cash, you will need to pay for the books in installments. If you do not have a credit card, you will need to pay for the books in cash. If you do not have cash, you will need to pay for the books in installments.

CONTENTS

PRECAUTIONARY STATEMENTS	3
Hazards to Humans	3
Statements of Practical Treatment	3
Physical or Chemical Hazards	3
Environmental Hazards	4
PESTS IndoORS	5
General Information	5
Spray Duster Chart	6
Application With:	
Food Handling Establishments	7
Residential and Nonfood Areas	8
Control of Carpet Beetles	8
Control of Brown Dog Ticks	9
PESTS OUT-OF-DOORS	10
General Information	10
Only to Surfaces of Buildings	10
Lawns, Ornamental and Turf Areas	11
Pests of Ornamentals	12
Wood Boring Insects	12
Use on Turf Charts	13
Patch Tree Borer	16
Native Elm Bark Beetle	16
Area Control of Ticks and Chiggers	17
Imported Fire Ants	17
Mosquitoes	18
Use Charts	
Ornamental	19-20
Mosquito Control	21-22
STORAGE AND DISPOSAL	23
NOTICE	23

KEEP OUT OF REACH OF CHILDREN

WARNING
PRECAUTIONARY STATEMENTS

Hazards to Humans
MAY BE FATAL IF SWALLOWED
MAY BE ABSORBED THROUGH SKIN
MAY BE INJURIOUS TO EYES AND SKIN

Do Not Take Internally • Do Not Get In Eyes or on Skin • Wash Thoroughly After Handling • Wash Contaminated Clothing Before Reuse • Flush Contaminated Eyes with Plenty of Water and Get Prompt Medical Attention • Avoid Breathing Vapors and Spray Mist • Avoid Spray on Food Crops • Keep Away from Food, Feedstuffs, and Domestic Water Supplies • Keep Container Closed • Handle Concentrate in a Well-Ventilated Area

Statement of Practical Treatment

[illegible]

Physical or Chemical Hazards
COMBUSTIBLE

**Do Not Use or Store Near Heat
or Open Flame
Do Not Cut or Weld Container**

SPRAY DILUTION CHART		
	Fine Droplets at 0.25" spray	FINEST 25 in dia 0.50" spray
Coarsest Spray	1	2
Medium Spray	3	4
Finest Spray	5	6
Water Spray	7	8

PESTS OUT-OF-DOORS

External Information The following information is available from the following sources:

Pests on Wetland Surfaces of Buildings

[illegible]

4511

As a result, the model is able to capture the effects of the various factors on the response variable. The model is then used to predict the response variable for a given set of input variables. The model is then used to predict the response variable for a given set of input variables. The model is then used to predict the response variable for a given set of input variables.

1. *Pharmaceutical industry* – The pharmaceutical industry is a major source of funding for research in the field of aging. The industry has a vested interest in developing new drugs and treatments for age-related diseases, and it often funds research that is likely to lead to the development of such products.

Feeds of Lyons and other Grapemetal and Macrometall Yarl Gross
Arms

[illegible]

Pulse of Treatment

[illegible]

Worksheet and Instructions for Use on 1097		
Pool	Overage FOURTH 22 per 1000 sq ft	Remarks
Blumgart's Building	22 0.00	Surveys of this building are not required. No additional fees.

Worksheet and Instructions for Use on 1097		
Pool	Overage FOURTH 22 per 1000 sq ft	Remarks
Blumgart's Building	22 0.00	Surveys of this building are not required. No additional fees.

Pool	Dosage FORMAN 2E per 1000 sq. ft.	Remarks
Turkeygrass (Hydrocotyle weevil)	3 to 4 oz.	Spray suspected areas immediately after rain. May or as recommended by your local Agricultural Extension Service specialist for your area.
White grubs (European chaffer and Japanese beetle larvae only)	3 to 6 oz.	For white grubs spray when grubs are young and active feeding near the soil surface, usually during late July or August or as recommended by your local Agricultural Extension Service specialist for your area. Immediately after spraying irrigate the treated area with 1/2 to 1 in. of water to wash the insecticide in the underlying soil.

Pool	Dosage FORMAN 2E per 1000 sq. ft.	Remarks
Turkeygrass (Hydrocotyle weevil)	3 to 4 oz.	Spray suspected areas immediately after rain. May or as recommended by your local Agricultural Extension Service specialist for your area.
White grubs (European chaffer and Japanese beetle larvae only)	3 to 6 oz.	For white grubs spray when grubs are young and active feeding near the soil surface, usually during late July or August or as recommended by your local Agricultural Extension Service specialist for your area. Immediately after spraying irrigate the treated area with 1/2 to 1 in. of water to wash the insecticide in the underlying soil.

Pool	Dosage FOSPH 2E per 1000 sq. ft.	Remarks
Ants	1 1/2 oz	Spills when pests first appear retreat when needed for 500 methomyls delay water spray follow the treated area at 12 to 24 hours after treatment
Armyworms		
Chiggers		
Crunch bugs		
Clover mite		
Crickets		
Cutworms		
Earwigs		
Fleas		
Grasshoppers		
Sod webworms		
Ticks		

Pool	Dosage FOSPH 2E per 1000 sq. ft.	Remarks
Ants	1 1/2 oz	Spills when pests first appear retreat when needed. For Sod webworms delay watering until after the treated area is 12 to 24 hours after treatment.
Armyworms		
Chiggers		
Crunch bugs		
Clover mite		
Crickets		
Cutworms		
Earwigs		
Fleas		
Grasshoppers		
Sod webworms		
Ticks		

Special Selling Tactics Including Smiths and Tiers used in Depressed Areas
Depressed Areas—The following are the names of the various depressed areas in the United States, as designated by the Federal Reserve Board, and the names of the various banks operating in these areas:

What About Insects Infesting Shrubs and Trees such as Dogwood

Amount of FOSPHATE or Water to Make			
Pests Controlled	3 gallons	10 gallons	100 gallons
Apple	1.0	1.0	1.0
Banana	1.0	1.0	1.0
Cashew	1.0	1.0	1.0
Citrus	1.0	1.0	1.0
Cocoa	1.0	1.0	1.0
Coffee	1.0	1.0	1.0
Custard Apple	1.0	1.0	1.0
Guava	1.0	1.0	1.0
Jackfruit	1.0	1.0	1.0
Lychee	1.0	1.0	1.0
Mango	1.0	1.0	1.0
Orange	1.0	1.0	1.0
Pineapple	1.0	1.0	1.0
Rubber	1.0	1.0	1.0
Sugarcane	1.0	1.0	1.0
Tea	1.0	1.0	1.0
Tomato	1.0	1.0	1.0
Watermelon	1.0	1.0	1.0

Amount of FOSPHATE or Water to Make			
Pests Controlled	3 gallons	10 gallons	100 gallons
Apple	1.0	1.0	1.0
Banana	1.0	1.0	1.0
Cashew	1.0	1.0	1.0
Citrus	1.0	1.0	1.0
Cocoa	1.0	1.0	1.0
Coffee	1.0	1.0	1.0
Custard Apple	1.0	1.0	1.0
Guava	1.0	1.0	1.0
Jackfruit	1.0	1.0	1.0
Lychee	1.0	1.0	1.0
Mango	1.0	1.0	1.0
Orange	1.0	1.0	1.0
Pineapple	1.0	1.0	1.0
Rubber	1.0	1.0	1.0
Sugarcane	1.0	1.0	1.0
Tea	1.0	1.0	1.0
Tomato	1.0	1.0	1.0
Watermelon	1.0	1.0	1.0

STORAGE AND DISPOSAL

Particle Disposal — The particle disposal system is designed to handle the waste generated by the particle beam. It consists of a series of collection and disposal units that are connected to the particle beam. The waste is collected in a series of containers and then disposed of in a safe and secure manner.

[illegible]

Condition of Area to be Treated	Amount to Use per Acre		Number Acres 1 Gallon of FOSBAN ZE Will Treat
	FOSBAN ZE (fluid ounces)	Chlorpyrifos (pounds)	
Aspen Forests Light to medium understory cover	16	0.025	80
Mountain to heavy understory cover	32 1/2	0.5 1/2	40

* For higher dosage and also for faster knockdown and longer residual control where there is more or less vegetation cover.

Condition of Area to be Treated	Amount to Use per Acre		Number Acres 1 Gallon of FOSBAN ZE Will Treat
	FOSBAN ZE (fluid ounces)	Chlorpyrifos (pounds)	
Aspen Forests light to medium understory cover	16	0.025	80
Mountain to heavy understory cover	32 1/2	0.5 1/2	40

* For higher dosage and also gun type knockdown and longer residual control where there
is no or low vegetation cover

* The higher dosage will also give faster knock down and longer residual control where there is no vegetative cover

Conditions of Area to be Treated	Amount to Use per Acre		Number Acres ± Gallons of FOSMAN 2E Will Treat
	FOSMAN 2E (Pound ounces)	Chlorpyrifos (ounces)	
1. 100% of area to be treated	8.0 - 10.0	1.0 - 1.5	1000 - 1500
2. 50% of area to be treated	4.0 - 5.0	0.5 - 0.75	500 - 750

Conditions of Area to be Treated	Amount to Use per Acre		Number Acres ± Gallons of FOSMAN 2E Will Treat
	FOSMAN 2E (Pound ounces)	Chlorpyrifos (ounces)	
1. 100% of area to be treated	8.0 - 10.0	1.0 - 1.5	1000 - 1500
2. 50% of area to be treated	4.0 - 5.0	0.5 - 0.75	500 - 750

Plants Controlled		Amount of FOSBAN 2E in Water to Use		
		3 gallons	10 gallons	100 gallons
Annual weeds	3-4" high	2 1/2 oz.	6 1/2 oz.	1 1/2 lb.
Annual weeds	4-6" high	3 1/2 oz.	9 1/2 oz.	2 1/4 lb.
Annual weeds	6-8" high	4 1/2 oz.	12 1/2 oz.	3 1/4 lb.
Annual weeds	8-10" high	5 1/2 oz.	15 1/2 oz.	4 1/4 lb.
Annual weeds	10-12" high	6 1/2 oz.	18 1/2 oz.	5 1/4 lb.
Annual weeds	12-14" high	7 1/2 oz.	21 1/2 oz.	6 1/4 lb.
Annual weeds	14-16" high	8 1/2 oz.	24 1/2 oz.	7 1/4 lb.
Annual weeds	16-18" high	9 1/2 oz.	27 1/2 oz.	8 1/4 lb.
Annual weeds	18-20" high	10 1/2 oz.	30 1/2 oz.	9 1/4 lb.
Annual weeds	20-22" high	11 1/2 oz.	33 1/2 oz.	10 1/4 lb.
Annual weeds	22-24" high	12 1/2 oz.	36 1/2 oz.	11 1/4 lb.
Annual weeds	24-26" high	13 1/2 oz.	39 1/2 oz.	12 1/4 lb.
Annual weeds	26-28" high	14 1/2 oz.	42 1/2 oz.	13 1/4 lb.
Annual weeds	28-30" high	15 1/2 oz.	45 1/2 oz.	14 1/4 lb.
Annual weeds	30-32" high	16 1/2 oz.	48 1/2 oz.	15 1/4 lb.
Annual weeds	32-34" high	17 1/2 oz.	51 1/2 oz.	16 1/4 lb.
Annual weeds	34-36" high	18 1/2 oz.	54 1/2 oz.	17 1/4 lb.
Annual weeds	36-38" high	19 1/2 oz.	57 1/2 oz.	18 1/4 lb.
Annual weeds	38-40" high	20 1/2 oz.	60 1/2 oz.	19 1/4 lb.
Annual weeds	40-42" high	21 1/2 oz.	63 1/2 oz.	20 1/4 lb.
Annual weeds	42-44" high	22 1/2 oz.	66 1/2 oz.	21 1/4 lb.
Annual weeds	44-46" high	23 1/2 oz.	69 1/2 oz.	22 1/4 lb.
Annual weeds	46-48" high	24 1/2 oz.	72 1/2 oz.	23 1/4 lb.
Annual weeds	48-50" high	25 1/2 oz.	75 1/2 oz.	24 1/4 lb.
Annual weeds	50-52" high	26 1/2 oz.	78 1/2 oz.	25 1/4 lb.
Annual weeds	52-54" high	27 1/2 oz.	81 1/2 oz.	26 1/4 lb.
Annual weeds	54-56" high	28 1/2 oz.	84 1/2 oz.	27 1/4 lb.
Annual weeds	56-58" high	29 1/2 oz.	87 1/2 oz.	28 1/4 lb.
Annual weeds	58-60" high	30 1/2 oz.	90 1/2 oz.	29 1/4 lb.
Annual weeds	60-62" high	31 1/2 oz.	93 1/2 oz.	30 1/4 lb.
Annual weeds	62-64" high	32 1/2 oz.	96 1/2 oz.	31 1/4 lb.
Annual weeds	64-66" high	33 1/2 oz.	99 1/2 oz.	32 1/4 lb.
Annual weeds	66-68" high	34 1/2 oz.	102 1/2 oz.	33 1/4 lb.
Annual weeds	68-70" high	35 1/2 oz.	105 1/2 oz.	34 1/4 lb.
Annual weeds	70-72" high	36 1/2 oz.	108 1/2 oz.	35 1/4 lb.
Annual weeds	72-74" high	37 1/2 oz.	111 1/2 oz.	36 1/4 lb.
Annual weeds	74-76" high	38 1/2 oz.	114 1/2 oz.	37 1/4 lb.
Annual weeds	76-78" high	39 1/2 oz.	117 1/2 oz.	38 1/4 lb.
Annual weeds	78-80" high	40 1/2 oz.	120 1/2 oz.	39 1/4 lb.
Annual weeds	80-82" high	41 1/2 oz.	123 1/2 oz.	40 1/4 lb.
Annual weeds	82-84" high	42 1/2 oz.	126 1/2 oz.	41 1/4 lb.
Annual weeds	84-86" high	43 1/2 oz.	129 1/2 oz.	42 1/4 lb.
Annual weeds	86-88" high	44 1/2 oz.	132 1/2 oz.	43 1/4 lb.
Annual weeds	88-90" high	45 1/2 oz.	135 1/2 oz.	44 1/4 lb.
Annual weeds	90-92" high	46 1/2 oz.	138 1/2 oz.	45 1/4 lb.
Annual weeds	92-94" high	47 1/2 oz.	141 1/2 oz.	46 1/4 lb.
Annual weeds	94-96" high	48 1/2 oz.	144 1/2 oz.	47 1/4 lb.
Annual weeds	96-98" high	49 1/2 oz.	147 1/2 oz.	48 1/4 lb.
Annual weeds	98-100" high	50 1/2 oz.	150 1/2 oz.	49 1/4 lb.

Plants Controlled		Amount of FOSBAN 2E in Water to Which		
		3 gallons	10 gallons	100 gallons
Annual weeds	3.00	1.00	1.00	1.00
Perennial weeds	3.00	1.00	1.00	1.00
Grasses	3.00	1.00	1.00	1.00
Shrubs	3.00	1.00	1.00	1.00
Trees	3.00	1.00	1.00	1.00
Other plants	3.00	1.00	1.00	1.00