



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Avoid inhalation. May be absorbed through skin. Do not apply to household pets or humans. If sprayed on skin, or in eyes, wash thoroughly. Harmful if swallowed. Do not use on lactating dairy animals or in dairy barns or milking rooms. Do not use on newborn, very young animals or calves under 3 months old. Areas near eyes should be treated with extreme caution. Avoid contamination of food or feedstuffs. Use minimum amount of material to treat wounds.

PHYSICAL OR CHEMICAL HAZARDS

Flammable! Contents under pressure. Do not use near fire, sparks or flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not store at temperatures below 35°F. If container is exposed to freezing temperatures, can should be warmed at room temperature for four hours before use.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.
If on skin: Remove contaminated clothing and wash affected areas with soap and water.
If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABEL.
GENERAL DIRECTIONS: Remove safety cap, hold can about 4 to 8 inches from area to be treated with nozzle pointing at wound. Spray the area until it is completely wet. 10 to 15 seconds are usually enough. This product is not recommended for use on deep wounds, as directed on superficial, surface wounds or as an antiseptic treatment. Some strains of screwworms may develop resistance to lindane. This will be observed by lack of response to treatment.
INFESTED SUPERFICIAL WOUNDS: Spray the wound until it is completely wet. The area around the wound should also be sprayed with particular attention to that area below the injury where drainage might occur.
NEW UNINFESTED SURFACE WOUNDS: New superficial wounds should be sprayed promptly to help in preventing screwworm infestation. For minor wounds such as slight cuts, scratches, abrasions, etc., spray them once as directed.
Treatment should be repeated two times a week until healing is complete. Two treatments may be sufficient if no drainage occurs. Flies attracted to the wound for several days after treatment will normally be killed by the insecticide. EAR TICKS: Thoroughly spray or immerse the ear in the solution enough to spray through the ear and back of the head. Massage the ear thoroughly with coverage inside the ear. The ear canal with spritz solution. After 10 to 15 seconds indicate that a thorough treatment should keep ear free of ticks for about 30 days. Retreatment after 10 days is recommended.

DISPOSAL

Do Not Re-Use Container. Break Container in Newspaper and Put in Trash Can.
EPA Reg. No. 43286-13 EPA Est. 43286-TX-1

OMNI LINDANE SCREWORM SPRAY

A pressurized Screwworm Killer and Ear Tick Treatment providing a convenient means of killing these pests of beef cattle, horses, swine, sheep and goats.

ACTIVE INGREDIENT:

Gamma Isomer of Benzene Hexachloride
from Lindane 3.0%

INERT INGREDIENTS:

97.0%

Blue Color Added
Prevent From Freezing

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Side Panel For Additional Cautions.

Net Weight: 10 Oz.

**OMNI DEVELOPMENT
Houston, Tx 77090**