

# Spartan

## Formula "O"

### ACTIVE INGREDIENTS

Dissodium cyanodithiocarbamate ..... 1.00%  
Potassium N-methylidithiocarbamate ..... 1.15%

### INERT INGREDIENTS

82.85%

### TOTAL

100.00%

This product contains 1.67% of active ingredients per gallon and weighs 9.2 lbs. per gallon.

### KEEP OUT OF REACH OF CHILDREN

### WARNING

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING:** Causes eye irritation and skin irritation. Avoid contact with skin or on clothing. Wear goggles on face, splash water into eyes immediately. Harmful if swallowed. Avoid contamination of food.

**FIRST AID:** In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, give patient 1/2 cup of powdered charcoal immediately, or 1/2 cup of water with 1/2 raw egg white, milk, or oil, or flour and water. Then induce vomiting with salt water or mustard in warm water. Call a physician immediately.

**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to fish and other aquatic life. Do not discharge into lakes, streams, ponds, or other waters unless you are in compliance with an NPDES permit. For guidance, contact your Regional Office of the EPA.

### DIRECTIONS FOR USE GENERAL CLASSIFICATION

This is a restricted use pesticide. Do not use this product in a manner inconsistent with its labeling.

Spartan Formula "O" is a systemic fungicide for use on ornamental plants and trees. It is a systemic fungicide that is absorbed by the plant and moves through the vascular system. It is effective against a wide range of fungi and other plant pathogens. It is also effective against some insects and mites. An application of 1.0 to 1.5 gallons of Spartan Formula "O" per 1000 square feet of water will provide adequate control of most plant diseases. Control is evident within 7 to 14 days after application.

Reapply at intervals of 14 to 21 days. Apply 1.0 to 1.5 gallons of Spartan Formula "O" per 1000 square feet of water. Reapply every 1 to 3 days, or as needed. The frequency of application depends on the relative amount of bleedoff and the severity of the disease. Additional spray applications should be made in the summer months.

**ACCEPTED**  
22 MAY 1980

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTRY  
EPA REG. NO. 43392-1

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not store in water, or use for food, or storage, or disposal of hazardous waste.

**PESTICIDE DISPOSAL:** Do not use spray nozzles or hoses that cannot be used for other purposes. They should be disposed of in a landfill approved for hazardous waste disposal. Do not mix with water supplies.

**CONTAINER DISPOSAL:** Do not reuse, or recycle, and offer for recycling. Do not use for disposal of hazardous waste. Do not bury in a safe place. Metal containers should be triple-rinsed, sealed, and offered for recycling.

**GENERAL:** Do not use in areas where disposal authorities for approved pesticides are not available.

Manufactured by  
**SPARTAN ENVIRONMENTAL SERVICES, INC.**  
43 West 42nd Avenue  
Brooklyn, N.Y. 11220

EPA REG. NO. 43392-1

EPA EST. NO.

NET CONTENTS

25