



DELTA BT
Biological Insecticide

ACCEPTED
AUG 5 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 43567-1

Active ingredients:	
Bacillus thuringiensis, Berliner Strain	26.4%
Potency:	
130 thousand International Units per mg.	
125 billion IU per qt	
(500 billion IU per gal.)	
Inert ingredients	73.6%
	100%

PRECAUTIONARY STATEMENT
Keep Out of Reach of Children

HAZARDOUS TO HUMANS: Avoid inhalation or contact with eyes or open wounds. Avoid breathing spray mist.

ENVIRONMENTAL HAZARD WARNING: Avoid unnecessary use near lakes, ponds, or streams.

STORAGE AND DISPOSAL

DISPOSAL: Do not reuse empty container. Dispose of in a sanitary land fill or by incineration if approved by state and local authorities. Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Delta BT must be stored in a cool place. Do not allow spray solution to stand more than 12 hours before use.

Mfg by Delta Biological Products, Inc
Box 133
Mer Rouge, La 71261
EPA Establishment No. 43567-LA-01
EPA Registration No. 43567-1

NET CONTENTS 1 Gal

RECEIVED
RUC 3

Delta BT

1. Not harmful to beneficial insects.
2. Not harmful to plants.
3. Treated worms usually stop eating within one hour of ingestion and die within 1-5 days.
4. A suitable exempt spreader sticker may be added to Delta BT.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Delta BT is a highly concentrated insecticide. The recommended rate for control of the following pests on the following crops is:

10 billion IUs of Delta BT per acre — OR — 1 gallon per 50 acres

SHAKE WELL BEFORE USE

Delta BT is recommended for aerial applications at rates of 1-5 gallons total spray solution per acre and for ground spray machine application at rates of 10-20 gallons total spray solution per acre. Complete coverage of plants is essential for effective pest control by Delta BT.

CROP	PEST
Cotton	Cotton bollworm Tobacco budworm Cabbage looper Beet armyworm
Soybean	Green cloverworm Velvetbean caterpillar Soybean looper Cabbage looper Cotton bollworm
Tobacco	Tobacco budworm Cabbage looper Horn worm
Garden Vegetables beans, broccoli, brussel sprouts, cabbage, cauliflower, celery, cucumbers, lettuce, mustard greens, melons, potatoes, spinach, tomatoes, turnip greens	Cabbage looper Imported cabbageworm Diamondback moth
Greenhouse Flowers	Cabbage loopers Omnivorous looper
Rangeland, Pastureland pasture, hay, small grains, forage crops, rangeland	European Skipper Rangeland caterpillar Fall armyworm
Orchards, Fruit Trees, Forest Land pecan, walnut, almond, apple, orange, shade trees, spruce	Fall webworm Red humped caterpillar Tent caterpillar Leaf rollers Orangedog Fruitree leafroller Eastern spruce budworm Gypsy moth Bag worm

*For hand sprayers used to treat garden vegetables use:
1/2 ounce or approximately 15 milliliters of Delta BT in a gallon of water.

**Trees should be sprayed using high volumes of water to provide uniform coverage.

Notice: Delta Biological Products, Inc. makes no warranty, expressed or implied, concerning this material except those which are contained on the label.