

DELTA BT

DELTA BT-BIOLOGICAL INSECTICIDE

ACTIVE INGREDIENTS:

<u>Bacillus thuringiensis</u> , subspecies <u>kurstaki</u>	
Lepidopteran active toxin	26.4%

INERT INGREDIENTS:	73.6%
	100.0%

EPA REGISTRATION NO. 43567-1
 EPA ESTABLISHMENT NO. 43567-LA-01

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT:

If swallowed: Induce vomiting and call a physician.

If on skin: Wash thoroughly with soap and water.

If in eyes: Hold eyelids open and flush with copius amounts of clean water for 10-15 minutes.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Avoid inhalation or contact with eyes. Avoid breathing spray mist.

ENVIRONMENTAL HAZARDS: Do not contaminate water when disposing of equipment washwaters.


SHAKE WELL BEFORE USING

MANUFACTURED BY:
 Delta Biological Products, Inc.
 Rt. 1, Box 639, Airport Road
 Bastrop, La. 71220

NET CONTENTS:

1 Quart

-1-



JUN 04 1992

43567-1


STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool place. Do not allow spray solution to stand more than 12 hours before use. Do not use or store near heat and/or open flame.

PESTICIDE DISPOSAL: Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent), adding rinsate to spray tank. Offer rinsed containers for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Do not reuse container.



Delta BT is a highly selective, highly concentrated biological insecticide for use against larvae (caterpillars or worms) of lepidopterous insects. Delta BT must be eaten by the targeted larvae to be effective. Once a lethal dose is ingested, Delta BT paralyzes the midgut of the larvae. Usually within 1 hour the larvae stop feeding, thus crop damage is halted. Affected larvae move slowly, discolor, shrivel, darken and die from starvation within several days.

Larvae must be actively feeding on treated, exposed plant parts. Complete coverage of plant foliage where larvae are feeding is essential for effective pest control. A suitable spreader sticker, cleared for use on growing and harvested crops, may be added to increase spray coverage, uniform deposition, and adherence. Treat when larvae are young (1st and 2nd instars) before crop is extensively damaged. Monitoring insect infestation by close scouting and early attention to infestation is highly recommended. For heavy infestation and mature worms, use the higher label rates. Re-treat at intervals necessary to maintain control, usually 3-14 days depending on factors such as plant growth rate, rainfall after treatment, and moth activity. Delta BT may be tank mixed with other labelled insecticides to enhance control when mixed or overlapping generations of larvae are present. Check tank mix compatibility of Delta BT and other insecticide before tank mixing.

Delta BT does not harm Honeybees or other beneficial or predaceous insects.

DIRECTIONS FOR USE:


It is a violation of federal law to apply this product in a manner inconsistent with its labelling.

Do not apply this product through any type of irrigation equipment.

Delta Bt may be applied up to the day of harvest.

Delta BT may be applied by aerial equipment or conventional ground equipment with quantities of water necessary to ensure complete, uniform coverage of infested plant foliage. Apply Delta BT in at least 1 to 10 gallons of total spray volume per acre by air and in at least 10 to 20 gallons of total spray volume per acre by ground. Higher spray volumes increase coverage. Ground application on trees should be made with 100 to 400 gallons of spray volume. Add Delta BT to agitated spray solution. Agitate tank mixes thoroughly and continuously.

Do not mix more Delta BT than can be used in a 12 hour period.



The following pests may be treated with Delta BT. Complete coverage of plants and trees is essential for effective control of the following pest:

CROP	PEST	RATE (fl.oz. per acre)
Cotton, tobacco, peanuts, safflower, soybeans, sunflower, sugarbeets, rice	Cotton bollworm, tobacco budworm, cabbage looper, armyworm, green cloverworm, hornworm, saltmarsh caterpillar, velvetbean caterpillar, soybean looper	2.5-4

- NOTE:**
- For heavy infestations and mature worms use up to 4.0 fl. oz. per acre (1 gal. per 32 acres).
 - For additional control, tank mix Delta BT with a labelled ovicide or other labelled products such as pyrethroids, carbamates, organophosphates in accordance with labelling recommendations.

Vegetables: beans, broccoli, brussel sprouts, cabbage, cauliflower, celery, cucumbers, lettuce, mustard greens, melons, potatoes, spinach, tomatoes, turnip greens, beets, carrots, chard, chinese cabbage, cole crops, cucurbits, endives, escarole, eggplant, kale, kohlrabi, lentils, mint, onions, parsley, peas, peppers, turnips, watermelons	Cabbage looper, imported cabbage worm, diamondback moth, corn earworm, green cloverworm, melonworm, rindworm, pickleworm, saltmarsh caterpillar, variegated cutworm, webworm.	2.5-4
--	--	-------

- NOTE:**
- For heavy infestations and mature worms use up to 4 fl. oz. per acre (1 gallon per 32 acres).
 - For additional control, tank mix Delta BT with a labelled ovicide or other labelled products such as pyrethroids, carbamates, organophosphates in accordance with labelling recommendations.

- For hand held sprayers used to treat vegetables use: ½ fl. oz. or approximately 15 milliliters of Delta BT in a gallon of water.


<p>Greenhouse flowers, ornamentals, plants, shade trees, forests</p>	<p>Cabbage loopers, omniverous looper, bagworm, cankerworm, oakworm, oleanderworm, spanworm, saddleback caterpillar, tussock moth, armyworm, gypsy moth, spruce budworm, bedworm, diamondback moth, hornworm, leafroller, red-humped caterpillar, tent caterpillar</p>	<p>2.5-4</p>
---	--	--------------

- NOTE:**
- For heavy infestations and mature worms use up to 4 fl. oz. per acre (1 gallon per 32 acres).
 - For hand held sprayers used to treat vegetables use: ½ fl. oz. or approximately 15 milliliters of Delta BT in a gallon of water.

<p>Rangeland, Pastureland pasture, hay, small grains, forage crops, rangeland, alfalfa, turf</p>	<p>European skipper, rangeland caterpillar, fall armyworm, sod webworm alfalfa caterpillar, loop r</p>	<p>2.5-4</p>
---	--	--------------

- NOTE:**
- For heavy infestations and mature worms use up to 4 fl. oz. per acre (1 gallon per 32 acres).

<p>Orchards, fruit trees, forestland timber, pecan, walnut, almond, apple, orange, shade trees, spruce, grapes, berries, strawberries, pome, stone fruit, tropical fruits</p>	<p>Fall webworm, redhumped caterpillar, tent caterpillar, leaf roller, orangedog, fruit-tree leafroller, eastern spruce budworm, gypsy moth, bagworm, cankerworm, fruitworm, hornworm, leaffolder, saltmarsh caterpillar, skeletonizer, tuffed apple bedmoth, walnut caterpillar</p>	<p>2.5-4</p>
--	--	--------------



6 9 5

- NOTE:**
- For heavy infestations and mature worms use up to 4 fl. oz. per acre (1 gallon per 32 acres).
 - For hand held sprayers used to treat vegetables use: 1/2 fl. oz. or approximately 15 milliliters of Delta BT in a gallon of water.
 - Trees should be sprayed using high volumes of water to provide uniform coverage.
-

Stored products grains, legumes, peanuts, oilseeds, other seeds	Indian mealmoth, almond moth, mediterranean flour moth	21 fl. oz. per 100 gallon
---	--	---------------------------------

- NOTE:**
- For heavy infestations and mature worms use up to 4 fl. oz. per acre (1 gallon per 32 acres).
-

WARRANTY

Delta Biological Products, Inc. makes no warranty, expressed or implied, concerning this material except those which are contained on this label.

