

DEC 1 1962

43679-5

EPA Reg. No. 43679-~~X~~

A Concentrated PYRENONE® Micro Emulsion

Effective, Fast Acting Insecticide

For Use in Homes, Institutions, Restaurants.

Food Processing Plants, Dairies, Livestock and Outdoors for Mosquito Control

Apply with Conventional Mechanical or Compressed Air Equipment

ACTIVE INGREDIENTS:

Pyrethrins	0.1%
Piperonyl Butoxide Technical	1.0%
Petroleum Distillate	0.4%
INERT INGREDIENTS	98.5%
	100.0%

*Equivalent to 0.8% (butylcarbonyl) 6-propylpiperonyl ether and 0.2% related compounds.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See side panel for additional cautions.

DO NOT TRANSPORT OR STORE UNDER 32° F.

PYRENONE — Registered trademark of Fairfield American Corporation

NET CONTENTS _____

INDUSTRIAL SPECIALTIES INC.

BOX 1113

TOPEKA, KANSAS 66601

DIRECTIONS FOR USE

violation of Federal law in that the document is a matter of confidence and is being

INDOORS APPLICATIONS

The use of live product in food processing establishments should be confined to the use of the product as an ingredient in food. The use of live product in food processing establishments should be confined to the use of the product as an ingredient in food. The use of live product in food processing establishments should be confined to the use of the product as an ingredient in food.

For maximum effectiveness, a combination of 1704 & 1705 is recommended in a logging or reclamation area.

[illegible]

To control carpet beetles: Spray edges of carpeting and under carpeting and rug. Make an extra application to floor and baseboards. Spray quickly and protect children and infants from inhaling. Repeat treatment as needed.

To control moldings: Spray moldings lightly with water around hole and seams. Tear ends open and spray in all joints. Trim baseboards, moldings and doors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray affected areas per label, testing a small nearby cracks and crevices along and behind baseboards, stairways, window and door sills, and entire areas of floor and base casing. Such handling should be placed in store at 180°F. for 10 hours following treatment. Concurrent treatment of animals with an approved topical or injectable insecticide is advised.

[illegible]

SPACE SPRAYING: Closes doors and windows and shut off venting systems. Acts as a gas-tight machine adjusted to deliver a hot mist at long (up to 100 ft) range of flame. Heat flux, photo-microscopes, ultrasonic film, mass, harmful substances from flammable hydrocarbons. Heat, motion, infrared heat, motion, and infrared motion. Direct spray of an infrared single-acting spray of infrared, throughout the entire area of a room of 200 sq ft. 1000 sq ft. 1000 sq ft. 1000 sq ft. 1000 sq ft.

POGGING. To an exposed or accessible stage of the above named clearing which is -- be a tree or one gallon to 20,000 to 40,000 cubic feet of space. Contact direct to direct, or even to pass the

[illegible]

To control blood-eating ticks, apply to the selected areas of the animal body a tick spray or tick dip. Repeat every 2 to 3 weeks if required.

OUTDOOR APPLICATION

ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. But best results treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery, or vegetation where mosquitoes may roost in shady places. Slightest fog or drizzle will help as a carrier of 1 gallon per acre (4" drizzle per 1000 square feet).

Using truck-mounted conventional mapping equipment such as Ruland's® models 41C, 51C or 64C traveling at 5 m.p.h. to load a 200 type punch catalog equipment to collect & gather on small STORAGE AND DISPOSAL.

Do not touch empty container. Wear container and put in waste collection

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS
Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritating persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is under way. After spraying in food processing plants, wash with potable water all equipment, benches, shelving, etc., where exposed food will be handled. All food processing surfaces and tunnels should be exposed during treatment or thoroughly washed before use.

Remove tanks and birds, and remove fish enclosures before spraying.

STANDARD FORM NO. 64

This product is toxic to fish. DO NOT APPLY DIRECTLY TO MARINE. Do not contaminate water by cleaning of equipment or disposal of wastes.

Buyer assumes all risks of use, storage or handling of this product and its contents in accordance with directions given hereon.

BATCH

BEST DOCUMENT AVAILABLE