

PRECAUTIONARY STATEMENT ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product as specified on this label.

STORAGE AND DISPOSAL

PROHIBITIONS

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL. OPEN DUMPING IS PROHIBITED. DO NOT REUSE EMPTY CONTAINER.

PESTICIDE DISPOSAL

PESTICIDE SPRAY MIXTURE OR PREPARATION THAT CANNOT BE USED OR CHEMICALLY REPROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL

TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE IN AN INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS OR BURY IN A SAFE PLACE.

GENERAL

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES SUCH AS LIMITED OPEN BURNING.

GENERAL DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Shu-Fly's Devil Dust is a ready-to-use insecticide dust which can be applied through the use of self-treating dust bags or by hand. But do not apply with bare hand. Use dust mitt or other protective glove.

Devil Dust For Livestock and Poultry with RABON® Pesticide Insecticide

Patent Pending

A ready-to-use insecticide dust. Aids in control of face flies on horses and beef and dairy cattle. Controls lice on swine. Controls northern fowl mites, chicken mites and lice on or around poultry. Controls ticks and fleas on dogs and cats.

ACTIVE INGREDIENT

2-chloro-1-(2,4,5-trichlorophenyl)

vinyl dimethyl phosphate

INERT INGREDIENTS

BY WEIGHT

3.00%

97.00%

TOTAL

100.00%

Net Weight: 4 lbs. (1.8Kg)

KEEP OUT OF REACH OF CHILDREN

CAUTION:

HAZARDS TO HUMANS

Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes. If the irritation persists, see a physician. Wash thoroughly with soap and water, after handling and before eating or smoking. Avoid contamination of feed and food-stuffs.

Do not apply in dwellings.

Manufactured by Shu-Fly Industries - Animal Health Division, Reno, Nevada
EPA REG. 43688-1

TOLL FREE
(800)648-5600

ACCEPTED
NOV 6 1980
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under
EPA Reg. No. 43688-1

DIRECTIONS FOR USE:

BEEF CATTLE AND DAIRY CATTLE

for the control of house flies and lice and to aid in the control of face flies.

HAND DUSTING - Apply approximately 2 oz. of dust by shake can, rotary duster or by spoon to the upper portions of the back, neck and poll and to the face as an aid in the control of face flies. Rub lightly to carry the dust beneath the hair. Repeat as necessary.

SELF-TREATING DUST BAG-FORCED USE

Put RABON® Livestock Dust in cotton cloth or double burlap bags or use pre-packed weather-proof cattle dust bags and hang in barn door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap from weather. **FREE CHOICE USE** - Use the same dust bags as above but place in feeding sheds, holding pens, feeders near water holes or other areas where cattle gather. The free choice aids in the control of lice.

POULTRY

for the control of

SWINE

for the control of lice. **HAND DUSTING** - Apply 3-4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary, but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of 3% dust should be applied per 150 sq. ft. of bedding.

HORSES

for aid in the control of face flies, for the control of stable flies, house flies, horn flies, horse flies, deer flies, lice, ticks, gnats and fly larvae. **HAND DUSTING** - Apply approximately 2 oz. of Devil Dust by shake can, hand duster, grooming brush, or dust mitt. Cover upper portions of back and neck and to the face, mane and tail for added control of face flies, rub or brush in lightly to carry dust beneath the hair. Repeat as necessary. **PREMISE**

TYPE HOUSING	APPLICATION	REMARKS
Stable	Hand dusting	For individual treatment of individual animals. Group treatment of entire area. Dust should be applied to the entire area every 14 days.
Stable	Free choice	For individual treatment of individual animals. Dust should be applied to the entire area every 14 days.
Stable	Free choice	For individual treatment of individual animals. Dust should be applied to the entire area every 14 days.
Stable	Free choice	For individual treatment of individual animals. Dust should be applied to the entire area every 14 days.
Stable	Free choice	For individual treatment of individual animals. Dust should be applied to the entire area every 14 days.
Stable	Free choice	For individual treatment of individual animals. Dust should be applied to the entire area every 14 days.
Stable	Free choice	For individual treatment of individual animals. Dust should be applied to the entire area every 14 days.
Stable	Free choice	For individual treatment of individual animals. Dust should be applied to the entire area every 14 days.
Stable	Free choice	For individual treatment of individual animals. Dust should be applied to the entire area every 14 days.
Stable	Free choice	For individual treatment of individual animals. Dust should be applied to the entire area every 14 days.

WARRANTY STATEMENT

SHU-FLY INDUSTRIES MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product. The user of this product assumes all liability for any damage or loss resulting from the use of this product.

RABON® is a trademark of Shu-Fly Industries, Inc. P.O. Box 10631, Reno, NV 89510

