

PRECAUTIONARY STATEMENT ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product as specified on this label.

STORAGE AND DISPOSAL

PROHIBITIONS
DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL
PESTICIDE DISPOSAL
PESTICIDE THAT CANNOT BE USED ACCORDING TO LABEL INSTRUCTION, MUST BE DISPOSED OF ACCORDING TO FEDERAL, STATE OR LOCAL PROCEDURES UNDER THE RESOURCE CONSERVATION AND RECOVERY ACT
CONTAINER DISPOSAL
TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE OF IN A SANITARY LAND FILL OR BY INCINERATION IF PERMITTED BY STATE OR LOCAL AUTHORITIES

GENERAL DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Ag Products' Devil Dust is a ready-to-use insecticide dust which can be applied through the use of self-treating dust bags or by hand. But do not apply with bare hand. Use dust mitt or other protective glove.

DEVIL DUST

FOR LIVESTOCK AND POULTRY

with RABON®
Pesticide Insecticide

Patent Pending

A ready-to-use insecticide dust. Aids in control of face flies on horses and beef and dairy cattle. Controls lice on swine. Controls northern fowl mites, chicken mites and lice on or around poultry. Controls ticks and fleas on dogs and cats.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate.....	3.00%
INERT INGREDIENTS.....	97.00%
TOTAL	100.00%

Net Weight **5LBS. (2.2KG)**

KEEP OUT OF REACH OF CHILDREN CAUTION: HAZARDS TO HUMANS

Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes. If the irritation persists, see a physician. Wash thoroughly with soap and water, after handling and before eating or smoking. Avoid contamination of feed and foodstuffs.
Do not apply in dwellings.

Manufactured by Ag Products, Inc., 1000 North Division Street, St. Paul, MN 55101
EPA REG. # 416-01 EPA EST. 43668-NV-01

DIRECTIONS FOR USE:

BEEF CATTLE AND DAIRY CATTLE for the control of horn flies, and lice, and to aid in the control of face flies.

HAND DUSTING—Apply approximately 2 oz. of dust by shaker can, rotary duster or by spoon to the upper portions of the back, neck and poll, and to the face as an aid in the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary. **SELF-TREATING DUST BAG-FORCED USE**—Put RABON® Livestock Dust in cotton cloth or double burlap bags or use pre-packed weather-proof cattle dust bags and hang in barn door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap from weather. **FREE CHOICE USE**—Use the same dust bags as above but place in feeding sheds, holding pens, feedlots, near water holes or other areas where cattle gather. The free choice aids in the control of lice.

POULTRY for the control of northern fowl mites, chicken mites, and lice. Dangling treated corn (bottle).

SWINE for the control of lice. **HAND DUSTING**—Apply 3-4 oz. dust by conventional hand or rotary duster to each animal with special attention given to the neck and around the ears. Repeat as necessary, but not more often than every 14 days. In severe infestations, both individual animals and barns may be treated. One lb. of 3% should be applied per 150 sq. ft. of barning.

HORSES for aid in the control of face flies, for the control of stable flies, house flies, horn flies, flies, deer flies, lice, ticks, gnats, fly larvae. **HAND DUSTING**—approximately 2 oz. of Devil Dust by shaker can, hand duster, or using brush, or dust mitt. Cover portions of back and neck. For added control of face flies, also the face, mane and tail. Rub or brush in lightly to carry dust beneath the hair. Repeat as necessary. **PREMISE DUSTING**—Apply to barn or stall area for control fly larvae and flies in stable area.

TYPE HOUSING	APPLICATION	
Wire Cages	1 lb./300 birds	For individual vent and fluff. may be preferred. Do not treat every 14 days.
	Plunger rotary type duster or shaker-can duster	Wire rungs are treated.
Notes: For northern fowl mites on roosters, thorough individual application, long-lasting control and reduce reinfestation of birds.		
FLOOR MANAGEMENT		
Litter	1 lb./100 sq. ft. Plunger or rotary type duster	Treat evenly. Roosts, cracks may hide.
Dust Box	2 lb./100 birds	Use box above.
Roosts and	1 lb./100 ft. line the A side of the roost with 1 lb. of dust with 1 qt. of water. Continually stir.	Treat roosts, cracks and

WARRANTY STATEMENT

SHU-FLY MAKES NO WARRANTY OF MERCHANTABILITY OR OTHERWISE, EXPRESS OR IMPLIED OF THE PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE USE OF THE PRODUCT OR ITS USE, AND WITH THE STATEMENTS MADE ON THIS LABEL.

Ag Products, Inc., St. Paul, MN 55101

DEVIL DUST

FOR LIVESTOCK AND POULTRY

with RABON®
Pesticide Insecticide

Patent Pending

A ready-to-use insecticide dust. Aids in control of face flies on horses and beef and dairy cattle. Controls lice on swine. Controls northern fowl mites, chicken mites and lice on or around poultry. Controls ticks and fleas on dogs and cats.

ACTIVE INGREDIENT	BY WEIGHT
chloro-1-(2,4,5-trichlorophenyl)	
diethyl phosphite	3.00%
INERT INGREDIENTS	97.00%
TOTAL	100.00%

Net Weight 5LBS. (2.2KG)

KEEP OUT OF REACH OF CHILDREN
CAUTION:
HAZARDS TO HUMANS

Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes. If the irritation persists, see a physician. Wash thoroughly with soap and water, after handling and before eating or smoking. Avoid contamination of feed and foodstuffs.
Do not apply in dwellings.

Manufactured by Shu-Fly Industries, A Division of Shu-Fly Corporation
EPA EST. 43666-NV-01

DIRECTIONS FOR USE:

BEEF CATTLE AND DAIRY CATTLE for the control of horn flies and eye gnats and to aid in the control of face flies.

HAND DUSTING—Apply approximately 2 oz. of dust by shaker can, rotary duster or by spoon to the upper portions of the back, neck and poll, and to the face as an aid in the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary. **SELF-TREATING DUST BAG-FORCED USE**—Put RABON® Livestock Dust in cotton cloth or double burlap bags or use pre-packed weather-proof cattle dust bags and hang in barn door exits or alleyways leading from animal buildings, stall or mineral blocks, or watering holes. Protect cloth or burlap from weather. **PREMISE DUSTING**—Use the same dust bags as above but place in rotating pens, holding pens, feedlots, near water holes, or other areas where cattle gather. The free choice aids in the control of lice.

POULTRY for the control of northern fowl mites, chicken mites, and lice. Drenching beetle (other pests).

SWINE for the control of lice. **HAND DUSTING**—Apply 3-4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary, but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of 3% dust should be applied per 150 sq. ft. of bedding.

HORSES for aid in the control of face flies, for the control of stable flies, house flies, horn flies, horse flies, deer flies, lice, ticks, gnats and fly larvae. **HAND DUSTING**—Apply approximately 2 oz. of Devil Dust by shaker can, hand duster, grooming brush, or dust mitt. Cover upper portions of back and neck. For added control of face flies, apply to the face, mane and tail. Rub or brush in lightly to carry dust beneath the hair. Repeat as necessary. **PREMISE DUSTING**—Apply to barn or stall area floors where manure accumulates to control fly larvae and flies in the stable area.

FOR CONTROL OF TICKS AND FLEAS ON CATS AND DOGS: Dust powder liberally over the animal and rub thoroughly through the hair coat to skin. Apply also to feet, between the toes and on the legs for ticks. Fleas will die quickly and can be combed from hair in a few hours if desired. Ticks are somewhat more difficult to kill and can be combed out of pet's hair a day or two after the application of this product.

FOR TICKS AND FLEAS IN KENNELS AND DOG HOUSES: Occasional dusting in and around sleeping quarters and other areas will keep free area of ticks and fleas. Dust bedding as well.

TYPE HOUSING	APPLICATION	REMARKS
Wire Cages	1 lb./500 birds	For individual treatment direct dust to vent and foot area. Group treatment may be preferred. Dust should not be in skin. Do not repeat more often than every 14 days.
	Rotary type dust or shaker can	Wire cages and corners should be treated.
Note: For northern fowl mites or lice, a thorough individual application of the dust will assure the mites or lice and reduce reinfestation of breeding flocks.		
FLOOR MANAGEMENT		
Litter	1 lb./100 sq. ft. of litter	Treat evenly and thoroughly. Also treat roosts, cracks and crevices where pests may hide.
Dust box	2 lb./100 sq. ft.	Use box about 2" x 2 ft. by 1 ft. deep.
Water trough	1 lb./100 sq. ft.	Treat evenly and thoroughly. Also treat cracks and crevices.

WARRANTY STATEMENT

SHU-FLY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions and in accord with the statement of directions on the label.