

Sanitizing of Food Processing Equipment and Contact Surfaces in Food Contact Locations

For sanitizing food processing equipment, contact surfaces, food utensils, dishes, silverware, glasses, cooking utensils, tops, refrigerated storage containers, equipment, and other surfaces.

Wash and rinse in hot, soapy water. Then apply 2 to 4 oz. **Sana Cage Wash** in 4 gallons of water (200 ppm) to the surfaces. Surfaces should remain wet for at least 10 minutes. Wash, adequate draining and air drying. For 2 to 4 gallons of prepared daily use solution, use mechanical application or mechanical application. Use solution may not be reused for sanitizing applications.

Apply to sink tops, counter tops, refrigerated storage and other equipment and other stationary hard surfaces. For dishes, glasses, silverware, glasses, cooking utensils and other small size food processing equipment can be sanitized by immersion in a 2 to 4 gallon solution of **Sana Cage Wash**.

No Potable water rinse is required.

At 2 to 4 gallons **Sana Cage Wash** has a hard water tolerance of 750PPM of hardness (calculated as CaCO₃) when measured by the AOAC Germicidal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*.

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 2 to 4 oz. **Sana Cage Wash** in 4 gallons of warm water.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towels back into sanitizing solution. When solution becomes dirty, change and provide fresh solution.

Precautionary Statements

Hazards to Humans and domestic animals

DANGER

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

OPEN DUMPING IS PROHIBITED.
DO NOT REUSE EMPTY CONTAINER.

PESTICIDE DISPOSAL

PESTICIDE OR RINSATE THAT CANNOT BE USED OR CHEMICALLY REPROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES, OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL

TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE IN AN INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS OR BURY IN A SAFE PLACE.

GENERAL

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES SUCH AS LIMITED OPEN BURNING.

SANA CAGE WASH

**DISINFECTANT-SANITIZER
FUNGICIDE DEODORIZER**

Disinfectant-Sanitizer-Fungicide-Deodorizer for Hospitals and Tuberculosis, Industrial, School, Dairy and Other, Earth and Marine Use.

Active Ingredients

Alkyl(C₁₂, C₁₁, C₁₀, and C₉) (as C₁₀) 5.0% 2.5%

Dimethyl Benzyl Ammonium Chloride

Alkyl(C₁₂, C₁₁, C₁₀, and C₉) 2.5%

Dimethyl Ethylbenzyl Ammonium Chloride

Inert Ingredients

95.0%

100.0%

KEEP OUT OF REACH OF CHILDREN.

DANGER

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No. 43736-1

Net Contents 5 GAL.

Manufactured By:

See Corporation, Milltown, New Jersey 08850

Directions for Use

GENERAL CLASSIFICATION

Disinfectant-Sanitizer-Fungicide-Deodorizer for Hospitals and Tuberculosis, Industrial, School, Dairy and Other, Earth and Marine Use.

Sana Cage Wash will disinfect and deodorize surfaces. When applied with a brush, it is especially effective on surfaces. Surfaces may be sprayed with the spray, washed, treated, and surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the solution becomes dirty.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution.