

EASTERN MINERALS'
FLY CONTROL MINERAL
With Rabon® Oral Larvicide

DM 15

To prevent the development of horn flies, face flies, house flies, and stable flies in the manure of treated cattle.

43757-1

ACTIVE INGREDIENT

Tetrachlorvinphos: 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphite

BY WEIGHT

1.00%*
99.00%**
100.00%


100%

INERT INGREDIENT

TOTAL

*Each pound contains 4.54 g of Rabon
**Refers only to ingredients which are not larvicidal

READ ENTIRE LABEL FOR DIRECTIONS
AND PRECAUTIONARY STATEMENTS



GUARANTEED ANALYSIS

Calcium (Ca) Min.....	20.00%
Calcium (Ca) Max.....	23.00%
Phosphorus (P) Min.....	6.00%
Salt (NaCl) Min.....	23.00%
Salt (NaCl) Max.....	25.00%
Iron (Fe) Min.....	1.00%
Manganese (Mn) Min.....	0.20%
Zinc (Zn) Min.....	0.40%
Copper (Cu) Min.....	0.05%
Cobalt (Co) Min.....	0.005%
Iodine (I) Min.....	0.005%
Fluorine (F) Max.....	0.06%
Vitamin A I.U.'s per Lb., Min.....	200,000
Vitamin D-3 I.U.'s per Lb., Min.....	20,000

FEED INGREDIENTS

Defluorinated Phosphate, Calcium Carbonate, Salt, Cottonseed Meal, Cane Molasses, Iron Oxide, Ferrous Carbonate, Manganous Oxide, Zinc Oxide, Copper Oxide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, and Anise Oil.

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE BACK PANEL FOR FEEDING DIRECTIONS

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear long-sleeved shirt and pants; chemical resistant gloves; shoes and socks for protection when handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of waste.

Do not use manure/droppings on land where food crops are to be grown during the current growing season.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Store in a dry place in the original container.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty into mixing equipment by shaking and tapping sides and bottom to loosen clinging particles. Then dispose of bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE:

7 of 2

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Feed EASTERN MINERALS' FLY CONTROL MINERAL WITH RABON® ORAL LARVICIDE at the rate of 2.4 ounces per 1000 pounds of body weight per head per day by mixing it into the ground grain or complete daily ration.

This product will mix uniformly in feeds when standard mixing procedures are followed. Thus, usual problems that are common to all feed preparation and which cause stratification such as excessive free-fall or excessive handling are to be avoided. It is recommended that appropriate preblending techniques be employed to assure adequate distribution throughout the feed mix. The premix should be preblended with ground grain, protein supplements, mineral mixes, etc., before being added to roughages, such as chopped hay or silage. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if this product is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e. vertical mixers, horizontal blenders, mixer/feeder truck) may be used to prepare formulated feeds.

Feeds prepared with this product should not be pelleted, nor be mixed with feeds containing predominantly pellets. Further, this product should not be mixed in liquid feed supplements.

Start feeding EASTERN MINERALS' FLY CONTROL MINERAL WITH RABON® ORAL LARVICIDE early in the spring before flies begin to appear and continue feeding throughout the summer into the fall until cold weather restricts fly activity.

EASTERN MINERALS' FLY CONTROL MINERAL WITH RABON® ORAL LARVICIDE MAY BE FED UP TO SLAUGHTER AND TO LACTATING DAIRY COWS.

EASTERN MINERALS' FLY CONTROL MINERAL WITH RABON® ORAL LARVICIDE PREVENTS THE DEVELOPMENT OF HORN FLIES, FACE FLIES, HOUSE FLIES, AND STABLE FLIES IN THE MANURE OF TREATED CATTLE BUT IS NOT EFFECTIVE AGAINST EXISTING ADULT FLIES. When starting a feed program during the fly season, it is desirable to use other control measures to reduce the population of existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, EASTERN MINERALS' FLY CONTROL MINERAL WITH RABON® ORAL LARVICIDE should be used in conjunction with other good management and sanitation practices.