# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS) CAUTION

Read the entire label before using. Use only as recommended for claims listed

During commercial exposure in seed treatment and loading operations, wear goggles clean synthetic rubber goves protective clothing, and a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR part II

Wash hands, arms, and face thoroughly with soap and water after handling and before eating or smoking. Houtinely shower or bathe after work and wash all clothing with soap and hot water before reusing.

### **ENVIRONMENTAL HAZARDS:**

Thir fuct is toxic to fish. Keep out of any body of water. Do not contaminate water cleaning of equipment or disposal of wastes. Bury wastes and empty containers in isolated area away from water supplies.

## DIRECTIONS FOR USE AND RECOMMENDATIONS:

If is a violation of State and Federal Law to use this product in a manner inconsistent with its labeling.

Flead all directions before using and use only as specified on this label

DIRECTIONS Thoroughly mix the recommended amount of Captan-Methoxyclor 300-20 Flowable Seed Protectant into the required amount of water for the slurry treater equipment and dilution rate to be used. Clean equipment thoroughly between treating operations.

#### SPECIAL USE PRECAUTIONS:

IMPORTANT: Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled

## TREATED SEED DO NOT USE FOR FEED, FOOD, OR OIL PURPOSES

THE Federal Government and some states have issued regulations for the application of special tags or labols to be affixed to bags containing treated seed. Dispose of excess treated seed by burial.

#### PRODUCT INFORMATION:

Captan-Methoxyclor 300-20 Flowable Seed Protectant is a liquid suspension that is especially formulated for the treatment of seed prior to storage and planting to protect seed from molds and other fungi causing storage loss and to protect seed from seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights. Also for protection of stored seed against Cade'le, confused flour beetle, flat grain beetle, granary weevil, lesser grain borer, red flour beetle, rice weevil, and saw-toothed beetle. At the recommended dosage Captan-Methoxyclor 300-20 Flowable Seed Protectant is noningunous to seeds or seedlings and the germination of treated seed is frequently stimulated, giving a superior result as compared to results from untreated seed.



CAPTAN - METHOX VCHLOR 300-20 FLOWABLE SEED PROTECTANT

Agricultural Fungicide - ACTIVE INGREDIENTS:

- Liquid Suspension BY WT.

CAPTAN: N-[(Trich'oromethyl) thio]-4-Cyclohexene-1, 2-dicarboximide

\*\*Boharythan, Technicar\*

INERT INGREDIENTS

30.0% 2.0% 68.0% 100.0%

\*Equivalent to 1,76% of 2, 2 is ignifiction; phenylin 1, 1, 1, thic libit ename and 0,24% other isomers, and related noncounds.

# CAUTION: KEEP OUT OF REACH OF CHILDREN

\*IN CASE OF EYE CONTACT: Immediately flush eyes with running water for at least 15 minutes. Call a Physician.

\*IN CASE OF SKIN CONTACT: Wash the affected area with soap and warm water.

\*Causes eye and skin irritation.

\*Harmful if swallowed, inhaled, or absorbed through skin.

\*Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling.

\*Avoid inhaling of spray mist.

\*Do not use or store near food or feed. See side panel for additional precautionary statements.

STORE IN A COOL, DRY PLACE.
STIR THOROUGHLY TO MIX CONTENTS
BEFORE USING.

5 GAL. NET

AGRICULTURAL PRODUCTS COMPANY, Harlingen, Texas 78550

The Following seeds in **OU** 

Alfalfa Clossians of Beets (garden) Cereal Grains (I Barley Iye Oats Wheat Cole Crops (E. 2)

Corn Field Corn Sweet Cor Cotton Acid delinte:

Reginned a of Cruchiers (Mc. st. Flax Grasses Milo Muskmelon Via Peanuts (unsn-ii Peas Rice Sorotium chi. st. Flax Grantium chi. st. flax

Rice Sorghum (hu Soybeans Spinach Sugar Beets, Wat £ s

Sunflowers Swiss Chard Tomato Cucu - a

PROMIBITIONS
ed --Dr no. -u
PESTICIDE DIS

2 PESTICIDE DI Shou'd te disc si 3 CONTAINER D

gesticide compre GENERAL: Et limited open burn

CONDITIONS OF S buser and a lusers ( ) a by wither agreem WARRANTY LIW T GESCRIPTOR OF THE ; a DE INTERENTIALISM FORM

WARRANTY OF MI nerect WHERENT RISKS. HOMERENT CLISHT; HOWERENT CLISHT APP VALCE OF THE LIMITATION OF ALL CONSEQUENTIAL DAYS OF ALL CONSEQUENTIAL DAYS OF ALL

EPA Reg 1.3 EPA Est. 4.7

amount than the co



## Captam - Methoxychlor 366-26 FIGHABLE SEED PROTECTANT

Agricultural Fungicide — Liquid Suspension ACTIVE INGREDIENTS: CAPTAN: N-((Trichloromethyl) thio)-4-Cyclohexene-1, 2-dicarboximide 30.0%

Millionrychiar, Technical

2.0% INERT INGREDIENTS 2.0%

100.0% rEquivarent to 1.76% of 2, 2 bis in-methoxyphenyth 1, 1, 1-trichlorustrate and 0,24% other isomers and related compounds.

# CAUTION: KEEP OUT OF REACH OF CHILDREN

- 'IN CASE OF EYE CONTACT: Immediately flush eyes with running water for at least 15 minutes. Call a Physician.
- 'IN CASE OF SKIN CONTACT: Wash the affected area with soap and warm water.
- \*Causes eye and skin irritation.

Ø \*

\* \* \* \* \* \* \* \*

- \*Harmful if swallowed, inhaled, or absorbed through
- \*Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling.
- \*Avoid inhaling of spray mist.
- \*Do not use or store near food or teed.

See side panel for additional precautionary statements.

STORE IN A COOL, DRY PLACE. STIR THOROUGHLY TO MIX CONTENTS BEFORE USING.

5 GAL. NET

AGRICULTURAL PRODUCTS COMPANY, Harlingen, Texas 78550

SEED TREATMENT DOSAGE RATES

Ozore 190 is The Following Table gives the amount of this property in seeds in OUNCES per HUNDRED POUNDS of seed SEED

Allalia Crover Lespedeza Treto	:DOSAGE RATE
Beans is national conseas.  Beets igardeni	2 2 2
Cereal Grains (for seed & seeding rotuling) for for bent smo	
Oats Wineat	2t ce 15 -
Cole Crops (Broccoli Brussels sprouts Cabbage Caustioner	1 1 2 2 2 2
Corn	1 1 67 150
Field Corn	
Sweet Corn	さんしょ だんこうがん
Cotton	ន អំ ធ្វី ស៊ីន
Acia definted cotton	
Reginned and Fuzzy cotton	2 21 52 155 5
Crucifiers (Mustard Radishes Rape Turnips	37:11 62 150 2
161	1 21 02 15 5
Grasses	3 21: 62 155:15
Milito	6 ptr ag 106 a
Muskmelon Watermelon Pumpuin Squash	5# 62 100 is 14f 62 100 s
Peanuts runsnehed. Peas	14 fr oz 100 ± 6fr oz 100 ±
Rice	2':1 02 100 t
Sorghum (hull-less	31 62 109 2
Soybeans	3'21 02 105.E
Spinach	21/ oz 100 is
Sugar Beets West	51 oz 100 s
East '	5 f oz 100 a
Sunflowers	10 fr oz 100 6
Swiss Chard	3th oz 100 5
Tomato Cucumbers Cantaloupe	101/02 100/5
	2 . 1 52 1.0 %

## STORAGE AND DISPOSAL:

- 1 PROMBITIONS. —De not comum nate water food ler feed by storage or dispose. —— den dumb no equino n
- PROMISETIONS. Do not company have when how on source and source of the post GENERAL: Consult Federa. State of 1000 dispose authorizes for approved aremative procedures such as

## NOTICE - READ CAREFULLY

NOTICE — READ CAREFULLY

CONDITIONS OF SALE. Application Products Company, and select over a majorical flow service of the conditions of selection flows according to the selection of conditions of selection flows according to the selection of conditions of selections of the conditions of the conditi

58 EPA Reg. No. 43789-2 EPA Est. 43789-TX-1