# PRECAUTIONARY STATEMENTS 'S HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

. .

auses irreversible eye danage. Maraful if swallowed or inhales, May sume sliergic skin reactions. On not get in eyes, Year goggles or some shield when handling. Not sould contact with skin and clothing. Not see and separately launder clothing before resse.

issh hands, arms and face thoroughly with soap and water after handling and before eating, drinking, using tobacco products or applying commetics, butingly shower or bathe after working. Do not contaminate food and feedatuffs. This product must be applied in a well ventilated area.

### ENVIRONMENTAL HAZARDS: .

This posticide is toxic to fish. Reep out of any body of water, lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Exposed treated seed may be hazardous to birds and other wildlife. Dispose of excess treated seeds by burial away from bodies of water.

#### ENDANGERED SPECIES PRECAUTIONS

The use of this product in a manner which may harm endangered species or adversely modify their habitat is prohibited by Federal law. The following endangered species and the general location of their habitats have been identified by county:

Slackwater Darter, Alabama (Lauderdale, Limestone and Hadison)
Tennessee (Lawrence and Wayne)

Pecos gambusia,

New Hexico (Chaves and Eddy) Texas (Reeves, Jeff Davis and Pecos)

Preshwater Mussels, Tennessee (Mancock, Claiborne, Hawkine, Sullivan)
virginia (Smyth, Scott, Lee, Washington, Grayson)

Commanche Springs Funfish.

Texas (Reeves and Jeff Davis)

Prior to use of this product in the above counties you must determine that these species are not located in, or immediately adjacent to, the areas to be treated with captan. Consult your regional U.S. Fish and wildlife Service office (Endangered Species Specialist) or Statu Wildlife Agency If there is any doubt as to the areas where the outdoor use of captas is prohibited.

### DIRECTIONS FOR USE:

It is a violation of federal law to use this product to a manner inconsistant with its labeling.

#### Use Precaution Statements


Mixers, loaders, and applicators, when mixing, loading, and applying, must wear mid-forearm to elbow-length natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan; boots or overshoes; one-piece overalls which have long sleeves and long pants; face shield or goggles and a hat or other appropriate head covering.

A dust mask must be worn when mixing and loading this product.

A chemical-resistant apro $\ \$  must be worn when mixing and loading this product.

Clothing worn while loading, mixing and applying this product must be laundered separately from other clothing before reuse. Clothing that may have been drenched or heavily contaminated must be disposed of in accordance with State and local regulations.

Applicator's protection may also be obtained by use of an enclosed, tractor cab with a properly filtered air supply.



PM12 43789-5 192

### PRODUCT INFORMATION:

Captan' - Methoxychlor 65 - 10 WP Seed Protectant is a microfine wettable powder especially formulated for the treatment of seed prior to storage and planting to protect seed from molds and other fungi causing storage loss and to protect seed from seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights. At the recommended dosage Captan - Methoxychlor/65 - 10 WP Seed Protectant is noninjurious to seeds or seedlings and this gennination of treated seed is frequently stimulated, giving a superior result as compared to results from untreated seed.

# SEED TREATMENT DOSAGE RATES Oz. per 100 lb.

The Following Table gives the amount of this product to use on various seeds, in OUNCES per HUNDRED POUNDS of seed.

SEED	DOSAGE RATE
Alfalfa, Clover, Lespedeza, Trefoil	7 oz./100 (b.
Beans (snap, dry, cowpeas)	2% oz./100 lb
Beets (garden)	10 oz /5 0 lb
Cereal Grains (for seed & seedling rots - not for burnt s	muti
Barley, rye	2 oz /100 lb
Oats	2½ oz /100 lb.
Wheat.	14 oz /100 lb.
Cole Crops	
(Broccoli, Brussels sprouts, Cabbage, Cauliflower)	14 oz (100 lb.
Com	
Field Corn	1% oz /100 Ib
Sweet Corn	24 or (100 lb.
Cotton	
Acid delinted cotton	2 02 (100 15
Reginned and Fuzzy cotton	2 oz /100 lb
Crucifiers (Mustard, Radishes, Rape, Turnips)	17 02 / 20 16
Flax	2V or (100 lb.
Grasses	61/ a= (100 lp
Milo	
Muskmelon, Watermelon, Pumpkin, Squash	5 OZ./100 ID.
Peanuts (unshelled)	1% 02./100 lb.
Peas.	
Rice.	2 02./100 15.
Sprobum (hull-less)	3 oz./100 lb.
Sorghum (hull-less)	3% oz./100 lb.
Soybeans	2 oz./109 lb.
Spinach	5 oz./100 lb.
Sugar Beets, West	5 oz./100 lb.
East	10 oz./100 lb.
Sunflowers	3 oz./100 lb.
Swiss Chard	10 oz./100 lb.
Tomato, Cucumbers, Cantaloupe	2¼ oz./100 lb.

SEED TREATMENT: DO NOT USE TREATED SEED FOR POOD, FEED OR OIL PURPOSES.
Treated seed should be labelled in accord with all the applicable requirements of the Federal Seed Act. The Federal Seed Act requires that seed treated with a posticide must contain a dys which imparts an unnatural color to the seed if the seed is intended to be moved in interstate commerce. Labels attached to treated seed must state, "Treated seed must not be used for food, feed or oil purposes." and "Exposed treated seed may be hazardous to birds and other wildlife. Dispose of excess treated seed by burial away from bodies of mater."

# CAPTAN-METHOXYCHLOR 10 WP DIRECTIONS FOR SE:

It is a violation of State and Federal Law to use this product in a manner inconsistent with its labeling.

Read all directions before using and use only as specified on this label.

### STIR CONTENTS THOROUGHLY INTO WATER BEFORE USE.

FOR SLUBBY TREATMENT, apply in a slurry-type treater specifically designed and approved for that purpose or apply by hand in any suitable mixing equipment, such as home made rotary drums.

Thoroughly mix the recommended amount of this product into the required amount of water for the mlurry treater equipment and dilution rate to be used. Clean equipment thoroughly between treating operations.

Only seed of high Aermination with good vigor should be treated with this product. Poor seed may suffer greater injury. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Store treated seed in a dry place until planted so that excess moisture will evaporate. If seed is stacked when wet, heat may develop which could impair germination.

IMPORTANT - Be certain treated seeds are completely dry before sacking or storing.

STORE ALL TREATED SEED IN A COOL, DRY PLACE.

IMPORTANT: Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled;

### TREATED SEED

### DO NOT USE FOR FEED, FOOD, OR OIL PURPOSES

## STORAGE AND DISPOSAL

Do not contaminate water, look or look by startes or dispasal. Open dymping is prohibited.

\$TORAGE Project from excessive heat. Store in a cool (temperature below \$5°F/29°C), dry, well ventilated area away from flasmable materials and sources of heat or flame. Store away from foodstuffs or animal feed in an area designated for pesticides. Exercise due caution to prevent damage to or leakage from the container. Store in original container only. Keep container tightly closed when not in use.

PESTICIOE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticides, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINED MAPOSAL: Paper and plastic bags — Completely empty bag into application equipment. Then dispose of empty bag in a sanilary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### NOTICE-READ CAREFULLY

CONDITIONS OF SALE: Satisf(s) offer(s) this product for sale subject to, and buyer and all users are germed to have accepted. the following conditions of sale and warranty which may only be varied by written agreement of a duly authorized representative of Seller(5)

WARRANTY LIMITATION: Seller(s) warrant(s) that this produce conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the tabel subject to the inherantirisks referred to below. Seller(s) make(s) no other express warranges. THERE IS NO IMPLIED WARRAN-TY OF MERCHANTAINLITY and there are no warranties which extend beyond the description on the label

IMMERENT RISKS: The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with its use. Buyer assumes all risks associated ciated with use or application of this product contrary to label instructions or resulting from extraordinary wee-

LIMITATION OF LIABILITY: In no case shall Seller(s) be liable for special, indirect, or consequential demages resulting from the use or handling of this product and no claim of any kind shall be greater in amount than the purchase price of the product in respect of which such damages are claimed.

EPA REG. NO. 43789-5 EPA Est. No.



. .