Campata eve and skin irritation. Maraful if avallowed, tohated or absorbed through skin. Do not get in eyes, on skin, or on clothing, Mear goodles or face shield and subber gloves when handling. Avoid imbaling of spray mist or dust, Do not use or store near food or feed.

Dering commercial exposure in seed treatment and loading operations. wear goodles, clear subber gloves, protective clothing, and a pesticide respirator jointly approved by the Mine Bafety and Health administration (formerly the U.S. Bureau of Mines) and by the National Institute of Occupational Safety and Bealth under provisions of 10 CFR. pact 11.

Much hards, arms, and face thoroughly with most and water after handling and before eating or smoking. Routinely shower or bathe after works wash all clothing with soap and water before results.

Consumption of alcoholic beverages increases the toxic effect of +hiram

ENVIRONMENTAL HAZARDS

This pesticide product is toxic to fish, birds and wildlife. Do not contaminate water by cleaning of equipment or disposal of wastes, Drift or runoff from treated areas may be harardous to squatic organisms in neighboring areas. Do not apply directly to water or wetlands. Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all excess treated seeds by burial away from bodies of water.

DIRECTIONS FOR USE

It is a violation of State and Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal, Open dumping is prohibited. Do not seuse empty container.

STORAGE: Store in a cool, dry place, but do not allow to freeze. Store in an area designated for peaticides, and away from food or feed. Keep pesticide in original container. Keep container firmly closed when not in use.

PESTICIDE DISPOSAL: Pesticide vastes are toxic. Improper disposal of excess pesticide, spray mixture, or rineate is a violation of Yederal Law. If these wastes cannot be disposed of by use according to label instructions, contact your state Festicide or Environmental Control Agency, of the Mazardous Maste representative at the nearest 27A Regional Office for guidance.

CONTAINER DISPOSAL: (plastic) Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. (Metal) Triple rines (or equivalent). Then offer for recycling or

reconditioning, or puncture and dispose of in a sanitary landfill, or by other Procedures approved by state and local authorities.



THIRAM 300 DYED

FLOWABLE SEED PROTECTANT

ACTIVE INGREDIENTS: BY WT. INERT INGREDIENTS: 70.0% THIS PRODUCT CONTAINS THE TOXIC INERT INGREDIENT 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or two classes of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Call a Physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-tomouth.

IF ON SKIN: Wash the affected area with soan and Warm water.

IF IN EYES: Immediately flush eyes with running water for 15 minutes. Call a Physician

See side panel for additional drecautionary statements.

EPA Est. EPA Reg. No. 43789-35

5 GAL. NET

MEG'd. FORE AGRECULTURAL MIDUCTS CON 2 43789-35

* * * * * * * * * *

्रा प्रश्ना हरू।

43784-35

GENERAL INFORMATION
ommended for the treatment of listed seeds, and when us and it is directed, will usually and it. This prof increase 4 ... d years by reducing loss from feed decay, damping-off, and washing control smaller of sorghum.

DIRECTIONS FOR USE

AR A SEED TREATMENT: This product should be applied with water as a suspension in the slurry-type treater designed and approved for this purpose or may be applied by hand in any surjects flexing equipment such as home-made rotary drums. Seed may be treated anytime after being well cured.

FOR SLURRY TYPE TREATER: Use by mixing with water in a slurry treater at the concentration necessary for the type of slurry equipment being used and the seed being treated

FOR HAND SLURRY TREATMENT: Consult the following table for the number of fluid ounces recommended for each 100 pounds of seed. Add this amount to 1 pint of water for 100 pounds of seed. Stir to a smooth suspension. Place seed ma mixing container or spreading 4-inch layer on the floor on a clean dry cloth or paper. Sprinkle the measured amount of sharry succension over the specified amount of seed and mix immediately. Turn seed at least 5 times and allow to dry before planting

IMPORTANT: Treated seed must not be used for or mixed with food or animal feed or processed for oil Treat only those seeds needed for immediate use, finhimiting the interval between treatment and planting Do not store excess treated seeds beyond planting time. Seed commercially treated with this product must be labeled in accord with all applicable requirements of the Federal Seed Act. Labels attached to treated seed must state. Treated seed must not be used for or mixed with food or animal feed, or processed for oil." and "Exposed treated seed may be hazardous to bride and other wildlife. Dispose of all excess treated seed and seed packaging by bunal away from bodies of water

SEED TREATMENT DOSAGE RATES

Type of seed	Fluid ounces per 100 pound
Barley	
Seans (Snap & Dry	
Beans (Lima)	
Seets (Garden & Sugar)	
Cantaloupe, Cucumber, Pumpkin.	•
Squash, Watermelon	6
Cestor Beans	
Corn	
Cotton (Acid delinted)	
Cotton (Reginned & Fuzzy)	
Comoss	
Cowpess	
ggplant	
lax	
Grasses	11
Legumes (Small seeded)	
Millet	.,, 5
Okra, Oniona	B
Peanuts (Shelled)	6
Peas	
Rice	
Rye	
Safflower	
Carema	
Sesame	
Sorghum	
Soybeans	
Tomato	
Wheat	
Broccoli, Brussel Sprouts, Carrots, Cabbage, Caulif Kohlrabi, Lettuce, Mustard, Peppers, Radish, Spinac	lower, Collards, Endive, Kale,
FOR TREATMENT OF RICE SEED IN CALIFORNIA:	
Used according to directions, this product will usually increa	and along and sinide by radiusing love at fr
reed decay, demping-off, and seed no blights caused by see	

For treatment of nice seed in California, use this product at the rate of 5 0 ounces per 100 pounds of seed. It may be applied to well cured rice seed prior to pre-plant soaking. It should be used with water as a suspension in freaters suitable for adequate mising and accurate application to seed

HOTE: To prevent water containmeation in irrigation or drainage canals soak water should be retained in settling basins. Do not use treated seed for food or feed

NOTICE - READ CAREFULLY

CONDITIONS OF SALE: Seller(s) offer(s) this product for sale subject to, and buyer and all users are deemed to have accepted, the following conditions of sale and warranty which may only be varied by written agreement of a duly authorized representative of Seller(a).

WARRANTY LIMITATION: Seller makes no varianty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all tisk of use and/or handling of this material when such use and/ Or handling is contrary to label instructions.

LIMITATION OF LIABILITY: Seller's quarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of misuse or improper handling and accepts the product On these conditions.