

THIRAM 30% UNDYED FLOWABLE SEED PROTECTANT

ACTIVE INGREDIENT: Thiram (Tetramethylthiuram disulfide) **BY WT. 30.0%**
INERT INGREDIENTS: **70.0%**
100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

Statement of Practical Treatment

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit fluid is clear. Call a Physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

IF ON SKIN: Wash the affected area with soap and warm water.

IF IN EYES: Immediately flush eyes with running water for at least 15 minutes. Call a Physician.

See side panel for additional precautionary statements.

Mfg. by Agricultural Products Co.
Ft. Worth, Texas 76112

EPA Est. 43789-TX-01
EPA Reg. No. 43789-36

Net Contents

ACCEPTED
MAY 26 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 43789-36

GENERAL INFORMATION

This product is recommended for the treatment of listed seeds and, when used as directed, will increase stands and yields by reducing losses from seed decay, damping-off, and seedling blights. It also will help control the loose and curved kernels known as "mushy" kernels.

DIRECTIONS FOR USE

AS A SEED TREATMENT: This product should be applied with water as a suspension in the slurry-type treater designed and approved for this purpose, or may be applied by hand in any suitable mixing equipment such as home-made rotary drums. Seed may be treated anytime after being well cured.

FOR SLURRY TYPE TREATER: Use by mixing with water in a slurry treater at the concentration necessary for the type of slurry equipment being used and the seed being treated.

FOR HAND SLURRY TREATMENT: Consult the following table for the number of fluid ounces recommended for each 100 pounds of seed. Add this amount to 1 pint of water for 100 pounds of seed. Stir to a smooth suspension. Place seed in a mixing container or spread in a 4-inch layer on floor on a clean dry cloth or paper. Sprinkle the measured amount of slurry suspension over the specified amount of seed and mix immediately. Turn seed at least 5 times and allow to dry before planting.

IMPORTANT: Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Seed commercially treated with this product must be labeled in accord with all applicable requirements of the Federal Seed Act. Labels attached to treated seed must state: "Treated seed must not be used for or mixed with food or animal feed, or processed for oil," and "Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water."

NOTE TO USER

The purchaser of this product is responsible for insuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21 CFR Part 225. Any dye added to treated seed must be cleared for use under 40 CFR Part 180.1001.

SEED TREATMENT DOSAGE RATES

Type of Seed	Fluid ounces per 100 pounds
Barley	5.9
Beans (Snap & C y)	2.8
Beans (Lima)	4.2
Beets (Garden & Sugar)	11.2
Cantaloupe, Cucumber, Pumpkin, Squash, Watermelon	6.3
Castor Beans	6.3
Corn	4.2
Cotton (Acid deinted)	6.3
Cotton (Reginned & Fuzzy)	6.3
Cowpeas	2.8
Eggplant	8.4
Flax	7.4
Grasses	11.2
Legumes (Small seeded)	11.2
Millet	5.6
Oats, Onions	3.4
Peanuts (Shelled)	6.3
Peas	4.2
Rice	3.5
Rye	5.2
Safflower	5.6
Sesame	4.2
Sorghum	3.8
Soybeans	2.8
Tomato	8.4
Wheat	4.6

Broccoli, Brussels Sprouts, Carrots, Cabbage, Cauliflower, Collards, Endive, Kale, Kohlrabi, Lettuce, Mustard, Peppers, Radish, Spinach, Swiss Chard, Turnip All 11.2

FOR TREATMENT OF RICE SEED IN CALIFORNIA:

Used according to directions, this product will usually increase stands and yields by reducing losses from seed decay, damping-off, and seedling blights caused by seed-borne and soil-borne organisms.

For treatment of rice seed in California, use this product at the rate of 5.0 ounces per 100 pounds of seed. It may be applied to well cured rice seed prior to pre-plant soaking. It should be used with water as a suspension in treaters suitable for adequate mixing and accurate application to seed.

Note: To prevent water contamination in irrigation or drainage canals, soak water should be retained in settling basins. Do not use treated seed for food or feed.

NOTICE — READ CAREFULLY

CONDITIONS OF SALE: Seller(s) offer(s) this product for sale subject to, and buyer and all users are deemed to have accepted, the following conditions of sale and warranty which may only be varied by written agreement of a duly authorized representative of Seller(s).

WARRANTY LIMITATION: Seller(s) warrant(s) that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the label subject to the inherent risks referred to below. Seller(s) make(s) no other express warranties. THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY and there are no warranties which extend beyond the description on the label hereof.

INHERENT RISKS: The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with its use. Buyer assumes all risks associated with use or application of this product contrary to label instructions or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY: In no case shall Seller(s) be liable for special, indirect, or consequential damages resulting from the use or handling of this product and no claim of any kind shall be greater in amount than the purchase price of the product in respect of which such damages are claimed.