

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(AND DOMESTIC ANIMALS)
CAUTION**

Causes eye and skin irritation. Harmful if swallowed, inhaled, or absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling. Avoid inhaling of dust. Do not use or store near food or feed.

Wash hands, arms, and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work and wash all clothing with soap and hot water before reusing.

ENVIRONMENTAL HAZARDS


This product is toxic to fish, birds, and other wildlife. Do not apply directly to lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

- 1. PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.
- 2. PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State, or Local procedures under the Resource Conservation and Recovery Act.
- 3. CONTAINER DISPOSAL:** Dispose of liners according to approved Federal, State, or Local procedures under the Resource Conservation and Recovery Act. If container cannot be reused, dispose of in a sanitary landfill or by incineration if permitted by State and Local Authorities.
- 4. GENERAL:** Consult Federal, State, or Local Disposal Authorities for approved alternative procedures such as limited open burning.
- 5. STORAGE:** Store in a cool, dry place.



ZINEB 75% FUNGICIDE

Controls Diseases on Vegetables, Fruits & Ornamentals.

ACTIVE INGREDIENT:	BY WT.
Zinc Ethylene Bisothiolate	75.0%
INERT INGREDIENTS:	25.0%
	100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Statement of Practical Treatment

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit fluid is clear. Call a Physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

IF ON SKIN: Wash the affected area with soap and warm water.

IF IN EYES: Immediately flush eyes with running water for at least 15 minutes. Call a Physician.

See side panel for additional precautionary statements.

Mfg. by Agricultural Products Co.
Ft. Worth, Texas 76112

EPA Est 43789-TX-01
EPA Reg No 43789-62

Net Contents _____

Mix the recommended amount of water. Thorough and uniform coverage on both top and bottom surfaces of plants.

VEGETABLES - Unless otherwise specified, apply for thorough coverage. Start application when symptoms appear and repeat as directed.

BEANS - Anthracnose, downy mildew. Apply at 7-14 day intervals to within 7 days of harvest.

CARROTS - Alternaria leaf blight. Apply in 7 days of harvest if tops are to be eaten.

CELERY - Early and late blight. Apply weekly. Remove residue after harvest.

CORN - (SWEET) - Helminthosporium leaf blight. Apply at 3-7 day intervals.

LETTUCE - Downy mildew. Apply at 7-10 day intervals.

PEPPERS - Anthracnose, late blight. Apply at 7-10 day intervals.

TOMATOES - Anthracnose, late blight, spots and Stemphylium. Use 2-3 times if disease is severe. Do not apply within 14 days of harvest.

FRUITS - Apply at 1-2 week intervals.

APPLES - Apple blotch, bitter rot, fly speck, frog-eye, scab. Apply in delayed dormant and curative stages.

CITRUS - To control rust mite. Apply when rust mite is present. Do not apply after petal fall.

PEACHES, APRICOTS - Control rust mite. Apply when rust mite is present. Do not apply after petal fall.

STRAWBERRIES - Leaf spot. Apply when new growth starts in spring.

ORNAMENTALS - Use 1 level of application.

AZALEAS, CAMELLIAS - Petal blight. Direct spray into flowers. Spray at 7-14 day intervals.

CARNATIONS, CHRYSANTHEMUMS - leaf spot, rust. Apply at 7-14 day intervals. For control of Botrytis, apply weekly during the blooming season.

GLADIOLUS - Botrytis blight. Apply often under severe disease conditions.

ROSES - Black spot. Apply at 7-14 day intervals.

TURF GRASSES - Fading-out. Use 2 oz. in 3-5 gal. per 1,000 sq. ft. and repeat every 7-14 days until green.

- Use 2 oz. in 3-5 gal. per 1,000 sq. ft. and repeat every 5-7 days when green. Continue applications until disease is controlled. See side panel for further details on use.

NOTICE

CONDITIONS OF SALE: Seller is not deemed to have accepted the following agreement of a duly authorized representative of the buyer. **WARRANTY LIMITATION:** Seller warrants that the product is reasonably fit for the use intended. This warranty is limited to the description of the product on the label hereof. **INHERENT RISKS:** The precautions for use are stated on the label hereof. However, it is impossible to eliminate all risks associated with use or application of this product under all weather conditions. **LIMITATION OF LIABILITY:** In no case shall the seller be liable for any damages resulting from the use or handling of this product in excess of the purchase price of the product.

MENTS
(S)

u, inhaled, or ab-
n clothing. Wear
ding. Wash with
st. Do not use or

and water after
shower or bathe
water before reus-

ot apply directly to
leaning of equip-
ed on this label.

ner inconsistent

L
food, or
is prohi-

mixture
to label
to Fed-
resources

s accord-
cedures
recovery
se of in a
mitted by

ical Dis-
ve proce-



ZINEB 75% FUNGICIDE

Controls Diseases on Vegetables, Fruits & Ornamentals.

ACTIVE INGREDIENT:	BY WT
Zinc Dinocrotylate	75.0%
INERT INGREDIENTS:	25.0%
	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

Statement of Practical Treatment

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit fluid is clear. Call a Physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

IF ON SKIN: Wash the affected area with soap and warm water.

IF IN EYES: Immediately flush eyes with running water for at least 15 minutes. Call a Physician.

See side panel for additional precautionary statements.

Mfg. by Agricultural Products Co.
Ft. Worth, Texas 76112

EPA Est. 43789-TX-01
EPA Reg. No. 43789-62

Net Contents _____

DIRECTIONS

Mix the recommended amount of this product in sufficient water to produce good coverage. Thorough and uniform coverage is essential for disease control. Spray uniformly, covering both top and bottom surfaces of foliage and fruit.

VEGETABLES - Unless otherwise indicated on vegetables, use 2 1/2 tbsp. per gal. of water for thorough coverage. Start applications at first sign of or prior to appearance of disease symptoms and repeat as directed.

BEANS - Anthracnose, downy mildew, rust. Begin when first blossoms appear and repeat at 7-14 day intervals to within 7 days of harvest.

CARROTS - Alternaria leaf blight, Cercospora leaf spot. Apply at 7-14 day intervals to within 7 days of harvest if tops are to be used as food or feed.

CELERY - Early and late blight. Spray plant beds at 3-5 day intervals. Continue field applications weekly. Remove residues remaining at harvest by stripping, washing or trimming. Do not apply within 14 days of harvest.

CORN - (SWEET) - Helminthosporium leaf blight, Puccinia rust. Apply when disease appears and repeat at 3-7 day intervals.

LETTUCE - Downy mildew. Apply at weekly intervals to within 14 days of harvest.

PEPPERS - Anthracnose, ripe rot, Phytophthora. Use 2 1/2-5 1/2 tbsp. per gal. of water. Repeat at 7-10 day intervals.

TOMATOES - Anthracnose, Cladosporium Gray mold, early and late blight, Septoria leaf spots and Stemphylium. Use 2-2 1/2 tbsp. per gal. water. Repeat at 7-10 day intervals or more often if disease is severe. Do not apply within 5 days of harvest.

FRUITS - Apply at 1-2 week intervals beginning before bloom. Spray plants thoroughly.
APPLES - Apple blotch, bitter rot, black rot, Botrytis, phoma fruit rot, Brooks spot, cedar rust, fly speck, frog-eye, scab, sooty blotch, quince rust. Use 2-4 tsp. per gal. Begin applications in delayed dormant and continue through cover sprays.

CITRUS - To control rust mite and associated russeting and greasy spot. Use 2 tsp. per gal. water. Apply when rust mites appear.

PEACHES, APRICOTS - Coryneum blight or shot hole, peach leaf curl. Use 4 tsp. per gal. of water. Do not apply after petal fall on apricots or to within 30 days of harvest on peaches.

STRAWBERRIES - Leaf spot. Use 6 1/2 tsp. per gal. water for thorough coverage. Begin when new growth starts in spring and repeat as necessary. For leaf scorch apply in new beds in late summer using 2-3 weekly applications. Do not apply within 7 days of harvest.

ORNAMENTALS - Use 1 level tbsp. per gal. water. Spray plants thoroughly.

AZALEAS, CAMELLIAS - Petal blight. Spray 2-3 times a week while flowers are opening. Direct spray into flowers. Spray ground under bushes thoroughly.

CARNATIONS, CHRYSANTHEMUMS, HOLLYHOCKS, SNAPDRAGONS - Anthracnose, leaf spot, rust. Apply at weekly to 10 day intervals as soon as disease conditions threaten. For control of Botrytis petal spot and Ascochyta ray blight make applications twice weekly during the blooming season.

GLADIOLUS - Botrytis blight, Curvularia, Stemphylium. Spray at weekly intervals, or more often under severe disease conditions, from early stages until spikes begin to open.

ROSES - Black spot. Apply at weekly intervals.

TURF GRASSES - Fading out (Curvularia), Helminthosporium diseases, rust, slime molds. Use 2 oz. in 3-5 gal. per 1,000 sq. ft. of turf. Begin applications as soon as diseases appear and repeat every 7-14 days until disease threat is past. For Pythium blight or grease spot, use 2 oz. in 3-5 gal. per 1,000 sq. ft. of turf. Begin applications as soon as diseases appear and repeat every 5-7 days when disease is mild or every 3-4 days when disease is severe. Continue applications until disease threat is past. Follow State turf disease control recommendations for further details on application timing.

NOTICE — READ CAREFULLY

CONDITIONS OF SALE - Seller's offer of this product is subject to and may vary by state. Buyer shall have accepted the following conditions of sale and warranty which may only be varied by written agreement of the authorized representative of Seller.

WARRANTY LIMITATION - Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the label. Seller's liability is limited to the purchase price of the product. Seller makes no other express warranties. **THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY and there are no warranties which extend beyond the description on the label hereof.**

INHERENT RISKS - The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with its use. Buyer assumes all risks associated with use or application of this product contrary to label instructions or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY - In no case shall Seller be liable for special, direct or consequential damages resulting from the use or handling of this product and no claim of any kind shall be greater in amount than the purchase price of the product in respect of which such damages are claimed.