

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(AND DOMESTIC ANIMALS)**

CAUTION

Causes eye and skin irritation. Harmful if swallowed, inhaled, or absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling. Avoid inhaling of dust. Do not use or store near food or feed.

Wash hands, arms, and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work and wash all clothing with soap and hot water before reusing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

- 1. PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.
- 2. PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State, or Local procedures under the Resource Conservation and Recovery Act.
- 3. CONTAINER DISPOSAL:** Dispose of liners according to approved Federal, State, or Local procedures under the Resource Conservation and Recovery Act. If container cannot be reused, dispose of in a sanitary landfill or by incineration if permitted by State and Local Authorities.
- 4. GENERAL:** Consult Federal, State, or Local Disposal Authorities for approved alternative procedures such as limited open burning.
- 5. STORAGE:** Store in a cool, dry place.



MANEB 75% FUNGICIDE

Controls many plant diseases (fungus) on certain Fruits and Vegetables and Roses.

ACTIVE INGREDIENT: Maneb (Manganese ethylenebisdithiocarbamate) 75.0%
(Metal manganese equivalent - 15.5%)
INERT INGREDIENTS: 25.0%

BY WT.
75.0%
25.0%
100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Statement of Practical Treatment

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit fluid is clear. Call a Physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth.

IF ON SKIN: Wash the affected area with soap and warm water.

IF IN EYES: Immediately flush eyes with running water for at least 15 minutes. Call a Physician.

See side panel for additional precautionary statements.

Mfg. by Agricultural Products Co.
Ft. Worth, Texas 76112

EPA Est. 43/89-TX-01
EPA Reg. No. 43789-64

Net Contents

Best results are obtained by using a sprayer to foliage indicating one gallon of water for spray and make thorough coverage. If difficult to wet, add 8 to 10 mixture.

This product may also be dusted over all treated surfaces. Do dusting.

Apply dust or spray mix evenly, facing of foliage and fruit. Use when disease first appears and as specified below throughout entire season. When there are applications.

This product may be used on:
ROSES: For control of Canine. Application when the first leaf.
TOMATOES: For control of (Stemphylium). Do not use. For control of anthracnose, apply as specified above, starting 5 days of harvest.
POTATOES: For control of when plants are 2 to 6 inches.
CARROTS: For control of A. intervals.
CUCUMBERS: For control of V. at 3-5 day intervals when 5 days of harvest.

LIMA BEANS: For control of within 4 days of harvest.

ONIONS: For control of downy blight. Apply at 7-10 day intervals. Do not apply with more than one effective means.

SPINACH: For control of downy blight. Do not apply with more than one effective means.

GRAPES: Great Lakes Area after bloom and 7 to 10 days after bloom.

LETTUCE and ENDIVE: For Do not apply within 10 days of planting and from fruiting.

CABBAGE: For control of diseases 7 to 10 days after planting 7 days of harvest.

NOTICE

CONDITIONS OF SALE: See label for full details.
WARRANTY LIMITATION: See label for full details.
CEMENT MERCHANTS: See label for full details.
INHERENT RISKS: The user assumes all risks.
LIMITATION OF LIABILITY: See label for full details.



MANEB 75% FUNGICIDE

Controls many plant diseases (fungus) on certain Fruits and Vegetables and Roses

ACTIVE INGREDIENT:	BY WT.
Maneb (Manganese ethylenebis(3-thiocarbamate))	75.0%
(Metallic manganese equivalent - 15.5%)	
INERT INGREDIENTS:	25.0%

100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

Statement of Practical Treatment

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit fluid is clear. Call a Physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth to mouth.

IF ON SKIN: Wash the affected area with soap and warm water.

IF IN EYES: Immediately flush eyes with running water for at least 15 minutes. Call a Physician.

See side panel for additional precautionary statements.

Mfg. by Agricultural Products Co.
Ft. Worth, Texas 76112

EPA Est 43789-TX-01
EPA Reg No 43789-64

Net Contents



DIRECTIONS

Best results are obtained by mixing the product with water and applying with a suitable sprayer to foliage and rated below. Mix 1/2 ounce of this product with one gallon of water for spray applications. Keep mixture agitated well when spraying and make thorough coverage of all leaves, both top and bottom surfaces. If foliage is difficult to wet, add 8 to 10 drops of a spreader-sticker to each gallon of spray mixture.

This product may also be dusted or foliar dusted equipment is available to evenly distribute the product. Dust will adhere better if foliage is wetted slightly before dusting.

Apply dust or spray mix evenly and thoroughly, covering both bottom and top surfaces of foliage and fruit. Unless otherwise specified, always begin applications when disease first appears and by making applications at 7 to 10 day intervals or as specified below throughout the growing season you will protect your plants the entire season. When there is a severe outbreak, shorten the interval between applications.

This product may be used on the following fruits, vegetables, and flowers:

ROSES: For control of Cercospora Leaf Spot, Rose Rust, and Black Spot, begin application when the first leaves unfold. Repeat application at 7-10 day intervals.

TOMATOES: For control of early blight, late blight, septoria leaf spot, gray leaf spot (Stemphylium). Do not use on tender seedlings in greenhouse as injury may result. For control of anthracnose, mix 5.0 ounces of this product in 10 gallons of water and apply as specified above. Repeat applications at 7-10 day intervals. Do not apply within 5 days of harvest.

POTATOES: For control of early blight and late blight. Applications should begin when the plants start to form tubers.

CARROTS: For control of Alternaria and Cercospora leaf spots. Apply at 7-10 day intervals.

CUCUMBERS: For control of downy mildew, anthracnose. Apply at 7-10 day intervals. Do not apply within 10 days of harvest when severe disease conditions exist. Do not apply within 10 days of harvest.

LIMA BEANS: For control of downy mildew. Apply at 4-7 day intervals. Do not apply within 10 days of harvest.

ONIONS: For control of downy mildew, purple blotch, blight, and blast (Botrytis leaf blight). Apply at 7-10 day intervals.

SPINACH: For control of downy mildew, root rot, and white rust. Apply at 7-10 day intervals. Do not apply within 10 days of harvest and remove residues by washing in water before use.

GRAPES: For control of Anthracnose. For control of black rot, spray just before bloom, just after bloom, and 10-15 days after bloom. Do not apply within 10 days of harvest.

LETTUCE and ENDIVE: For control of downy mildew. Apply at 7-10 day intervals. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and from romaine and from leaf lettuce and endive by washing in water before use.

CABBAGE: For control of Alternaria leaf spot and downy mildew. Begin applications 10-15 days after planting and repeat at 5-7 day intervals. Do not apply within 7 days of harvest.

NOTICE-READ CAREFULLY

CONDITIONS OF SALE: This product is sold as a fungicide and is not intended for use as a herbicide. It is sold as a fungicide and is not intended for use as a herbicide. It is sold as a fungicide and is not intended for use as a herbicide.

WARRANTY LIMITATION: This product is sold as a fungicide and is not intended for use as a herbicide. It is sold as a fungicide and is not intended for use as a herbicide. It is sold as a fungicide and is not intended for use as a herbicide.

INHERENT RISKS: This product is sold as a fungicide and is not intended for use as a herbicide. It is sold as a fungicide and is not intended for use as a herbicide. It is sold as a fungicide and is not intended for use as a herbicide.

LIMITATION OF LIABILITY: This product is sold as a fungicide and is not intended for use as a herbicide. It is sold as a fungicide and is not intended for use as a herbicide. It is sold as a fungicide and is not intended for use as a herbicide.