

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(AND DOMESTIC ANIMALS)**

WARNING

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Wash contaminated clothing before reuse. Do not use or store near food or feed. Do not use on household pets or humans. Do not contaminate ornamental fishponds. Do not permit children or pets to go into treated areas until the spray has dried completely. **DO NOT USE IN THE HOME**

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Agricultural Extension Service.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

- 1. STORAGE:** Store in a cool place. Use contents within 6 months after opening. Keep container tightly closed.
- 2. CONTAINER DISPOSAL:** Do not reuse empty container. Wrap container and put in trash collection.

General Information:

Kills many outdoor insects which destroy lawns, roses, flowers, trees, and shrubs or annoy pets and people. (Refer to label directions for specific insects controlled.) Contains Diazinon, insecticide which provides positive control of many hard-to-kill sucking and chewing insects. This product works fast and remains effective for an extended period of time.

**DIAZINON* 25% EC
LAWN AND GARDEN INSECT CONTROL**

PROTECTS	CONTROLS
Lawns	Aphids
Vegetables	Chinch Bugs
Fruits & Nuts	Caterworms
Roses	Mites
Flowers	Sod Webworms
Trees	And many others
Shrubs	(See back label)

ACTIVE INGREDIENTS:	BY WT
0.000% Diazinon (25% EC)	25.00%
66.48% Inert Ingredients	66.48%
8.52% Inert Ingredients	8.52%
100.00%	100.00%

*Diazinon trademark of CIBA GEIGY Corp

**KEEP OUT OF REACH OF CHILDREN
WARNING**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. Do not induce vomiting. Vomiting may cause aspiration pneumonia.
IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.
IF ON SKIN: Wash the effected area with soap and warm water.
IF IN EYES: Flush eyes with plenty of running water. Call a physician if irritation persists.
See reverse panel for additional precautionary statements

Manufactured by Agricultural Products Company
Ft. Worth, Texas 76112

EPA Establishment Number 43789 TX-01
EPA Registration Number 43789-66

Net Contents

How to Mix and Apply:

Step # 1: Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Repeat applications as needed or as noted in the rate chart.

Step #2: Use clean sprayer. Carefully measure and mix the amount of this product and water as suggested in the chart below.

Step #3: Spray as directed in rate chart. Use coarse drenching sprays for lawns, soil and other surface applications. Use medium-fine mist sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

Lawns

Apply Diazinon 25% EC to lawns. Apply to lawns in early spring, late spring, and late summer. Apply to lawns in early fall. Apply to lawns in late fall. Apply to lawns in winter. Apply to lawns in spring. Apply to lawns in summer. Apply to lawns in fall. Apply to lawns in winter.

1 1/2 oz in 3 gals of water covers 125 sq ft

8 1/2 oz in 24 gals of water covers 1,000 sq ft

Thoroughly wet lawn grass a few hours before applying.

*Brown Dog Ticks: Do not apply to animals. In particular, spray under shrubbery and the grasses next to house.

**Lawn Billbugs: Water the lawn thoroughly immediately after treatment.

†In the North, Chinch Bugs are the only lawn insect problem. As a rate of application can be used: 1 1/2 oz in 3 gals of water covers 125 sq ft.

Milipedes: 2 1/2 oz in 3 gals of water covers 125 sq ft

Thoroughly wet grass a few hours before applying.

Rhodesgrass Scales

1 1/2 oz in 3 gals of water covers 125 sq ft

Apply when Scale Crawlers first emerge.

Ants, Digger Wasps*

1 1/2 oz in 3 gals of water covers 125 sq ft

Spot spray ant hills and wasp ground nest openings.

*Make the application at dusk.

††Billbugs: White Grubs** of Japanese Beetle, European Chaffer, Southern Chaffer.

1 1/2 oz in 3 gals of water covers 125 sq ft

Water grass thoroughly after application.

*Billbug control: Apply when activity is first observed or when chewed or brown grass is present, indicating damage from the insect.

**Grub control: Make application any time between late July and early October.

Hyperodes Weevil (New York only)

1 1/2 oz in 3 gals of water covers 125 sq ft

Water grass thoroughly after application. Treat problem areas in mid April and again in mid May.

Trees & Shrubs

Apply to trees and shrubs. Apply to trees and shrubs in early spring, late spring, and late summer. Apply to trees and shrubs in early fall. Apply to trees and shrubs in late fall. Apply to trees and shrubs in winter. Apply to trees and shrubs in spring. Apply to trees and shrubs in summer. Apply to trees and shrubs in fall. Apply to trees and shrubs in winter.

1 1/2 oz in 3 gals of water. Apply as a thorough foliar spray.

1 1/2 oz in 3 gals of water. Apply as a thorough foliar spray.

1 1/2 oz in 3 gals of water. Apply as a thorough foliar spray.

*Mimosa Webworms: Apply as a thorough foliar spray when insects first appear. Make two additional applications at 4 to 5 week intervals.

Roses & Flowers

Azaleas
Camellias
Clematis
Cypripedium
Garden Phlox
Hydrangea
Roses

(Note: Do not apply to any plants in bloom.)

Fruits & Nuts

Almonds
Figs
Pecans
Walnuts

Apples
Pears

Apricots
Peaches
Nectarines

Cherries

Figs

Plums
Prunes

Caneberries
Blackberries
Boulevardberries
Loganberries
Raspberries

Blueberries

Strawberries

Grapes

ACCEPTED
AUG 18 1981
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 43789-66.

DIAZINON* 25% EC LAWN AND GARDEN INSECT CONTROL

PROTECTS
Lawns
Vegetables
Fruits & Nuts
Roses
Flowers
Trees
Shrubs

CONTROLS
Aphids
Chinch Bugs
Cutworms
Mites
Sod Webworms
And many others
(See back label)

ACTIVE INGREDIENTS:
Diazinon 25.00%
Other Inert Ingredients 75.00%

BY WT
25.00%
66.48%
8.52%
100.00%

Diazinon trademark of CIBA-GEIGY Corp.

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED: Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken orally. Do not induce vomiting. Vomiting may cause aspiration pneumonia.
IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.
IF ON SKIN: Wash the effected area with soap and warm water.
IF IN EYES: Flush eyes with plenty of running water. Call a physician if irritation persists.
See inside panel for additional precautionary statements.

Manufactured by Agricultural Products Company
North, Texas 76112

Establishment Number 43789 TX-01
Registration Number 43789-66

Net Contents

Mix and Apply:

Determine size of spraying job. For lawns and vegetation measure the area to be sprayed. For all other applications, use a thorough cover spray. Make applications when insects are active. Repeat applications as needed or as noted in the rate chart.

Use clean sprayer. Carefully measure and mix the product and water as suggested in the chart below.

Spray as directed in rate chart. Use coarse drenching for lawns, soil and other surface applications. Use medium nozzles for trees and plants. Thorough coverage is important. Rinse sprayer with clean water after each use.

Use this rate chart for lawns and garden plants.

Lawns

Ants: Army Ants, Fire Ants, Imported Fire Ants, Brown Buggers, etc. (Cottonwood, etc.)
Chinch Bugs: (Cottonwood, etc.)
Cutworms: (Cottonwood, etc.)
Mites: (Cottonwood, etc.)
Sod Webworms: (Cottonwood, etc.)
Whiteties: (Cottonwood, etc.)

1 1/2 oz in 3 gals of water covers 125 sq ft
8 1/2 oz in 24 gals of water covers 1,000 sq ft

Thoroughly wet down grass a few hours before applying.

*Brown Buggers: Do not apply to animals. In particular, spray under shrubbery and the grass next to house.

**Lawn Billbugs: Water the lawn thoroughly immediately after treatment.

†In the North, Chinch Bugs are the only lawn insect problem; a lower rate of application can be used: 1 1/2 oz in 3 gals of water covers 125 sq ft.

Meepedes: 2 1/2 oz in 3 gals of water covers 125 sq ft.

Thoroughly wet grass a few hours before applying.

Rhodesgrass Scales:
1 1/2 oz in 3 gals of water covers 100 sq ft. Apply when Scale Crawlers first emerge.

Ants, Digger Wasps:
1 1/2 oz in 3 gals of water covers 125 sq ft. Spot spray ant hills and wasp ground nest openings.
*Make the application at dusk.

Grubs: White Grubs of Japanese Beetle, European Chaffer, Southern Chaffer.**
1 1/2 oz in 3 gals of water covers 125 sq ft.
Water grass thoroughly after application.

*Billbug control: Apply when activity is first observed or when chewed or brown grass is present, indicating damage from the insect.

**Grub control: Make application any time between late July and early October.

Hyperodes Weevil (New York only):
1 1/2 oz in 3 gals of water covers 125 sq ft.
Water grass thoroughly after application. Treat problem areas in mid April and again in mid May.

Trees & Shrubs

Azalea: Azalea Bark Scale, Azalea Bark Scale, Azalea Bark Scale, etc.
Berberis: Berberis Scale, Berberis Scale, etc.
Camellia: Camellia Scale, Camellia Scale, etc.
Cherry: Cherry Scale, Cherry Scale, etc.
Fig: Fig Scale, Fig Scale, etc.
Juniper: Juniper Scale, Juniper Scale, etc.
Maple: Maple Scale, Maple Scale, etc.
Oak: Oak Scale, Oak Scale, etc.
Pine: Pine Scale, Pine Scale, etc.
Poplar: Poplar Scale, Poplar Scale, etc.
Rhododendron: Rhododendron Scale, Rhododendron Scale, etc.
Spice: Spice Scale, Spice Scale, etc.
Wick: Wick Scale, Wick Scale, etc.

1 1/2 oz in 3 gals of water. Apply as a thorough four spray.

Webworms: Juniper, Mimosa, Whiteties.

1 1/2 oz in 3 gals of water. Apply as a thorough four spray.

*Mimosa Webworms: Apply as a thorough four spray when adults first appear. Make two additional applications at 4-5 week intervals.

Roses & Flowers:
Aphids, European Red, etc.
1 1/2 oz in 3 gals of water. Apply as a thorough four spray.

Roses & Flowers:
Aphids, European Red, etc.
1 1/2 oz in 3 gals of water. Apply as a thorough four spray.

Fruits & Nuts:
Aphids, Scale Crawlers, etc.
1 1/2 oz in 3 gals of water. Apply as a full cover spray.

Apples:
Aphids, Codling Moths, etc.
1 1/2 oz in 3 gals of water. Apply as a full cover spray.

Cherries:
Black Cherry Aphids, etc.
1 1/2 oz in 3 gals of water. Apply as a full cover spray.

Blueberries:
Aphids, Blueberry Maggots, etc.
1 1/2 oz in 3 gals of water. Apply as a full cover spray.

ACCEPTED
AUG 18 1981
Under the Federal Insecticide, Fungicide and Rodenticide Act, registered under EPA Reg. No. 227-226-01

Grapefruit
Lemons
Oranges

Citrus Thrips (see instructions on container with Gutter Spray)
1 fl oz in 3 gals of water. Apply as a full cover spray.
12 fl oz in 3 gals of water. Apply as a full cover spray after blossoming and again in the summer. Thorough spray coverage of tree trunk, limbs and twigs is essential.

*After spraying do not pick fruit within
5 days - Strawberries, Figs
7 days - Blueberries, Caneberries
10 days - Apricots, Cherries, Grapes, Nectarines, Plums, Prunes
14 days - Apples, Pears

Vegetables*

Foliar Application

Thoroughly spray all plant parts. Generally, applications at 7 to 10-day intervals may be required. Do not feed treated vines to livestock.

Beans
Peas

Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Mites
1 fl oz in 3 gals of water covers approximately 1,000 sq ft of garden

Cabbage,
Cauliflower,
Collards,
Broccoli,
Kale, Radishes,
Turnips

Aphids, Diamondback Moths, Flea Beetles, Imported Cabbage Worms, Dipterous Leaf Miners
1 fl oz in 3 gals of water covers approximately 1,000 sq ft of garden

Carrots, Celery,
Endive, Lettuce,
Parsnip, Spinach

Aphids, Flea Beetles, Leafhoppers, Dipterous Leaf Miners, Mites
1 fl oz in 3 gals of water covers approximately 1,000 sq ft of garden

Cantaloupes,
Cucumbers,
Muskmelon,
Squash (summer,
winter),
Watermelon

Aphids, Thrips, Mites
1 fl oz in 3 gals of water covers approximately 1,000 sq ft of garden

Peppers,
Potatoes,
Tomatoes

Aphids, Colorado Potato Beetles, Fall Armyworms (tomato), Flea Beetles, Leafhoppers, Southern Armyworms (potato and tomato), Vinegar Flies (*Drosophila* spp. (tomato))
1 fl oz in 3 gals of water covers approximately 1,000 sq ft of garden

Onions

Onion Thrips
1 fl oz in 3 gals of water covers approximately 1,000 sq ft of garden

Sweet Corn

Aphids, Flea Beetles, Grasshoppers, Mites
1 fl oz in 3 gals of water covers approximately 1,000 sq ft of garden

Vegetables*

Preplant Soil Application

Cutworms (Surface, Subterranean)

6 fl oz per 1,000 sq ft in sufficient water to achieve thorough coverage

Apply broadcast just before planting. Immediately work into soil 2-3 inches for Surface Cutworms, 3-6 inches for Subterranean Cutworms.

Garden Symphylans (Garden Centipedes)

15 fl oz per 1,000 sq ft in sufficient water to achieve thorough coverage

Apply broadcast just before planting. Immediately work into soil 4-8 inches.

Mole Crickets, Carrots, Corn, Peas, Pole Beans, Red Beets and Snap Beans Only

15 fl oz per 1,000 sq ft in sufficient water to achieve thorough coverage

Apply broadcast just before planting. Apply 1-2 days before sowing or transplanting. Immediately work into soil 2 inches.

Next Page: 10/10/01

4 fl oz per 1,000 sq ft in sufficient water to achieve thorough coverage

Apply broadcast just before planting. Immediately work into soil 4 inches before sowing, either in seed bed or in field or before transplanting.

Wireworms, Cabbage, Carrots, Corn, Lettuce, Peas, Snap Beans and Tomatoes Only

6 fl oz per 1,000 sq ft in sufficient water to achieve thorough coverage

Apply broadcast just before planting. Immediately work into soil 4-8 inches.

*After spraying do not pick vegetables within

1 day - Peas, Tomatoes, Sweet Corn

3 days - Cantaloupes, Muskmelons, Winter Squash, Watermelons

5 days - Broccoli, Cauliflower, Peppers

7 days - Beans, Cabbage, Cucumbers, Summer Squash

10 days - Carrots, Celery, Collards, Endive, Kale, Lettuce, Onions, Parsnips, Radishes, Spinach, Turnips

25 days - Potatoes

Nuisance Pests in Outside Areas

Ants†, Boxelder Bugs, Brown Dog Ticks†, Chiggers†, Clover Mites†, Cockroaches, Crickets†, Earwigs†, Fleas†, Flies†, Millipedes†, Sowbugs (Pillbugs)†, Spiders, Springtails (Collembola)†

8 fl oz in 3 gals of water

Thoroughly spray window and door screens, sills, foundations, porch and patio walls and ceilings, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide.

Do not treat plants or animals.

*Spray a 5 ft. band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet.

†See label recommendations for other control measures for these pests.

NOTICE — READ CAREFULLY

CONDITIONS OF SALE: Seler(s) offers, this product for sale subject to, and buyer and all users are deemed to have accepted, the following conditions of sale and warranty which may only be varied by written agreement of a duly authorized representative of Seler(s).

WARRANTY LIMITATION: Seler(s) warrant(s) that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the label subject to the inherent risks referred to below. Seler(s) make(s) no other express warranties. **THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY** and there are no warranties which extend beyond the description on the label hereof.

INHERENT RISKS: The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with its use. Buyer assumes all risks associated with use or application of this product contrary to label instructions or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY: In no case shall Seler(s) be liable for special, indirect, or consequential damages resulting from the use or handling of this product and no claim of any kind shall be greater in amount than the purchase price of the product in respect of which such damages are claimed.