## **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS** (AND DOMESTIC ANIMALS) CAUTION

May be harmful if swallowed. Do not breathe dust. Do not get in eyes, on skin, or on clothing. May cause eye or skin irritation. During exposure in treatment and loading operations wear goggles, clean synthetic rubber gloves, and protective clothing. Wash hands, arms, and face thoroughly with soap after handling and before eating or smoking Routinely shower or bathe after work and wash work clothes with soap and hot water before reusing.

#### **ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Exposed treated seeds may be hazardous to birds and other wildlife.

## **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

## STORAGE AND DISPOSAL

- 1. PROHIBITIONS: Do not contaminate water food or feed by storage or disposal. Open dumping is prohibited
- 2. PESTICIDE DISPOSAL: Pesticide, spray mixture or rinsate that cannot be used according to tabel instructions must be disposed of according to Federal. State, or Local procedures under the Resource Conservation and Recovery Act
- 3. CONTAINER DISPOSAL: Triple rinse for equivalent) and offer for recycling or reconditioning or dispose of in a same ary landfill or by incineration if permitted by State and Local Authorities
- 4. GENERAL: Consult Federal, State or Local Disposal Authorities for approved a ternative procedures such as limited open burning
- 5. STORAGE Store in a coor dry place



# PLANTER BOX SEED TREATER

**ACTIVE INGREDIENT: CAPTAN. N**-. Trick commethy  $(4.6 \, \text{GeV})^2$  at enemed 2-dicarboximide  $\pi$ 47.50% Related denvitives 2.50% **INERT INGREDIENTS** 50.00 % 100.00%

(Fungicide with MOLYBDENUM and non-oil lubricant)

## KEEP OUT OF REACH OF CHILDREN CAUTION

Statement of Practical Treatment

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit fluid is clear. Call a Physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

IF ON SKIN: Wash the affected area with soap and warm

IF IN EYES: Immediately flush eyes with running water for at least 15 minutes. Call a Physician.


See side panel for additional precautionary statements.


# STORE IN A COOL, DRY PLACE

Mfg. by Agricultural Products Co. Ft. Worth, Texas 76112

> EPA Est. 43789-TX-01 EPA Reg. No. 43789-74

\_\_\_\_ Net Contents





### DIRECTIONS

MEASURING SCOOP IS ENCLOSED. One level scroop holds approximately 4 ounces of product.

SOYBEANS: Use CAPTAN 50 + MOLY-LUBE SEED TREATER as a planter box treatment at the rate of 2 ounces per bushel or 312 ounces per hundred pounds of seed.

CAPTAN 50 + MOLY-LUBE SEED TREATER is a specifically designed formulation for the treatment of soybean seed. Protection is provided against seed-borne and soil-borne fungi which cause seed decay damping-off and seeding plights. At the recommended dosage CAPTAN 50 + MOLY-LUBE SEED TREATER is noninjurious to seeds or seedlings and the germination of treated seed is frequently stimulated. giving a superior result as compared to results from untreated seed.

FOR IMPROVED MIXING OF PRODUCT WITH SEED, it is recommended that the seed be premixed in a suitable container with the product, then transferred to the planter box. Similar results may be obtained by filling the planter box half full of seed, then adding half the required amount of product, followed by thorough mixing with a stick, and finally, the second half of the seed and product, repeating the mixing.

IMPORTANT: Treated seed must not be used for or mixed with food or animal feed or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Seed commercially treated with this product must be labeled in accord with all applicable requirements of the Federal Seed Act. Labels allached to treated seed must state. Treated seedmust not be used for or mixed with food or animal feed, or processed for oil, and. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial awary from bodies of water.

### NOTICE-READ CAREFULLY

CONDITIONS OF SA. Seller(s) offer(s) this product for sale subject to, and buyer and all users are ceemed to have accepted, the following conditions of sale and warranty which may only be varied by written agreement of a dury authorized representative of Seller(s):

WARRANTY LIMITATION: Seller(s) warrant(s) that this product conforms to the chemical description on the label and is reasonably lit for the purposes referred to in the directions for use on the label subject to the inherent risks referred to below. Seller(s) make(s) no other express warranties. THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY and there are no warranties which extend beyond the description on the label hereof

INHERENT RISKS: The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with its use. Buyer assumes all risks associated with use or application of this product contrary to label instructions or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY: In no case shall Seller(s) be liable for special, indirect, or consequential damages resulting from the use or handling of this product and no claim of unit kind shall be greater in amount than the purchase price of the product in resnect of which such damages are claimed.