

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS  
CAUTION**

May cause irritation of nose, throat, eyes, or skin. Do not breathe dust or spray mist. During exposure to seed treatment and loading operations, wear goggles, clean synthetic rubber gloves, and protective clothing.

Wash hands, arms, and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work and wash all clothing with soap and hot water before reusing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Exposed treated seeds may be hazardous to birds and other wildlife.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read all directions before using and use only as specified on this label.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool dry place.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state or local authorities.

**BEST AVAILABLE COPY**



**CHLOROVAX 200 Liquid**  
Planter Box Seed Treater

<b>ACTIVE INGREDIENT</b>	<b>BY WT.</b>
Chloroneb (1, 4-dichloro-2,5-dimethoxybenzene .....	20.0%
<b>INERT INGREDIENTS</b> .....	80.0%
	100.0%

THIS PRODUCT CONTAINS THE TOXIC INERT INGREDIENT RHODAMINE DYE

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**Statement of Practical Treatment**

**IF SWALLOWED:** Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit fluid is clear. Call a Physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

**IF ON SKIN:** Wash the affected area with soap and warm water.

**IF IN EYES:** Immediately flush eyes with running water for at least 15 minutes. Call a Physician.

See side panel for additional precautionary statements.

ACCEPTED

OCT 29 1987

EPA Est.  
EPA Reg. No. 43789-77

Net Contents

Made in U.S.A. by

AGRICULTURAL PRODUCTS CO., GREELEY, CO 80631

43789-77

Prad

10F1

**DIRECTIONS AND APPLICATION RATES**

This product is an agricultural fungicide formulated for seed treatment at planting time. It contains effective binding agents which reduce "dusting-off" of the treatment and yet is easily cleaned from equipment with water. This product should be used in combination with a standard seed treatment fungicide. Do not use excess treated seed for food, feed or oil purposes.

Dilute this product in the amount of water necessary for good coverage of the type of seed to be treated (usually 1 part of this product to 1-3 parts of water, depending upon type of seed - smaller seeds require higher dilution rate for good coverage). For best results it is recommended that seed be premixed in a suitable container with the product, then transferred to the planter box.

Best results are obtained when seed are allowed to dry prior to being transferred to the planter box.

**BEANS:** For control of Rhizoctonia solani, Pythium Spp., and Seed and seedling diseases use 2½-7 fluid ounces of this product per 100 pounds of seed.

**SOYBEANS:** For control of Rhizoctonia solani, Pythium Spp., and Seed and seedling diseases use 2½-7 fluid ounces of this product per 100 pounds of seed.

**COTTON:** For control of Rhizoctonia solani, Pythium Spp., Sclerotium rolfsii, and Seed and seedling diseases use 8-17½ fluid ounces of this product per 100 pounds of seed.

**SUGAR BEETS:** For control of Rhizoctonia solani, Pythium Spp., and Seed and seedling diseases use 3½-10½ fluid ounces of this product per 100 pounds of seed.

Use high rate on pathologically infested seed or for soils such as in California.

**NOTICE — READ CAREFULLY**

**CONDITIONS OF SALE:** Seller(s) offer(s) this product for sale subject to, and buyer and all users are deemed to have accepted, the following conditions of sale and warranty which may only be varied by written agreement of a duly authorized representative of Seller(s).

**WARRANTY LIMITATION:** Seller(s) warrant(s) that the product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the label subject to the inherent risks referred to below. Seller(s) make(s) no other express warranties. **THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY** and there are no warranties which extend beyond the description on the label hereof. **INHERENT RISKS:** The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with its use. Buyer assumes all risks associated with use or application of this product contrary to label instructions or resulting from extraordinary weather conditions.

**LIMITATION OF LIABILITY:** In no case shall Seller(s) be liable for special, indirect or consequential damages.