US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS	43797-7	NOV 4 1985
REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460	TERM OF ISSUANCE	
NOTICE OF PESTICIDE:	NAME OF PESTICIDE PRODUCT	
(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	Custom Insectio	ide Spray
	Concentrate	
ME AND ADDRESS OF REGISTRANT (Include ZIP code)	Г	
Custom Packaging Services Division of J. Coburn, Inc. 427 Evernia Street		
West Palm Beach, FL 33401	_1	

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cance: the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered

by other this product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment:

1.84

うったがいになる。アイリアはなどのというないないというようなわずないない

- a. Add the phrase "EPA Registration No. 43797-7."
- b. Delete the statement "Authorized for use in all departments of official establishments operating under the meat, poultry, shell egg grading and egg products inspection programs." Either obtain U.S. Department of Agriculture approval of your label before shipping this product or remove all mention of "Féderally inspected meat and poultry plants" included in the section under Directions For Use.
- c. Delete "Food additive tolerances have been established for the insecticide components in this product. When this product is used as directed, allowable tolerance levels will not be exceeded."

ATTACHMENT IS APPLICABLE SIGNATURE OF APPROVING OFFICIAL DATE 11-4-85

EPA Form 8570-6 (Rov. 5.76) 87802:Gardner/Schroeder:RD-85/25:KENCO: 10/28/85:11/6/85:DKD: VO

45-17- 215

(

2

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Ċ

Timothy A. Gardner Product Manager (17) Insecticide-Rodenticide Branch Registration Division (TS-767)

Enclosure

CUSTOM

AN INSELLICIDE SPPAY CONCENTRATE

FOR USE IN FOOD PROCESSING AND STORAGE, MUSHROOM HOUSES, HOSPITALS, HOTELS, MOTELS, RESTAURANTS, OTHER INSTITUTIONS AND IN RESIDENCES

Authorized for use in all departments of official establishments operating under the meat, poultry, shell egg grading and egg products inspection programs.

PROVIDES RAPID KILL OF INSECTS WITHOUT LEAVING OBJECTIONABLE RESIDUES.

Food additive tolerances have been established for the insecticide components in this product. When this product is used as directed, allowable tolerance levels will not be exceeded.

Kills Gypsy Moth Caterpillars and Japanese Beetles

ACTIVE INGREDIENTS:	
Tyretbring	1.01
"Piperonyl Butoxide, Technical	30.07
INERT INGREDIENTS:	67.01
	100 01

*Equivalent to 24% (butylcarbityl)(6-propylplperonyl)ether and 6% related compounds.

PYREHORE - Registered Trademark of Fairfield American Corporation.

KEEP OUT OF REACH OF CHILDREN CAUTION

See reverse side for additional precautions.

EPA Reg. No. EPA Est. No. 43797 FLOI

DIRECTIONS FCR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. NET CONTENTS: _____ To prepare dilutions the concentrate shi first be stirred or agitated well. Add required amount of concentrate to water blend thoroughly. This will produce a micro-emulsion.


For maximum effectiveness, a combination localized surface application and space ment is recommended.

AS A SURFACE SPRAY: For Initial clean severe insect infestation, dilute at a 1 part concentrate in 14 parts water (8 ounces per gallon). This dilution may used in edible product areas of federal inspected meat and poultry plants. In plants and for normal infestations dilu part concentrate in 29 parts water (4.3 per gallon).

To control cockroaches, ants, silverfis crickets, clover mites, cheese mites, g weevils, rice weevils, confused flour be rust red flour beetles, sawtoothed grai spider beetles, clgarette beetles, drug beetles, meal worms, grain mites and ca Use a good sprayer adjusted to deliver wet spray. Direct the spray into hidin cracks and cravices, under pallets, aro tainers of stored foods, around the bas machinery, behind shelves and drawers. bookcases for <u>silverfish</u>. Spray ant tr nests and points of entry. If surface cation only is to be used, spray floors and other surfaces applying at a rate o to 750 square feet of surface.

To control <u>carpet beetles</u>, spray edges ing and under carpeting and rugs. Make application to floor and baseboards. S directly into cracks, closets and infes of shelving. Repeat treatment as neede

To control bedbugs: Spray mattresses 1 particularly around tufts and seams. 1 apart and spray in all joints. Treat b moldings and floors. Repeat treatment



To control fleas, and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. To kill fleas and ticks on cats and dogs, and to obtain protection against reinfestation, dilute at 1 part concentrate to 19 parts water 5.4 fluid ounces per 1 gallon) and wet the animal by dipping or spraying.

AS A SPACE SPRAY: Dilute at a rate of 1 part concentrate to 19 parts water (6.4 <u>fluid</u> OZ. per gallon). Apply with mechanical or <u>com-</u> pressed air equipment (non-thermal) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems. For rapid control of <u>adult</u> house flies, <u>fruit</u> flies, <u>gnats</u>, <u>mosquitoes</u>, <u>skipper flies</u>, <u>wasps</u>, <u>hornets</u>. <u>Angoumois grain moths</u>, <u>Hediterranean flour moths</u>, <u>Indian meal moths</u>, and <u>tobacco roths</u>: <u>Birect</u> spray at an upward angle distributing it uniformly through the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Ventilate and sweep up dead insects before reoccupying.

For rapid kill of exposed or accessible stages of other insects named on this label; dilute 1 part concentrate in 1 - 9 parts water { 13 €0 64 ounces per gallon}. Apply using conventional, mechanical or compressed air equipment (non-thermel), following directions for space spraying. Apply the above dilutions at a rate of from 1 to 10 ounces per 1000 cubic feet of space. When used in meat and poultry processing plants, a dilution of no less than 1 to 1 is acceptable and it must be dispersed from equipment adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 802 less than 30 microns).

IN MUSHROOM PRODUCTION AND PROCESSING: To control adult <u>Mushroom flies</u>, or <u>Fungus gnats</u>. <u>Phorids</u>, <u>Sciards</u>, and <u>Lecidomylids</u> in growing houses. Dilute 1 part concentrate in 1 part water {.64 fluid ounces per gallon}. Using mechanical or compressed air equipment (non-tnermal) adjusted to delive a line misi, apply at a rate of 1 punce per 1600 kubic freet of space. Fogger should be operated into 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results rairigecation should be turned off. In packing and canding plants gas at a rate of 1 ounce to 2000 ouble feet of space.

...........

OUTDOORS: To kill gypsy roth caterpillars and adults, adult Japanese huntles and Eastern and forest tent caterpillars on contact: combine 3.0 fiuld ounce of concentrate per 1 gallon of water. Apply to insure thorough coverage of upper and lower surfaces.

- On Trees and Shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, meking sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.
- On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

· CAUTION

Harmful If swallowed. Remove pets, birds and cover fish aquaria before spraying. Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protective clothing, Wash exposed skin surfaces with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and water before reuse.

Bo not apply as a space spray while food processing is underway.¹ Except in Federally inspected meat and poultry plants, when applied as a surface spray with care and in accordance with the directions and precautions given above, food processing operations may continue. Do not apply at any time as a undiluted space spray in edible product areas of Federally inspected meat and poultry plants. Foods should be removed or covered before treatments. Thoroughly wash all food processing surfaces before reuse. All surfaces in processing areas must be washed and rinsed in potable water after spraying.

1

ENVIRONMENTAL HAZARDS

Page 2 of 2

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes, and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes.

CHEMICAL HAZARDS

Do not use or store near heat or open flame.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

> Custum Packaging Setuices West Palm Beach, Ft 33401

> > ٠.

shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Timothy A. Gardner Product Manager (7) Insecticide-Rodenticide Branch Registration Division (TS-767)

Enclosure