

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER**

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing. Wear goggles when handling. Wash thoroughly with soap and water after handling.

**ENVIRONMENTAL HAZARD**

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**DIRECTIONS FOR USE:**

Notice to User: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Prohibitions - Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal - Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Trademark of Janssen Pharmaceutica, 49340 Beerse, Belgium

Active ingredient is manufactured by Janssen Pharmaceutica, 49340 Beerse, Belgium. U.S. Patent No. 3,658,813

Janssen Pharmaceutica, 1980

ANTHRAKINONE DERIVATIVE

**Fungafloor**

An Emulsifiable Concentrate for Postharvest Treatment of Citrus Fruit Containing 6.77 lb. active ingredient/gal.

Active Ingredient (w/w)  
Irazalil: 1-[2-(2,4-dichlorophenyl)-2-(2-propenyloxyethyl)-1H-imidazole]\*.....680  
Inert Ingredients (w/w).....320

U.S. Pat. No. 3,658,813

KEEP OUT OF REACH OF CHILDREN

**DANGER**

**STATEMENT OF PRACTICAL TREATMENT**

If Swallowed: Induce vomiting, pump stomach, call a physician. Treat symptomatically.

If Inhaled: See a physician if an abnormal reaction occurs.

If on Skin: Wash thoroughly with soap and water.

If in eyes: Immediately flush eyes with plenty of water. Call a physician.

See label for additional precautionary statements.



JANSSEN PHARMACEUTICA  
49340 Beerse, Belgium  
Belgium, U.S. Patent No. 3,658,813

FIA Reg. No. 41813-1  
FIA Est. No. 41813-1102  
Patent No. 3,658,813

Intended for post-harvest control of *Penicillium digitatum* (green mold), *Penicillium italicum* (blue mold), *Diaporthe citri* (phomopsis stem-end rot), *Diplodia natalensis* (diplodia rot) and *Alternaria citri* on citrus fruits.

Used as directed, FUNGAFLOOR has curative and preventative action as well as anti-sporulation properties.

Dosage rates are given in weight/weight units.

In dips, wash tanks, and drenchers - Dilute with water at rates ranging from 1:1360 to 1:1020 to give treating concentrations of 500 to 750 ppm active ingredient. The treatment solutions should be inspected periodically to insure the proper concentration of active ingredient. Length of dip time should not exceed two minutes.

In non-recovery spray - add 1 part of FUNGAFLOOR to 680 parts of water to give a treating concentration of 1000 ppm active ingredient. Apply to freshly-cleaned citrus immediately prior to waxing.

For water spray brush application - after washing, dilute 1 part of FUNGAFLOOR to 680 parts of water to give a treating concentration of 1000 ppm active ingredient. Apply prior to storage wax application.

Waxer - Add 1 part of FUNGAFLOOR to 340 parts of diluted "ready to use" detergent material to give a treating concentration of 2000 ppm active ingredient. Apply to citrus fruit a mechanical framer. Use only as much as needed to clean fruit.

In Citrus wax - Add 1 part of FUNGAFLOOR to 340 parts of citrus wax to give a treating concentration of 2000 ppm active ingredient. Use without further dilution in conventional waxing equipment.

Remark 1: Combinations of above-listed treatments are possible, but never more than two treatments should be carried out on the same citrus lot.


Remark 2: In waxes and foams, sporulation control can be obtained at 4000 ppm active ingredient. Dilute 1 part of FUNGAFLOOR to 170 parts of citrus wax or diluted "ready to use" detergent material to give a treating concentration of 4000 ppm active ingredient. When this dilution is used, combinations of treatments are not allowed.

Remark 3: AFTER DILUTION OF FUNGAFLOOR IN PUMP OR WAX, STIR THOROUGHLY FOR AT LEAST 15 MINUTES WITH AN ELECTRIC MIXING DEVICE AT 1500 RPM.

**NOTICE OF WARRANTY**

Janssen warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as presence of other materials, or the manner of use or application, all of which are beyond the control of Janssen. In no case shall Janssen be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. JANSSEN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

BEST DOCUMENT AVAILABLE



Handwritten notes: 1st 2.5 10 43937-154-1