DANGER

Correctus. Causes irreversible eye damage. Maraful if availered, Do not get in eyes or on clothing. Year coegles when handling. Yesh thoroughly with some after handling.

INVIRORMENTAL MAZARDS

The nor apply directly to vater. Do not contaminate water by cleaning of equipment, or disposal of waster. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL BAZARDS

Do not use or store near heat or open flame.

FOR USE AND STORAGE BY AGRICULTURAL AND GERVICE PERSONS ONLY

DIRECTIONS FOR USE

It is a wielation of Faderal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Prohibitions - Do not contaminate water. food or faed by storage or disposal.

Storage - Store in a cool dry place.

Pesticide Disposal - Pesticide vastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or ripsate is a violation of Pederal Taw, If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest Regional Office for guidance.

container Disposal - Triple tinse for equivalent). Then offer for recycling or reconditioning, or nucture and dispose of in a senitary landfill, or inclineration. or. if allowed by state and local authorities, by burning, If burned, stay out of smoke.

JANSSEN PHARMACEUTICA

FECUNDAL 10EC

Active Ingredient
Imazalil: 1-(2-(2.4-dichlorophenyl)-2(2-propenyloxylethyl)-1X-imidazole**

Inert Ingredients 50 Contains 0.26 lbs. of Imagalil** Der gallon

10%

EFA keg. No. 43813-3 EFA Est. No. 43813-8L-2

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Induce vomiting, pump stomach, call a physician. Treat symptomatically.

If Inhiled: See a physician if an abnormal reaction occurs.

If on \$15; Yash thoroughly with soap and water.

If in Eyes: Immediately flush eyes with plenty of water. Call a physician

See side panel for additional precautionary statements.

Net Contents: 5 gallons



JAMSSEN PHARMACEUTICA 40 Kingsbridge Road Piscataway, NJ 08854 U.S.A.

ACCEPTED

JUL 9 1987

Under the Foderal Intenties to, Pungicide, and Rementing of Act, as attimented, for the postundo requested under EPA Reg. No. 43813-3

BEST AVARIABLE COPY

YMEAT & BARLEY: For control of common root rot (Weininthosporium satirum) and barley leaf stripe, use 0.8-1.5 fl. oz. FECUNDAL 10EC per 100 lbs of seed.

COTTOM: For control of Fuserium and Thieleviousis use 4-2 fl. oz. Fecundal 10EC per 100 lbs. of sead.

Use the higher dose when disease pressure is severe.

For ready mix treaters. Fecundal 102C may be applied directly to the seed. For slurry treaters, Fecundal 102C may be diluted with water.

SEED LABELLING: Seeds treated with Fecundal 10EC. Do not use treated seed for food, feed or oil purposes. Do not graze or feed foliage from treated area to livestock.

USE PRECAUTION: All seed treated with this product must be adequately dyed. Any dye added to treated seed must be cleared for use under 40 CFR. Part 180.1001.

NOTE TO USER: Combination treatments of FECUNAL 10 EC with products containing lindane may result in delayed emergence and stunting of wheat and barley, however. plants will usually outgrow this effect. Bo not carry over seed treated with 11 dans to the next crop season. Minimize time between treating and planting for such treatments.

NOTICE OF YARRANTY

Treatment of highly mechanically damaged seed, of known low vigor and poor quality, except for the Durbosss of curative control of existing disease or insect pests, can result in reduced dermination and/or reduction of seed and seadling vigor. Treat and dermination test a sample of the seed lot before treating with a selected chemical treatment. Excess seed should not be carried over from one season to the beat without re-testing germination. Due to seed quality conditions beyond the control of Janssen Pharmaceutics, no claims are made to quarantee dermination of carry-over seed.

Januaren Pharmaceutica warrents that this product conforms to the clemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury. ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the mapper of use or application, all of which are beyond the control of Janssen Pharmaceutics. In no case shall Janssen Pharmaceutics be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. JANESEN PRARMACTURICA MAKES NO VARRANTIES OF MERCHANTABILITY OR PITHESS FOR A PARTICULAR PURPOSE NOR MAY OTHER EXPRESS OR INPLIED VARBANTY EXCEPT AS STATED ABOVE.

10F/ Pmal 43813-3

1