

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing. Wear goggles when handling. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

FOR USE AND STORAGE BY AGRICULTURAL AND SERVICE PERSONS ONLY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Prohibitions - Do not contaminate water, food or feed by storage or disposal.

Storage - Store in a cool dry place.

Pesticide Disposal - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the nearest hazardous waste representative at the nearest Regional Office for guidance.

Container Disposal - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

JANSSEN PHARMACEUTICA

FECUNDAL[®] 10EC

Active Ingredient
Imazalil: 1-(2-(2,4-dichlorophenyl)-2-(2-propanoyloxy)ethyl)-1H-imidazole** 10%

Inert Ingredients 90%
Contains 0.85 lbs. of Imazalil** per gallon

EPA Reg. No. 43611-J
EPA Est. No. 43813-8L-2

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Induce vomiting, pump stomach, call a physician. Treat symptomatically.

If Inhaled: See a physician if an abnormal reaction occurs.

If on Skin: Wash thoroughly with soap and water.

If in Eyes: Immediately flush eyes with plenty of water. Call a physician

See side panel for additional precautionary statements.

Net Contents: 5 gallons

JANSSEN PHARMACEUTICA
40 Kingsbridge Road
Piscataway, NJ 08854
U.S.A.

ACCEPTED
DEC 17 1987
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
this pesticide
has been found to meet the
requirements of the Act.
EPA Reg. No. 43813-3

BEST AVAILABLE COPY

WHEAT & BARLEY: For control of barley leaf stripe, seedborne net blotch and *Septoria nodorum*, and protection against the seed and soilborne phases of the common root rot disease complex (*Helminthosporium*, *Fusarium*). Promotes stand establishment by protecting against seed and seedling blights caused by *Helminthosporium*, *Fusarium*, or *Penicillium*. Use 0.8 to 1.5 fl. oz. FECUNDAL 10EC per 100 lbs. seed.

COTTON: For control of *Fusarium* and *Thielaviopsis* use 4-8 fl. oz. FECUNDAL 10EC per 100 lbs. of seed.

Use the higher dose when disease pressure is severe.

For ready mix treaters, FECUNDAL 10EC may be applied directly to the seed. For slurry treaters, FECUNDAL 10EC may be diluted with water.

SEED LABELLING: Seeds treated with FECUNDAL 10EC. Do not use treated seed for food, feed or oil purposes. Do not graze or feed foliage from treated area to livestock.

USE PRECAUTION: All seed treated with this product must be adequately dyed. Any dye added to treated seed must be cleared for use under 40 CFR, Part 180.1001.

NOTE TO USER: Combination treatments of FECUNDAL 10 EC with products containing lindane may result in delayed emergence and stunting of wheat and barley, however, plants will usually outgrow this effect. Do not carry over seed treated with lindane to the next crop season. Minimize time between treating and planting for such treatments.

NOTICE OF WARRANTY

Treatment of highly mechanically damaged seed, of known low vigor and poor quality, except for the purposes of curative control of existing disease or insect pests, can result in reduced germination and/or reduction of seed and seedling vigor. Treat and germination test a sample of the seed lot before treating with a selected chemical treatment. Excess seed should not be carried over from one season to the next without re-testing germination. Due to seed quality conditions beyond the control of Janssen Pharmaceuticals, no claims are made to guarantee germination of carry-over seed.

Janssen Pharmaceuticals warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Janssen Pharmaceuticals. In no case shall Janssen Pharmaceuticals be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. JANSSEN PHARMACEUTICA MAKES NO WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

10F1
pm21
43813-3