

Fungaflor 6.5L** SEED TREATMENT

PRECAUTIONARY STATEMENTS:

Hazards to Humans and Domestic Animals

WARNING

Causes eye irritation. Wear goggles when handling the concentrate. May be fatal if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin, or on clothing.

ENVIRONMENTAL HAZARDS: Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

FOR USE AND STORAGE BY AGRICULTURAL AND SERVICE PERSONS ONLY

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal.

STORAGE

Store in a cool dry place.

PESTICIDE DISPOSAL - Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Trademark of Janssen Pharmaceutica, B 2340 Beerse, Belgium

ACTIVE INGREDIENT:

Imazalil: (1-(2-(2,4-dichlorophenyl)-2-(2-propenyloxy)ethyl)-1H-imidazole)*.....6.58

INERT INGREDIENTS:.....93.58

Contains 0.48 lbs. of Imazalil* per gallon. *Active ingredient is manufactured by Janssen Pharmaceutica n.v., Beerse, Belgium

EPA Reg. No.
EPA Est. No. R43813-BL02
U.S. Patent No. 3,658,813

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT
If swallowed: induce vomiting, pump stomach, call a physician. Treat symptomatically.

If inhaled: See a physician if an abnormal reaction occurs.

If on Skin: Wash thoroughly with soap and water.

If in eyes: Immediately flush eyes with plenty of water. Call a physician.

See side panel for additional precautionary statements.

Net Contents: 5 gallons



Janssen Pharmaceutica
40 Kingsbridge Road
Piscataway, N.J. 08854
U.S.A.

WHEAT & BARLEY: For control of common root rot (*Helminthosporium sativum*) and barley leaf stripe, use 1.0 to 1.5 fl. oz. of Fungaflor 6.5L per bushel of seed.

COTTON: For control of *Fusarium* and *Thielaviopsis* use 7.5-15 fl. oz. of Fungaflor 6.5L per 100 lbs. of seed.

For ready mix treaters, Fungaflor 6.5L may be applied directly to the seed.

For slurry treaters, Fungaflor 6.5L may be diluted with ethanol or a 20:80 water: ethanol solution not to exceed 6 fl. oz. additional liquid per 100 lbs. of wheat and barley seed or 15 fl. oz. per 100 lbs. of cottonseed.

SEED LABELING: Seeds treated with FUNGAFLO 6.5L. Do Not use treated seed for food, feed or oil purposes. Do not graze or feed foliage from treated area to livestock.

USE PRECAUTION - all seed treated with this product must be adequately dyed. Any dye added to treated seed must be cleared for use under 40 CFR, Part 180.1001.

NOTICE OF WARRANTY

Janssen Pharmaceutica warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Janssen Pharmaceutica. In no case shall Janssen Pharmaceutica be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. JANSSEN PHARMACEUTICA MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

BEST DOCUMENT AVAILABLE

