

PRECAUTIONARY STATEMENTS:
HAZARDS TO HUMANS AND DOMESTIC
ANIMALS

WARNING

Causes eye irritation. Wear goggles when handling the concentrate. May be fatal if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin, or on clothes.

ENVIRONMENTAL HAZARDS: Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS:
Flammable. Keep away from heat and open flame.

FOR USE AND STORAGE BY AGRICULTURAL AND SERVICE PERSONS ONLY

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS - Do not contaminate water, food or feed or storage or disposal.

STORAGE - Store in a cool dry place.

PESTICIDE DISPOSAL - wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration if it allows or state and local authorities, or burning, if burned, stay out of smoke.

*Active ingredient is manufactured by Janssen Pharmaceutica, Beerse, Belgium

JANSSEN PHARMACEUTICA

Fecundai® 6.5L SEED TREATMENT

Active Ingredient:

Imazalil-(1-(2-(2,4-dichlorophenyl)-2-(2-propenyl)oxyethyl)-1H-imidazole) 6.5%

Inert Ingredients: 93.5%

Contains 0.48 lbs. of Imazalil* per gallon

EPA Reg. No. 43813-5
EPA Est. No. 43813-06-2

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Induce vomiting, pump stomach call a physician, treat symptomatically.

If inhaled: See a physician if an abnormal reaction occurs.

If on skin: Wash thoroughly with soap and water.

If in eyes: Immediately flush eyes with plenty of water. Call a physician.

See side panel for additional precautionary statements.

Net Contents: 3 gallons



JANSSEN PHARMACEUTICA
40 Kingsbridge Road
Piscataway, NJ 08854
U.S.A.

ACCEPTED
AUG 24 1987
This is the Federal Insecticide, Fungicide, and Plant Quarantine Act, 7 U.S.C. 136, for the purpose of registration of under EPA Reg. No. 43813-5

BEST AVAILABLE COPY

43813-5 PM 21 1061

fl. oz. per 100 lbs. of seed. Fecundai 6.5L controls barley leaf stripe and suppresses common root rot (Cochliobolus sativus and Fusarium spp.), Septoria leaf spot, spot blotch (C. sativus), tan spot (Pyrenopeziza trichostoma), and net blotch sp. cereal by controlling seedborne and seedling infections.

COTTON: For control of Fusarium and Thielaviopsis use 7.5-15 fl. oz. of Fecundai 6.5L per 100 lbs. of seed.

Use the higher dose when disease pressure is severe.

For ready mix treaters, Fecundai 6.5L may be applied directly to the seed. For slurry treaters, Fecundai 6.5L may be diluted with ethanol or a 20:40 water: ethanol solution.

SEED LABELING: Seeds treated with Fecundai 6.5L. Do not use treated seed for food, feed or oil purposes. Do not graze or feed foliage from treated area to livestock.

USE PRECAUTION: All seed treated with this product must be adequately dried. Any dye added to treated seed must be cleared for use under 40 CFR, Part 160.1001.

NOTICE OF WARRANTY

Treatment of highly mechanically damaged seed, of known low vigor and poor quality, except for the purposes of curative control of existing disease or insect pests, can result in reduced germination and/or reduction of seed and seedling vigor. Treat and germination test a sample of the seed lot before treating with a selected chemical treatment. Excess seed should not be carried over from one season to the next without retesting germination. Due to seed quality conditions beyond the control of Janssen Pharmaceutica, no claims are made to guarantee germination of carry-over seed.

Janssen Pharmaceutica warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Janssen Pharmaceutica. In no case shall Janssen Pharmaceutica be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. JANSSEN PHARMACEUTICA MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.