

**PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating. Remove contaminated clothing and wash before reuse. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on the label. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permit authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your local State Water Board or Regional Office of the EPA.

STORAGE AND DISPOSAL

Store in original container in a cool, dry place. Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by other procedures approved by State and Local authorities.

*Trademark of Janssen Pharmaceutica.

*Active ingredient and formulation is manufactured by Janssen Pharmaceutica, Beerse, Belgium

Janssen Pharmaceutica

WOCOSEN™ 100SL
Wood Preservative

FOR INDUSTRIAL & COMMERCIAL USE ONLY

ACTIVE INGREDIENT: (w/w)

PROPICONAZOLE*:

1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl(methyl)-1H-1,2,4-triazole] 9.7%

INERT INGREDIENTS: (w/w) 90.3%

Contains 0.83 lbs. active ingredient/gallon

KEEP OUT OF REACH OF CHILDREN

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching in back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Treat symptomatically. Get medical attention.

IF IN EYES: Hold eyelids open and flush immediately with a steady gentle stream of water for at least 15 minutes. Get medical attention if irritation persists.

IF ON SKIN: Immediately flush exposed area with plenty of water. Remove contaminated clothing. Wash thoroughly with soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

NET CONTENTS: 55 gallons (208 liters)

EPA REG. NO. 43813-19

EPA EST. NO. 43813-BL-2



JANSSEN
Pharmaceutica
1125 Trenton-Harbourton Rd.
Titusville, NJ 08560-0200
(809) 730-2607

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Description: WOCOSEN 100SL is a wood preservative concentrate which requires dilution with water prior to use on wood products to protect against decay (brown and white rot) and staining fungi (blue stain).

Product Uses: For application to millwork, shingles & shakes, siding, plywood, and structural lumber & composites that are used in above ground applications only. Do not use on wood which will come in contact with food.

Wood Preparation: Machine, cut, glue, and sand wood as desired before treating. Do not use over previously finished or stained wood without stripping off old finish. Wood must be clean and dry before treatment. Wood for pressure treatment should be kiln dried or air seasoned to a moisture content, of about 25% or less, that will permit adequate penetration of preservative during treatment.

Treatment: Apply by immersion, roller coater, flood, spray, brush or pressure treatment processes (double vacuum, full-cell or modified full-cell). For non-pressure application methods, mix 5-10 gallons of Wocosen 100SL with water to achieve 100 gallons of treating solution at the concentration of 0.5 - 1% a.i., respectively. Wood products to be treated by the bundle-dipping process should be completely immersed for three (3) minutes. Individual wood pieces treated by immersion, roller-coater or flood shall be treated for not less than fifteen (15) seconds. Coverage for brush and spray application is approximately 150 square feet per gallon of treating solution, but varies by the species and condition of the wood. Pressure treatment should be performed according to the methods described in the American Wood Preservers' Association (AWPA) Standards. A minimum propiconazole retention level of 0.025 pounds per cubic foot (pcf) is recommended. All treated wood must be held on the drip pad until drippage has stopped. Any surface, cut after first treatment should be retreated using 1% a.i. concentration solution.

Finishing: Before glazing, staining, painting, varnishing, or application of water repellents, treated wood should be completely dry and odor-free. For non-pressure applications, allow 4-8 hours drying time under average temperature and humidity conditions; treated wood can be heat dried for plant production. Pressure treated wood requires drying after treatment to a moisture content suitable for the type of coating to be applied.

PM 21

43813-19

Page 1 of 2

282

Clean-up: Use water to remove wet WOCOSEN 100SL wood preservative from all surfaces, brushes and rollers.

Coatings, Caulks, Sealants, Adhesives, and Inks: Sufficient mixing to ensure adequate dispersion of the biocide is required. WOCOSEN 100SL should be used at levels between 0.5 and 12.5% based on the total weight of the formulation to inhibit fungal disfigurement and deterioration. For solvent-based formulations the WOCOSEN 100SL can be dissolved in aromatic solvents or combination of aromatic and aliphatic solvents and added in the letdown or added directly to the finished product. For water-thinned latex emulsion formulation, the WOCOSEN 100SL can be premixed with the wetting agent and added to the pigment slurry or simply added to the letdown or finished product.

NOTICE OF WARRANTY

Janssen Pharmaceutica warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Janssen Pharmaceutica. In no case shall Janssen Pharmaceutica be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. JANSSEN PHARMACEUTICA MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE

97 JAN -7 P 3 :10

REC'D

7/0P01